



Panafrican colloquium/Colloque panafricain (Dakar 2012)

Health-risk market garden production linked to heavy metals in irrigation water in Benin

Production maraîchère à risques liés aux métaux toxiques dans l'eau d'irrigation au Bénin

Luc Koumolou^{a,*}, Patrick Etorh^a, Sabine Montcho^a, Kodjo Aklirikou^b, Frédéric Loko^c, Michel Boko^d, Edmond E. Creppy^e

^a Laboratoire de toxicologie et de santé environnementale, université d'Abomey-Calavi, 03 BP 1463, Cotonou, Benin

^b Département de physiologie/pharmacologie, faculté des sciences, université de Lomé, BP 1515 Lomé, Togo

^c Direction des laboratoires et des pharmacies, ministère de la Santé, 01 BP 882 Cotonou, Benin

^d CIFRED, université d'Abomey-Calavi, 03 BP 1463, Cotonou, Benin

^e Laboratoire de toxicologie et hygiène appliquée/UFR des sciences pharmaceutiques, université de Bordeaux-2, 146, rue Léo-Saignat, 33076 Bordeaux cedex, France

ARTICLE INFO

Article history:

Available online 30 May 2013

Keywords:

Heavy metals
Irrigation water
Market production
Health risk
Benin

Mots clés:

Métaux lourds
Eau d'arrosage
Production maraîchère
Risques sanitaires
Bénin

ABSTRACT

Heavy metals in the Benin market garden products: is irrigation water the first factor in question, and what is the level of health risk linked to the consumption of these vegetables? Such are the essential problems that this survey attempts to solve. Comparison of the level of lead (Pb), cadmium (Cd) and arsenic (As) pollution shows that all the vegetables taken from three market sites are differently contaminated, as well as their irrigation water and the soil. But establishing that water is the first factor responsible for the presence of heavy metals in market garden products is not so obvious. Otherwise, the health risk assessment revealed that the total daily exposure dose (DED) of Cd, namely 8.05 $\mu\text{g}/\text{kg}/\text{day}$, is high compared to the daily dose defined by the WHO, which is 1 $\mu\text{g}/\text{kg}/\text{day}$. Also, the ensuing quotient of danger (QD) is 8.05; such a value poses public health risks for the consumer.

© 2013 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

RÉSUMÉ

La présence de métaux lourds dans les produits maraîchers du Bénin est-elle imputable à la qualité de l'eau d'irrigation et quel est le niveau d'exposition du consommateur de ces légumes à des risques sanitaires ? Telles sont les questions essentielles auxquelles cette étude tente de répondre. La comparaison des niveaux de pollution en plomb (Pb), en cadmium (Cd) et en arsenic (As) montre que tous les légumes prélevés sur trois sites maraîchers sont différemment contaminés, ainsi que les eaux d'irrigation et les sols. Établir, toutefois, que l'eau d'arrosage soit la cause principale de la contamination des légumes n'est pas aussi évident. Par ailleurs, une évaluation des risques liés au Cd a permis de calculer une dose journalière d'exposition totale (DJE) égale à 8,05 $\mu\text{g}/\text{kg}$ par jour, à comparer à la limite définie par l'OMS, qui est de 1 $\mu\text{g}/\text{kg}$ par jour. Aussi, le quotient de danger (QD) s'élève-t-il à 8,05, ce qui génère des risques de santé publique pour le consommateur.

© 2013 Académie des sciences. Publié par Elsevier Masson SAS. Tous droits réservés.

* Corresponding author.

E-mail address: heraluc@yahoo.fr (L. Koumolou).

1. Introduction

With the economic crisis in Benin, there has been an increase in the number of gardens as well as in the number of cultivated urban areas [1]. But the poor farming practice regulations pose also a threat to the health safety of garden products [2]. In fact, in Benin, market constraints, land and parasitic pressures force farmers to practice intensive phytosanitary treatments. It is well known that some farming practices are responsible for the introduction of traces of metal in gardening soils. Products intended to improve the physicochemical properties of soils are often richer in heavy metals than the soil itself [3]. Among them, we can cite fungicides [4], fertilizers, and composts [5]. Otherwise, water for market garden irrigation is a limiting factor for gardening, and many gardens are compelled to remain in the vicinity of swamps, areas that are used for dumping and are therefore sources of water as well as soil and vegetable contamination by xenobiotics [2]. This study attempts to establish that, among soils, use of phytosanitary products, location of the gardening site (rural or urban), irrigation water quality is the first factor responsible for the contamination of market garden products by heavy metals. In this study, we also make an assessment of health risks associated with the consumption of vegetables contaminated by toxic metals in the watering water.

2. Materials and methods

Two sites in Cotonou and another in Aplahoué, about 150 km far from Cotonou, have been our study framework (Fig. 1).

On two of the sites located in downtown Cotonou, Houéyiho (H) and Godomey (G), irrigation water comes from neighbouring swamps, and on the Aplahoué site (A), it comes from a small river streaming through cotton and crop fields.

2.1. Processing and analysis

Composite samples of eight different vegetables *Amaranthus hybridus* (amaranth), *Daucus carota* (carrot), *Lactuca sativa* (lettuce), *Spinacia oleracea* (spinach), *Allium cepa* (green onion), *Brassica oleracea* (cabbage), *Corchorus olitorius* (fiddle), *Salanum macrocarpum* (nightshade) were collected and treated with or without phytosanitary products, whereas water and soil samples were collected during the same periods on three major gardening sites in Benin.

The numbers of soils, water and vegetables samples are presented in Table 1.

The samples underwent the necessary treatment before they are tested for lead (Pb), cadmium (Cd) and arsenic (As) by atomic absorption spectrophotometry. First, all samples are cleaned and stored initially in the oven, first at 50 °C for




Fig. 1. Map of the study area (Cotonou [Houéyiho + Godomey] and Aplahoué).

Table 1
Numbers of soils, water and vegetables samples.

Sites	Godomey	Houéyiho	Aplahoué
Number of soils samples	7	7	7
Number of water samples	7	7	7
Number of vegetable samples	8	8	8

12 hours, then the temperature is increased to 120 °C and maintained for 24 hours. But water samples are left at 4 °C. After grinding in a mortar, 2 g of each lyophilized sample is soaked in a mixture of hydrogen peroxide (H₂O₂) and nitric acid for 24 hours, and then undergo an acid digestion in a digester (sand bath at 125 °C for 2 hours). The digest was recovered in a 100-ml flask and completed with distilled water to the mark. It is then filtered and stored at 4 °C until analysis for the extraction of heavy metals by atomic absorption spectrophotometry (Thermo Orion corrected Solaar S2), in the Laboratory of Management, Treatment and Recovery of Wastes of the University of Lomé in Togo according to the method described in [6] and [7] by electro-thermal atomic absorption spectrophotometry. The analysis was performed by interpolation on calibration

curves obtained with standard solutions. The results, displayed by software Solaar S2 on the computer screen connected to the spectrophotometer, are expressed in mg/kg or mg/l or ppm. The corrected concentration is obtained by the formula:

$$\text{Final concentration} = \frac{\text{Concentration on the screen} \times 100}{2}$$

The health risk assessment of Cd was performed using the standardized approach proposed by [8]. DED, the daily exposure dose, is expressed in µg/kg/day and calculated by crossing data of consumption of food contaminated with metal by the averaged poison content measured in the sample (Table 5). For the Cd-contaminated *Amaranthus hybridus* consumed by people, we have:

$$\text{DED} = \frac{Q_a \times C_a}{\text{BW}}$$

$$\text{QD} = \frac{\text{DED}_{\text{total}}}{\text{DAD}}$$

QD is the quotient of danger.

Table 2
Contamination of vegetables by toxic metals at the three sites.

Sites	Lead (ppm)			Cadmium (ppm)			Arsenic (ppm)		
	Norms	0.3 [9]			0.05 [10]			0.1 [10]	
	G	H	A	G	H	A	G	H	A
<i>L. sativa</i>	4.84	3.10	6.03	0.89	0.63	0.07	237.62	260.48	315.13
<i>A. hybridus</i>	5.01	3.73	5.92	0.93	5.13	0.32	171.63	259.92	302.15
<i>S. macrocarpum</i>	2.52	3.38	5.99	0.82	0.64	0.81	282.73	316.28	219.19
<i>B. oleracea</i>	6.69	3.12	6.32	1.73	0.55	0.79	230.81	325.30	195.08
<i>D. carota</i>	1.06	1.14	2.37	1.22	0.72	0.53	300.51	251.45	270.18
<i>C. olitorius</i>	3.5	4.76	5.52	0.91	1.17	1.10	271.76	241.41	255.12
<i>A. cepa</i>	2.85	3.36	4.80	0.43	0.26	0.34	358.67	323.52	196.4
<i>S. oleracea</i>	4.08	3.46	4.95	0.34	0.52	0.75	231.84	204.01	232.02
Average	3.82	3.25	5.24	0.91	1.20	0.59	260.70	272.80	248.16
± SD	± 1.73	± 1.00a	± 0.88b	± 0.43	± 1.60	± 0.33	± 56.04	± 44.25	± 45.56
C. factors	12.73	10.86	17.46	18.2	24	11.8	2607	2728	2481.6

G, H and A are respectively the initials for Godomey, Houéyiho and Aplahoué. SD: Standard Deviation; *t*-test: numbers (values) followed by letters are significantly different, others are not for the same metal. $P < 0.05$ for Pb between Houéyiho and Dokomey (they are significantly different for the Pb); for the rest, $P > 0.05$. C. factors or contamination factor (CF) is the ratio between heavy metal concentration and its concentration limits. $\text{CF} = [\text{Vegetable}]/\text{Norm}$.

Table 3
Soil contamination by toxic metals at the three sites.

Sites	Lead (ppm)			Cadmium (ppm)			Arsenic (ppm)		
	Norms [11] 85	0.80			29				
	G	H	A	G	H	A	G	H	A
Soils	9.527	49.75	4.62	0.74	0.09	0.63	151.2	138.79	112.34
	9.956	47.92	4.62	0.81	0.04	0.72	165.1	142.58	107.43
	11.24	50.72	5.02	0.74	0.03	0.56	158.8	151.33	115.55
	10.12	50.20	5.10	0.83	0.08	0.93	162.0	145.94	104.76
	9.75	48.51	4.92	0.75	0.04	0.89	160.8	139.84	92.69
	10.81	51.67	4.72	0.77	0.03	1.10	153.3	137.91	92.22
Average	9.90	49.10	4.91	0.64	0.04	0.68	163.2	153.34	115.28
	10.18	49.7	4.84	0.75	0.05	0.78	159.20	144.24	105.75
± S D	± 0.61 a	± 1.29b	± 0.19a	± 0.06a	± 0.02b	± 0.19a	± 5.16a	± 6.16 a	± 9.89 a
C. factors	0.11	0.58	0.05	0.98	0.06	0.98	5.48	4.97	3.64

t-test: means for identical letter for the same metal are not significantly different; those with different letters for the same metal are significantly different. So $P < 0.05$ for Cd and Pb between Houéyiho and the other two sites; for the rest, $P > 0.05$. $\text{CF} = [\text{Soils}]/\text{Norm}$.

Table 4
Contamination of irrigation water by toxic metals at the three sites.

Sites	Lead (ppm)			Cadmium (ppm)			Arsenic (ppm)		
	Norms [12] 0.4			0.21			0.1		
	G	H	A	G	H	A	G	H	A
Water	4.81	2.85	2.24	1.65	0.79	0.92	583.11	335.41	281.02
	5.07	3.51	3.10	1.45	0.82	0.62	592.82	344.52	280.51
	3.97	3.52	2.09	1.43	0.79	0.74	571.81	353.12	269.61
	4.96	3.45	2.51	1.16	0.92	0.90	542.51	343.94	282.41
	5.41	3.52	3.50	1.80	0.72	0.83	502.41	342.14	265.71
	5.71	3.54	2.81	1.62	0.83	0.65	573.71	336.54	275.91
Average	5.06	3.39	2.66	1.53	0.83	0.78	562.75	344.54	277.29
± SD	± 0.58b	± 0.24a	± 0.50a	± 0.20b	± 0.09a	± 0.11a	± 30.75b	± 7.76a	± 7.29a
C. factors	12.67	8.47	6.67	7.33	4	37.6	5626.6	3445.5	2773.0

t-test: the values for identical letters, for the same metal are not significantly different; those with different letters for the same metal are significantly different. $P < 0.05$ for Pb, Cd and As between Godomey and each one of the other two sites; for the rest, $P > 0.05$. CF = [Water]/Norm.

Statistical treatment is used for a pairwise and one-against-all comparison, using Student's *t*-test $P(T > t) = 0.05$.

3. Results

The results are given in ppm (mg/kg) and are compared in Tables 2–4 as mean ± standard deviation, and in Figs. 2–5

showed the presence of toxic metals in irrigation water, soils and vegetables on the three sites with some significant differences between the results. In addition, the risk assessment is summarized in Table 5 and a daily exposure dose of 8.05 µg/kg/day is shown, whereas the amount suggested is 1 µg/kg/day. The hazard quotient or quotient of danger (QD) therefore is shown to be 8.05.




Fig. 2. Vegetable contamination by site.




Fig. 3. Comparison between vegetable and soil contamination.




Fig. 4. Comparison between vegetable (légumes) and water (eaux) contamination.




Fig. 5. Contamination of pesticide-treated vegetables (T) compared to that of untreated ones (NT).

Table 5
Toxicological data about cadmium.

Q_a	C_a	BW	DAD [13]	DED	ADC [14]	DED total	QD
0.1	5.13	65	1	7.89	0.16	8.05	8.05

DED: daily exposure dose for cadmium in $\mu\text{g}/\text{kg}/\text{day}$; Q_a : average quantity of *Amaranthus* consumed in kg; C_a : average cadmium concentration measured in *Amaranthus* in $\mu\text{g}/\text{kg}$; BW: body weight of the consumer (adult) in kg; DAD: daily allowed dose (DAD) $\mu\text{g}/\text{kg}/\text{day}$ of cadmium or tolerable; ADC: average daily contributions by other contaminated foodstuffs.

4. Discussion

An analysis of the results revealed that multiple sources of contamination by toxic pollutants of the gardening soil and water from swamps and the river crossing agricultural fields affect the sanitary quality of the grown vegetables (Table 2). Comparison of the contamination level of all vegetables on the gardening sites showed no significant difference ($P > 0.05$) from one site to the other (Fig. 2). But apart from lead, the lowest levels were recorded on the rural site of Aplahoué (Fig. 2). We might think immediately that the polluted atmosphere of the city contaminates the vegetables the most. The results also showed that the irrigation water for irrigation of the vegetables as well as the soil on which they are grown is contaminated by toxic metals (Tables 3 and 4). By comparing the levels of metals in vegetables and in the soil (Fig. 3), we notice that an arsenic and cadmium bioaccumulation phenomenon has taken place (their contents in vegetables exceeded their level in the soil). This is not the case with lead levels in soil, which exceed those in vegetables. In fact, Cd and As are highly soluble and thus are more bioavailable to plants [15]. In this case, their concentrations in the soil solution or irrigation water are not significantly different ($P > 0.05$) from those found in the vegetables [16] because these plants grow faster and are quickly harvested. Apart from the only case of As level on the Godomey site, this hypothesis is verified (Fig. 4). It is not the case with lead because it is less soluble [17]. If it is found abundantly in vegetables; it probably comes from different origins. Mench et al. [18] believe that the Pb found in vegetables is especially atmospheric; this is the reason why smaller amounts were found in samples from the rural site in Aplahoué and these samples are apparently less polluted. Leafy vegetables will be more exposed than non-leafy ones: lead levels in *Daucus carota* and in *Allium cepa*, non-leafy vegetables, are among the lowest; this also proves this hypothesis (Table 2). However, this result does not allow us to conclude about the origin of vegetable contamination. As for the soil, the concentrations of toxic metals appear to be like ancillary data for plants harvested early, because the comparison between vegetables and soils (Fig. 3) showed significant differences ($P < 0.05$). Therefore, the contamination of the vegetables does not come only from the ground. On the contrary, the levels of toxic metals in irrigation water seem to influence the contamination of the vegetables for two over three sites; when we compare vegetables and water (Fig. 4), there is no

significant difference ($P > 0.05$) between the levels in water of Cd on the one hand and of As the other. This is justified because the ionic forms of these two metals are very soluble in the soil solution or the irrigation water that can be bioavailable for vegetables. An analysis of the soil has shown that its level of toxic metals is incidental. However, some results contradicted this argument, which lets us assume that the contamination of vegetables does not depend exclusively on the toxic levels in the soil or in the irrigation water. The contamination of *S. macrocarpum* whether pesticides are used or not (Fig. 4) showed a significant difference ($p < 0.05$) increase for lead and arsenic in treated vegetables (T) compared to untreated ones (NT). Therefore, the contamination of vegetables depends on the use of pesticides. Studies have highlighted other factors involved in the bioaccumulation of toxic metals by living organisms: metal speciation, the intrinsic nature of the organism bioaccumulative, biotic and physicochemical [19]. Finally, the content of Cd in vegetable (5.13 mg/kg) was used to calculate (Table 5) the daily exposure dose (EDI), taking into account the total exposure and the hazard quotient (QD). It appeared that the consumption of 100 g of this vegetable exposes an adult of 65 kg to 523 μg of Cd per day, when he only accepts a maximum of 65 μg . So, there is an annual accumulation of more than 167 g of Cd in the body of the consumer, hence the chronic risk whose magnitude is, for the moment, unknown. The QD greater than 1 confirms this observation.

5. Conclusion

This study showed that a contamination of vegetables by toxic metals depends on the soil, water, the environment of the gardening site and the use or not of pesticides. Many authors have highlighted that many factors – such as water – are primary responsible for the contamination of vegetables by pollutants. Our results prove this observation. In any case, due to the imminent risk of toxicity, the adoption of reasonable behaviour and the development of sustainable agriculture are needed to combine food security, economic development, environmental protection and public health.

Disclosure of interest

The authors declare that they have no conflicts of interest concerning this article.

References

- [1] F. Assogba-Komlan, P. Anihouvi, E. Achigan, R. Sikirou, A. Boko, C. Adje, V. Ahle, R. Vodouhe, A. Assa, Pratiques culturales et teneur en éléments anti nutritionnels (nitrates et pesticides) du *Solanum macrocarpum* au sud du Bénin, Afr. J. Food. Agric. Nutr. Dev. 7 (4) (2007) [21 p.].
- [2] P. Jacobi, J. Amend, S. Kiango, Urban agriculture in Dar es Salaam: providing an indispensable part of the diet, in : N. Bakker, et al. (Eds.), Growing cities, growing food: urban agriculture on the policy agenda, a reader on urban agriculture, ETC/DSE, Feldafing, Germany, 2000, pp. 257–283.
- [3] B.J. Alloway, Heavy metals in soils, 2nd ed., Blackie Academic & Professional, Glasgow, UK, 1995, [368 p.].

- [4] A. Deluisa, P. Giandon, M. Aichner, P. Bortolami, L. Bruna, A. Lupetti, F. Nardelli, G. Stringari, Copper pollution in Italian vineyard soils, *Commun. Soil. Sci. Plant. Anal.* 27 (1996) 1537–1548.
- [5] M. Robert, C. Juste, Enjeux environnementaux et industriels. Dynamique des éléments traces dans l'écosystème sol, Spéciation des métaux dans le sol, *Les Cahiers du Club Crin*, Paris, 1999 15–37.
- [6] R. Anane, M. Bonini, M.J. Grafeille, E.E. Creppy, Bioaccumulation of water soluble aluminium chloride in the hippocampus after transdermal uptake in mice, *Arch. Toxicol.* 69 (1995) 568–571.
- [7] C.O. Vaidya, T.T.R. Rantala, A comparative study of analytical methods determination of heavy metal in mussels (*Mytilus edulis*) from eastern Canada, *Int. Environ. Anal. Chem.* 63 (1996) 179–185.
- [8] C. Ricoux, B. Gasztowitz, Évaluation des risques sanitaires liés à l'exposition de forts consommateurs de produits de la pêche de rivière contaminés par des toxiques de l'environnement, *La Documentation française*, 1995 56 p.
- [9] INERIS, Données toxicologiques et environnementales des substances chimiques, Plomb et ses dérivés, Fiches INERIS, version N° 3, 2006.
- [10] Règlement CE/466/2001, Fixation de teneurs maximales pour certains contaminants dans les denrées alimentaires, Union européenne, 2001.
- [11] Smit H., Pollution prevention policies for agricultural soils in the Netherlands. Proc IIIrd Int. Conf. Biogeochem. Traces metals, Paris In: R. Prost (Ed.), Contaminated soils, « les Colloques » 85, INRA Éditions, Paris, 1998.
- [12] GESAMP, Group of experts on the scientific aspects of marine pollution, the health of the oceans, *Rep. Stud.* 15 (1982) 1–108.
- [13] WHO/Joint FAO/Expert Committee on Food Additives, Evaluation of certain food additives and contaminants: twenty-ninth report of the Joint FAO/WHO Expert Committee on Food Additives [meeting held in Geneva from 3 to 12 June 1985]. Food and Agriculture Organization of the United Nations. World Health Organization, 1986.
- [14] Agence française de sécurité sanitaire des aliments (AFSSA), Institut de veille sanitaire, Le méthylmercure, Programme « mortalité et morbidité des maladies d'origine alimentaire » Volet toxicologique-Contaminants chimiques et risque alimentaire en France, document de travail, 2003.
- [15] V.J. Camobreco, B.K. Richards, T.S. Steenhuis, J.H. Peverly, M.B. McBride, Movement of heavy metals through undisturbed and homogenized soil columns, *Soil Sci.* 161 (1996) 740–750.
- [16] A. Tessier, D.R. Turner, Metal Speciation and Bioavailability in Aquatic Systems, 3, John Wiley & Sons, Chichester, 1995, p. 609–646.
- [17] Ablain F., Rôle des activités lombriciennes sur la redistribution des éléments traces métalliques issus de boue de station d'épuration dans un sol agricole, thèse de doctorat, université de Rennes-1, UMR ECOBIO, Paris, 2002, 256 p.
- [18] M. Mench, V. Amans, D. Arrouays, V. Didier-Sappin, S. Fargues, A. Gomez, M. Löffler, P. Masson, B. Spiteri, I. Weissenhorn, In situ remediation of soils contaminated by Pb fallout (risk assessment near a lead tetra-ethyl production facility, and effectiveness of several inorganic compounds for decreasing Pb mobility and plant availability), *Env-srae*, 1993 61 p..
- [19] S.N. Luoma, Bioavailability of trace metals to aquatic organisms – a review, *Sci. Total Environ.* 28 (1983) 1–22.