



Evolution/Évolution

Alkaloid secretion inhibited by antibiotics in *Aphaenogaster* ants

La sécrétion d'alcaloïdes inhibée par un traitement antibiotique chez les fourmis Aphaenogaster

Alain Lenoir*, Séverine Devers

IRBI, Institut de recherche sur la biologie de l'insecte, Université de Tours, Faculté des sciences, parc de Grandmont, 37200 Tours, France

ARTICLE INFO

Article history:

Received 16 April 2018

Accepted after revision 12 June 2018

Available online 30 June 2018

Keywords:

Aphaenogaster

Poison gland

Dufour gland

Alkaloids

Cuticular hydrocarbons

Bacteria

Mots clés :

Aphaenogaster

Glande à poison

glande de Dufour

Alcaloïdes

Hydrocarbures cuticulaires

Bactéries

ABSTRACT

Although alkaloids are frequent in the poison glands of ants of the genus *Aphaenogaster*, this is not the case for *A. iberica*. Hypothesizing that in the genus *Aphaenogaster*, alkaloids are produced by symbiotic bacteria, except for *A. iberica*, we treated an experimental lot of both *A. iberica* and a 'classical' *Aphaenogaster* species, *A. senilis*, with an antibiotic. Compared to workers from a control lot, this treatment reduced considerably alkaloid production in *A. senilis*, whereas *A. iberica* did not react to the treatment. Furthermore, the treatment induced an increase in cuticular hydrocarbon quantities in *A. senilis*, but not in *A. iberica*. An analysis of the ant microbiota will be the next step to confirm our hypothesis. © 2018 Published by Elsevier Masson SAS on behalf of Académie des sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

R É S U M É

On trouve de nombreux alcaloïdes dans les glandes des fourmis, particulièrement chez les *Aphaenogaster*, mais pas chez *A. iberica*, qui fait exception. Nous avons émis l'hypothèse que la production d'alcaloïdes pouvait être liée à l'activité de bactéries qui seraient absentes chez *A. iberica*. Pour cela, nous avons traité les fourmis avec un antibiotique. Chez *A. senilis*, qui représente le cas « classique », la production d'alcaloïdes chute considérablement, alors que le traitement est sans effets sur *A. iberica*. Le traitement induit aussi une augmentation des hydrocarbures cuticulaires chez *A. senilis*, mais pas chez *A. iberica*. Ces données indiquent un rôle bactérien, le microbiome de ces fourmis restant à étudier.

© 2018 Publié par Elsevier Masson SAS au nom de Académie des sciences. Cet article est publié en Open Access sous licence CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail addresses: alain.lenoir@univ-tours.fr (A. Lenoir), severine.devers@univ-tours.fr (S. Devers).




Fig. 1. Hydrocarbon quantities (ng/thorax) in *Aphaenogaster* ants. se: *Aphaenogaster senilis*, ib: *Aphaenogaster iberica*, C: control, R: Rifampicin treatment. se R is significantly different from the other groups.

196 ng; 152 ng; 225 ng versus 230 ng; 160 ng; 263 ng for the controls and the treated ants, respectively; $U = 10$; $p = 0.69$). Furthermore, no difference appeared between the controls of the two species ($U = 13$; $p = 0.16$).

A cluster analysis of the hydrocarbon profiles did not indicate a difference between control and treated ants in both species (figures not presented). It also showed that the antibiotic treatment increases the production of hydrocarbons in *A. senilis* without changing the profile, and had no effect on *A. iberica*.

3.2. Glands

The *A. senilis* poison glands contain alkanes (mean: 3%), alkenes (6.6%), aldehydes (2.6%), and a great quantity of alkaloids (87.5%) as previously presented in [3], whereas the Dufour glands, devoid of alkaloid (0%), contain alkanes (51%), alkenes (47%), and 1% of aldehydes (Table 1).

The *A. iberica* poison glands contain alkaloids (6.1%), alkanes (24.5%), alkenes (62.1%), and small quantities of aldehydes (3%), the composition of the Dufour gland being quite similar, containing alkaloids (1.8%), alkanes (38.4%), alkenes (54.6%), and aldehydes (3.1%).

The quantities of substances in the glands are given in Table 1. The poison glands of *A. senilis* are significantly bigger than those of *A. iberica* and contain more substances ($n = 18$ and $n = 10$ respectively; median, lower and upper quartiles; 383.8 ng versus 41.6 ng; for *A. senilis* and *A. iberica*, respectively; $U = 25$; $p = 0.001$). The same is true for the Dufour glands ($n = 17$ and $n = 10$, respectively; 842.5 ng versus 23.2 ng for *A. senilis* and *A. iberica*, respectively; $U = 0$; $p < 0.001$).

The total content of the *A. senilis* poison gland did not change after the antibiotic treatment ($n = 18$ and $n = 20$, respectively; 383.8 ng versus 90.4 ng, $U = 125$; $p = 0.11$; Table 1). If we consider the classes of substances, there is

no difference for alkanes ($U = 163$; $p = 0.63$), alkenes ($U = 168$; $p = 0.74$), and aldehydes ($U = 177$; $p = 0.94$), but the alkaloid quantities were significantly higher in the control compared to workers treated with the antibiotic (289 ng versus 17.3 ng for controls and treated ants, respectively; $U = 90$; $p = 0.008$; Fig. 2), representing a decrease of 94%.

Still concerning *A. senilis*, the total content of the Dufour gland increased after the Rifampicin treatment (842.5 ng versus 1827.7 ng for the control and the treated ants, respectively; $U = 109$; $p = 0.50$; Table 1).

The content of the *A. iberica* glands did not change with the treatment (Table 1; poison gland: 41.6 ng versus 78.9 ng; $U = 34$; $p = 0.09$; Dufour glands: 23.2 ng versus 24.0 ng; $U = 37$; $p = 0.14$).

4. Discussion

We observed a general effect of the antibiotic in *A. senilis* with an increase in the cuticular hydrocarbon quantities likely corresponding to a reaction of the ants to the stress induced by the decrease in bacteria quantities. This is reminiscent of the case of *Camponotus* ants that have mutualistic endosymbiotic bacteria with alimentary and immune functions [7]. Here too, the antibiotic treatment results in higher production of cuticular hydrocarbons compared to controls, plus a melanization of the cuticle [8]. In *Camponotus* spp. as well as *A. senilis*, the cuticular hydrocarbon profiles are not modified by the antibiotic treatment, but by a larger quantity of hydrocarbons produced, likely in response to a decrease in the number of symbiotic bacteria.

We also observed a spectacular effect of the antibiotic on the production of alkaloids by the poison gland of *A. senilis* with a decrease of 94%. This can be due to a direct production by bacteria which may be endosymbiotic in the




Fig. 2. Alkaloid quantities (ng/gland) in poison glands of *Aphaenogaster senilis*. C: control, R: Rifampicin treatment. The difference is highly significant ($p=0.008$).

membrane of the poison gland or by other bacteria in the ant body acting on the synthesis pathway of the alkaloids. In their review on toxins, Touchard et al. [9] said that “It is also possible that a microbial symbiont may have been involved in the production of intermediary compounds, as was suggested for some alkaloids found in sponges and briefly hinted at in ants by Saporito et al. [10].” Also, the production of pyrazines depending on bacteria was noted in Attine ants [11], and it was suggested by different researches in ant-fungus species [12,13]. Nevertheless, the production of alkaloids by free bacteria was noted only in a few marine species [14].

Aphaenogaster iberica differs from *A. senilis* in its geographical distribution and its mode of recruitment of nestmates [15]. We saw here that these two ant species also differ in their cuticular hydrocarbons and glandular contents, *A. iberica* lacking alkaloids, or they are present only as traces [3]. The fact that *A. iberica* did not react to the antibiotic treatment is likely due to the absence of symbiotic bacteria in this species. More research is necessary to know in detail the microbiome of these ant species.

Acknowledgments

This work was supported by the PRES Centre Val-de-Loire Université (APR-IA 2012). We are grateful to Andrea Yockey-Dejean for proofreading the manuscript.

References

- [1] J.C. Braekman, D. Daloze, J. Pasteels, Alkaloids in animals, in: M.F. Roberts, M. Wink (Eds.), *Biochemistry, Ecology and Medical Applications*, Plenum Press, New York, 1998, pp. 349–378.
- [2] S. Leclercq, J.-C. Braekman, D. Daloze, J. Pasteels, The defensive chemistry of ants, *Prog. Chem. Org. Nat. Prod.* 79 (2000) 116–229.
- [3] A. Lenoir, A. Benoist, A. Hefetz, W. Francke, X. Cerdá, R. Boulay, Trail-following behaviour in two *Aphaenogaster* ants, *Chemoecology* 21 (2011) 83–88.
- [4] M.O. Ramalho, O.C. Bueno, C.S. Moreau, Species-specific signatures of the microbiome from *Camponotus* and *Colobopsis* ants across developmental stages, *PLoS One* 12 (2017) e0187461.
- [5] J.A. Russell, C.F. Funaro, Y.M. Giraldo, B. Goldman-Huertas, D. Suh, D.J.C. Kronauer, C.S. Moreau, N.E. Pierce, A veritable menagerie of heritable bacteria from ants, butterflies, and beyond: broad molecular surveys and a systematic review, *PLoS One* 7 (2012) e51027.
- [6] R. Koga, T. Tsuchida, M. Sakurai, T. Fukatsu, Selective elimination of aphid endosymbionts: effects of antibiotic dose and host genotype, and fitness consequences, *FEMS Microbiol. Ecol.* 60 (2007) 229–239.
- [7] D. de Souza, A. Bézier, D. Depoix, J.-M. Drezen, A. Lenoir, *Blochmannia* endosymbionts improve colony growth and immune defence in the ant *Camponotus fellah*, *BMC Microbiol.* 9 (2009) 29.
- [8] D. de Souza, S. Devers, A. Lenoir, *Blochmannia* endosymbionts and their host, the ant *Camponotus fellah*: cuticular hydrocarbons and melanization, *C.R. Biologies* 334 (2011) 737–741.
- [9] A. Touchard, S. Aili, E. Fox, P. Escoubas, J. Orivel, G. Nicholson, A. Dejean, The biochemical toxin arsenal from ant venoms, *Toxins* 8 (2016) 30.
- [10] R.A. Saporito, H.M. Garraffo, M.A. Donnelly, A.L. Edwards, J.T. Longino, J.W. Daly, Formicine ants: an arthropod source for the pumiliotoxin alkaloids of dendrobatid poison frogs, *Proc. Natl. Acad. Sci. USA* 101 (2004) 8045–8050.
- [11] E.A. Silva-Junior, A.C. Ruzzini, C.R. Paludo, F.S. Nascimento, C.R. Currie, J. Clardy, M.T. Pupo, Pyrazines from bacteria and ants: convergent chemistry within an ecological niche, *Sci. Rep.* 8 (2018) 2595.
- [12] E.J. Choi, S.-J. Nam, L. Paul, D. Beatty, C.A. Kauffman, P.R. Jensen, W. Fenical, Previously uncultured marine bacteria linked to novel alkaloid production, *Chem. Biol.* 22 (2015) 1270–1279.
- [13] D.-C. Oh, M. Poulsen, C.R. Currie, J. Clardy, Dentigerumycin: a bacterial mediator of an ant-fungus symbiosis, *Nat. Chem. Biol.* 5 (2009) 391.
- [14] S. Haeder, R. Wirth, H. Herz, D. Spiteller, Candidicin-producing *Streptomyces* support leaf-cutting ants to protect their fungus garden against the pathogenic fungus *Escovopsis*, *Proc. Natl. Acad. Sci. USA* 106 (2009) 4742–4746.
- [15] X. Espadaler, J.M. Riasol, Distribución, variabilidad y sinonimias en *Aphaenogaster iberica* [sic] Emery, 1908 y dos adiciones a la fauna ibérica (Hymenoptera, Formicidae), *Actas del I Congreso Ibérico de Entomología*, Vol. 1, Universidad de León, León, Spain, 1983 pp. 219–228.