

the *Wolbachia* microorganism, which reduces the ability of the mosquito to transmit dengue, Zika, and chikungunya. The *Wolbachia* method is safe, natural, and self-sustaining, and has the potential to achieve significant public health impact in areas endemic to these viruses.

WMP was formerly known as “Eliminate Dengue: Our Challenge.” The name has been changed due to rapid global program expansion and also to reflect that the method works against several diseases, not just dengue. WMP is active in 12 countries: Australia, Vietnam, Indonesia, Pacific Islands (Fiji, Vanuatu, Kiribati, New Caledonia), India, Sri Lanka, Colombia, Mexico, and Brazil, and discussions are at an advanced stage with several others. In Brazil, it has been conducted since 2012 by the Fundação Oswaldo Cruz (Fiocruz), based in Rio de Janeiro.

The aim of WMP is to perform controlled releases of *Aedes aegypti* containing *Wolbachia*, so there will be a gradient substitution of local the *Aedes aegypti* population by *Aedes aegypti* with *Wolbachia*. This is possible, through both the CI mechanism and the vertical transmission, guaranteeing the self-sustainability of the method.

Steps of *Wolbachia* Method

To achieve the goal of substitution of the local population of *Aedes aegypti* with *Aedes aegypti* with *Wolbachia*, the actions of WMP are divided into three main stages. (i) In the Community Engagement phase, project teams interact with the population and institutions (health clinics, schools, community leaders) for disseminating information about the initiative. At this stage, surveys are conducted to measure understanding and acceptance of the local population about the method. In addition, a Community Reference Group is set up, a local committee who monitors all actions taken in the locality, and communication channels are established with the community, including telephone, e-mail, face-to-face interactions, and social media. (ii) After the Community Engagement phase and community approval begins the release of *Aedes aegypti* mosquitoes with *Wolbachia*. Mosquitoes are bred in the Fiocruz insectary, which has similar humidity and temperature characteristics as those that mosquitoes encounter in the external environment. Adult releases happen in the early hours of the morning and are made by car. In some regions, this work is done on foot, by health surveillance agents and community health agents of the municipal governments of Rio de Janeiro and Niterói. (iii) Monitoring of the *Aedes aegypti* population in these areas initiate during mosquito releases. Traps are installed in homes or commercial establishments that voluntarily provide a location for the traps. Weekly, technicians go to these sites to collect mosquitoes. At the project facilities, the captured mosquitoes are separated, identified and the *Aedes aegypti* are sent to the Diagnostic Laboratory, where they are checked individually. This analysis aims to identify the DNA of *Wolbachia* bacteria in the mosquito organism and is an indicator of the establishment of the local mosquito population with *Wolbachia*.

Large-scale expansion

WMP began preparative studies in Brazil in 2012 and ethical and regulatory approvals. In 2014, releases began in the pilot areas, Jurujuba, in Niterói-RJ, and Tubiacanga, in the city of Rio de Janeiro. The last releases of mosquitoes at these sites were carried out in January 2016, and since then monitoring has revealed *Wolbachia*'s establishment of more than 90%. In November 2016, large-scale expansion began for 28 neighborhoods in Niterói, to protect 270,000 people.

In Rio de Janeiro, large-scale releases began in August 2017, with the plan of reaching 19 neighborhoods, where 0.9 million people live. Concomitantly with community engagement, mosquito release and monitoring activities, an epidemiological study is underway to measure the impact of our work in reducing disease transmission. Preliminary results show already a reduction on the incidence of dengue, Zika, and chikungunya in areas where *Wolbachia* is present.

Through a partnership with the city of Belo Horizonte, WMP Brazil will start activities in an pilot area with 60,000 inhabitants, probably in late 2019.

Regulatory and ethical approvals

For pilot projects, regulatory and ethical approvals were granted in 2014. The Temporary Special Registry (Registro

Especial Temporário–RET–, in Portuguese) was granted after evaluating the project simultaneously by three governmental areas: National Agency of Sanitary Surveillance–ANVISA–, Ministry of Agriculture, Livestock and Supply–MAPA–, and the Brazilian Institute of Environment and Renewable Natural Resources–IBAMA. Ethical approval was also granted in early 2014, following a thorough evaluation by the National Commission for Research Ethics (CONEP). For the large-scale expansion, all the regulatory documents have been renewed.

In February 2016, the Ministry of Health recommended the *Wolbachia* method as promising in vector control in response to Zika's national emergency. This recommendation was reinforced in March of that year by the World Health Organization (WHO) through its Vector Control Advisory Group (VCAG).

Funding

The project has been historically financed in Brazil by the Ministry of Health, with an in-kind contribution from Fiocruz and, internationally, by the Bill and Melinda Gates Foundation, through Monash University, Australia. The municipalities of Rio de Janeiro and Niterói provided an important contribution with local infrastructure and human resources.

Disclosure of interest The author declares that he has no competing interest.

Further reading

L.A. Moreira, I. Iturbe-Ormaetxe, J.A. Jeffery, G. Lu, A.T. Pyke, L.M. Hedges, B.C. Rocha, S. Hall-Mendelin, A. Day, M. Riegler, L.E. Hugo, K.N. Johnson, B.H. Kay, E.A. McGraw, A.F. van den Hurk, P.A. Ryan, S.L. O'Neill, A *Wolbachia* symbiont in *Aedes aegypti* limits infection with dengue, Chikungunya, and *Plasmodium*, Cell 139(7) (2009) 1268–1278.

H.L. Dutra, M.N. Rocha, F.B. Dias, S.B. Mansur, E.P. Caragata, L.A. Moreira. *Wolbachia* Blocks Currently Circulating Zika Virus Isolates in Brazilian *Aedes aegypti* Mosquitoes, Cell Host Microbe. 19(6) (2016) 771–774.

S.L. O'Neill, The Use of *Wolbachia* by the World Mosquito Program to Interrupt Transmission of *Aedes aegypti* Transmitted Viruses, Adv. Exp. Med. Biol. 1062 (2018) 355–360.

<https://doi.org/10.1016/j.crv.2019.09.020>

20

Ecosystem services provided by insects for achieving sustainable developmental goals

Olivier Dangles

Département “Écologie, biodiversité et fonctionnement des écosystèmes continentaux” (ECOBIO), Montpellier, France
E-mail address: olivier.dangles@ird.fr

Ecosystem services underpin all dimensions of human well-being. As a consequence, it is crucial to integrate ecosystem services into strategies for achieving Sustainable Development Goals (SDGs). Because insects and other invertebrates have profound and well-identified influences on many ecosystem services (e.g., pollination and biological control) and SDGs (e.g., crop pest and disease vectors), insect research and development have a great potential to address current global challenges. We argue that time is ripe to put more efforts in developing integrated research on the ecosystem services provided by insects, as they may result in solutions to achieve many SDGs. We provide evidence of insects' utility to address global challenges and propose a framework of the needed shift in the perception of insects from enemies to allies, providers of ecosystem services, and then to solutions to achieve SDGs. We further advocate that making a place for SDG-relevant research on insects' ecosystem services requires transforming existing academic knowledge into application-driven science, a potential upscaling of local solutions and socio-economic relevance.

Disclosure of interest The author declares that he has no competing interest.

<https://doi.org/10.1016/j.crv.2019.09.021>

