



INSTITUT DE FRANCE
Académie des sciences

Comptes Rendus

Biologies


Jacques Demongeot and Hervé Seligmann

Theoretical minimal RNA rings mimick molecular evolution before tRNA-mediated translation: codon-amino acid affinities increase from early to late RNA rings

Volume 343, issue 1 (2020), p. 111-122.

<<https://doi.org/10.5802/crbio.1>>

© Académie des sciences, Paris and the authors, 2020.
Some rights reserved.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*Les Comptes Rendus. Biologies sont membres du
Centre Mersenne pour l'édition scientifique ouverte*
www.centre-mersenne.org



News and events / *Actualités*

Theoretical minimal RNA rings mimic molecular evolution before tRNA-mediated translation: codon-amino acid affinities increase from early to late RNA rings

Des ARNs circulaires possibles chaînons de l'évolution de la traduction médiée par les ARNs : les affinités codons-acides aminés augmentent en fonction de l'ancienneté des ARN circulaires

Jacques Demongeot^{*, a} and Hervé Seligmann^{a, b}

^a Université Grenoble Alpes, Faculty of Medicine, Laboratory AGEIS EA 7407, Team Tools for e-Gnosis Medical, F-38700 La Tronche, France

^b The National Natural History Collections, The Hebrew University of Jerusalem, 9190401 Jerusalem, Israel.

E-mail: jacques.demongeot@yahoo.fr (J. Demongeot).

Abstract. Nucleotide affinities for noncovalent interactions with amino acids produce associations between mRNAs and cognate peptides, potentially regulating ribosomal translation. Correlations between nucleotide affinities and residue hydrophobicity are explored for 25 theoretical minimal RNA rings, 22 nucleotide-long RNAs designed *in silico* to code for each amino acid once after three translation rounds, and forming stem-loop hairpins. This design presumably mimics life's first RNAs. RNA rings resemble consensual tRNAs, suggesting proto-tRNA function, predicted anticodon and cognate amino acid. The 25 RNA rings and their presumed evolutionary order, deduced from the genetic code integration order of the amino acid cognate to their predicted anticodon, produces noteworthy associations with several ancient properties of the cell's translational machinery. Here we use this system to explore the evolution of codon affinity-residue hydrophobicity correlations, assuming these reflect pre-tRNA and pre-ribosomal translations. This hypothesis expects that correlations decrease with genetic code inclusion orders of RNA ring cognates. RNA ring associations between nucleotide affinities and residue hydrophobicities resemble those from modern natural genes/proteins. Association strengths decrease with genetic code inclusion ranks of proto-tRNA cognate amino acids. *In silico*

* Corresponding author.

design of minimal RNA rings didn't account for affinities between RNA and peptides coded by these RNAs. Yet, interactions between RNA rings and translated cognate peptides resemble modern natural genes. This property is strongest for ancient RNA rings, weakest for recent RNA rings, spanning a period during which modern tRNA- and ribosome-based translation presumably evolved. Results indicate that translation lacking tRNA-like adaptors based on codon-amino acid affinities and the genetic code pre-existed tRNA-mediated translation. Theoretical minimal RNA rings appear valid prebiotic peptide-RNA world models for the transition between pre-tRNA- and tRNA-mediated translations.

Résumé. Les affinités nucléotidiques des interactions non covalentes avec les acides aminés produisent des associations entre les ARNm et les peptides apparentés, régulant potentiellement la traduction ribosomale. Les corrélations entre les affinités nucléotidiques et l'hydrophobicité des résidus sont explorées pour 25 ARNs circulaires théoriques d'une longueur minimale de 22 nucléotides, conçus *in silico* pour coder chaque acide aminé une fois et une seule après trois tours de traduction et former des structures « hairpin » (épingles à cheveux) maximales. Cette conception imite vraisemblablement les premiers ARNs de la vie. Les ARNs circulaires ressemblent à des ARNs consensus, suggérant une fonction de proto-ARNt, mettant en jeu un anticodon prédit et un acide aminé apparenté. Les 25 ARNs circulaires et leur ordre d'évolution présumé, déduits de l'ordre d'intégration dans le code génétique de l'acide aminé apparenté à leur anticodon prédit, montrent des associations remarquables avec plusieurs propriétés anciennes du mécanisme de traduction de la cellule. Ici, nous les utilisons pour explorer l'évolution des corrélations entre hydrophobicité d'un résidu et affinité à son codon, en supposant qu'elles reflètent des propriétés de la traduction pré-ARNt et pré-ribosomique. Cette hypothèse suppose que les corrélations diminuent avec les ordres d'inclusion dans le code génétique du codon apparenté à un ARN circulaire. Les associations entre les affinités des nucléotides et les hydrophobicités des résidus ressemblent à celles existant actuellement entre gènes et protéines. Les forces d'association diminuent avec les rangs d'inclusion dans le code génétique des acides aminés apparentés au proto-ARNt. La conception des ARNs circulaires minimaux ne tient pas compte des affinités entre ceux-ci et les peptides qu'ils codent. Cependant, les interactions entre les ARNs circulaires et leurs peptides apparentés ressemblent à celles observées pour les gènes naturels actuels. Cette propriété est plus forte pour les ARNs circulaires anciens et plus faible pour les récents, couvrant une période au cours de laquelle la traduction moderne à base d'ARNts et de ribosomes a probablement évolué. Les résultats indiquent que la traduction usant d'adaptateurs fondés sur les affinités codon-acide aminé et sur le code génétique primitif a préexisté à une traduction médiée par l'ARNt. Les ARNs circulaires minimaux théoriques apparaissent ainsi comme des modèles prébiotiques pour la transition entre les traductions à médiation pré-ARNt et ARNt.

Keywords. Secondary structure, Cyclic code, Circular RNAs, Circular code, Nonribosomal translation.

Mots-clés. Structure secondaire, Code cyclique, ARNs circulaires, Code circulaire, Traduction non ribosomale.

Manuscript received 15th June 2019, accepted 21st February 2020.

1. Introduction

Codon-amino acid assignments according to the genetic code are not random [1] because many codon and anticodon properties correlate with properties of cognate amino acids [2–4]. Numerous analyses indicate the genetic code minimizes error impacts for various processes and properties: mutations on protein structure [5, 6], how proteins fold [7–9], tRNA misloading by tRNA synthetases [10–13], frameshifts during translation before and after these occur (before occurrence [14–17], after occurrence [18–23]). The genetic code maximizes the diversity of physicochemical properties of coded amino acids [24, 25] and optimizes error according to several properties at the same time [1].

1.1. Genetic code evolution

Numerous hypotheses on inclusion orders of amino acids in the genetic code have been deduced from different properties and experiments. These hypotheses predict orders by which amino acids were assigned to codon(s) [26, 27]. Most of these unrelated hypotheses predict congruent orders, justifying a consensual order of amino acid inclusion in the genetic code [26, 27]. Ulterior analyses produced genetic code inclusion orders that converge with these amino acid inclusion ranks. The average position of amino acid types in modern proteins correlates with these inclusion ranks. The mean position of ancient amino acids is closer to the protein C-terminus, recently included amino acids are on average closer

to the protein N-terminus, which corresponds to the gene's 5' extremity with the initiation/start codon [28].

1.2. Stereochemical genetic code codon-amino acid assignments

This shows that structures of modern biomolecules embed fossilized information on the origins of life. Indeed, ribosome X-ray crystal structures show that contacts between ribosomal RNA nucleotide triplets and ribosomal protein residues favor amino acid-codon/anticodon assignments. Amino acids that integrated early the genetic code favor contacts with their codons, late amino acids favor contacts with their anticodons [29]. This shows two different steps during the formation of the genetic code: (1) an adaptor/tRNA-free translation where codons interacted directly with amino acids, and (2) tRNA-based translation. Both results confirm, at the level of ribosomal structure, the hypothesis that (at least some of) the genetic code's assignments result from nucleotide-amino acid affinities due to stereochemical interactions.

The importance of stereochemical interactions is not surprising, as these also determined homochirality of life's major molecules, right-handed RNA (R-RNA) and left-handed amino acids (L-amino acids), which interact better when these have opposite handedness [30–34]. The first step in genetic code evolution deduced from ribosomal crystal structures suggests direct codon-amino acid stereochemical interactions. This is in line with observations that in most mRNAs, nucleotide affinities for amino acids correlate with amino acid polar requirements [35–37]. These observations imply that frequencies of observed contacts between nucleotides and amino acids, from which nucleotide-amino acid affinities are deduced, are approximated by the amino acid hydrophobicity, as these interactions are broadly spoken hydrophobic interactions. This implies that genetic code assignments resulted from stereochemical interactions before proto-tRNAs switched from their function as initiators of RNA/DNA polymerizations [38, 39] to their main modern function in translation. Translations by direct codon-amino acid interactions are in line with a primordial peptide-RNA world, rather than a primordial RNA world [40].

Ribosomal translation of RNA




Figure 1. Translation in three rounds of theoretical minimal RNA ring. Numbers indicate codon order. * indicates stop codon.

1.3. Theoretical minimal RNA rings as models for the primitive RNA world

Here we study associations between codon affinities for their cognate amino acid and that amino acid's hydrophobicity, specifically in the context of 25 theoretical minimal RNA rings [41–45]. These 22-nucleotide long RNAs were designed to code for a start and a stop codon and once for each of the genetic code's 20 amino acids by three consecutive translation rounds (Figure 1). This ensures maximal coding capacity over the shortest possible length. Their design was also constrained to form a stem-loop hairpin. Independent secondary structure analyses confirmed most theoretical minimal RNA rings resemble proto-tRNAs [44, 45]. A cognate amino acid, in the sense of tRNA acylation, can be assigned to each RNA ring, based on their predicted anticodon sequence [46]. Theoretical RNA rings also fit predictions that circular RNAs had critical roles in the early formation of protogenomes [47]. Most importantly, their design assumes the genetic code's pre-existence to tRNA-assisted translation. It is of note that some RNA ring secondary structures resemble more rRNAs than tRNAs [46], and that maturation of some modern tRNAs includes circularization [48].

The design of theoretical minimal RNA rings minimizes sequence length and maximizes the diversity of coded amino acids. Circularized tRNAs are intermediates in post-transcriptional modifications of some tRNAs [40]. Circular RNAs regulate gene expression in modern cells [49–55] and are occasionally translated [56]. They code for proteins with different functions, frequently in muscles of mammals [57]. The assumption of overlap coding is also realistic: artificially designed genes coding over three overlapping frames for three predefined proteins from different protein domains are relatively easy to produce [58]. This fits with observations suggesting that

overlap coding is a common property of genes [59–67]. Indeed, the genetic code is optimized for coding after frameshifts [1, 65–67].

We expect that associations between RNA ring codon affinities with cognate amino acid hydrophobicity in the peptides coded by the RNA rings resemble those observed in natural modern genes. Two evolutionary scenarios exist in this context. (1) This association appeared spontaneously and, in modern genes, is a carryover from an earlier translation mechanism lacking tRNA and ribosome. In this case its strength should decrease with the genetic code inclusion order of cognate amino acids of RNA rings. (2) This association results from an evolutionary process and reflects processes that still occur in modern cells. According to this scenario, the strength of this association should increase from early to late RNA rings.

2. Materials and methods

We translated the 25 peptides coded by the theoretical minimal RNA rings (Table 1) according to the standard genetic code and estimated the strength of associations between their codon-amino acid affinity and the hydrophobicity of their cognate amino acids forming the peptide. Codon-amino acid affinity is calculated by summing the affinity of single nucleotides forming the codon with the amino acid coded by the codon [35]. These single nucleotide affinities for amino acids are estimated from nucleotide-amino acid contact frequencies in crystal structures of interacting RNA-protein complexes [35].

These linear sums of affinities of single nucleotides with amino acids produce the same trinucleotide affinity for trinucleotides N1N2N3, N1N3N2, N2N1N3, N2N3N1, N3N1N2 and N3N2N1, though in most cases, these permutations do not code for the same amino acid. Hence this sum of single nucleotide affinities must be considered as an approximation of trinucleotide-amino acid affinities. Nevertheless, biases for codon-amino acid contacts observed in ribosomal crystal structures [29] correlate with these codon affinities calculated by summing single nucleotide affinities (Figure 2). This preliminary observation justifies the linear sum of single nucleotide affinities to approximate codon-amino acid affinities.

Pearson correlation coefficients r were calculated between codon affinities and several estimates of amino acid polar requirements, experimental and calculated polar requirements [69–71], two estimates of proportions of accessible versus buried amino acid surfaces in proteins [70, 71], and several estimates of affinities of side chains of amino acids for solvent water [68, 72], other solvents [73, 74] and for water at different temperatures (at 25 °C versus 100 °C [75]).

3. Results and discussion

3.1. Codon affinity-amino acid hydrophobicity correlations at specific codons

Our first analysis considers each codon across the 25 theoretical minimal RNA rings and calculates the correlation between codon affinity and hydrophobicity for the amino acid coded by each codon. This produces 21 Pearson correlation coefficients r for each estimate of hydrophobicity, for all 21 codon positions coding for an amino acid, and aligned as shown in Table 2. These correlations are expected to reflect those observed for regular mRNAs and their cognate proteins. Affinities are estimated from nucleotide-residue contact frequencies and log-transformed to correspond to contact enthalpies. If affinities are proportional to amino acid hydrophobicities as observed in regular genes, correlations with log-transformed estimates of affinities should produce negative correlations with amino acid hydrophobicities.

Among seventeen different hydrophobicity scales tested, the highest number of positions for which negative associations between codon affinity and amino acid hydrophobicity is 20 among 21 positions, for the scales based on the distribution of equilibria between the dilute aqueous solution and the vapor phase at room temperature (log (RHvap/RHwater), hydration potential [72]). Note that Pro is absent from these scales. Hence the pattern observed in regular mRNAs and proteins (negative correlations) is also observed for most positions in theoretical RNA rings according to these hydrophobicity scales. This majority is statistically significant ($P = 0.00001$, two tailed sign test) and remains so also after considering test multiplicity, even according to Bonferroni's over-conservative correction ($P = 0.05/17 = 0.002941$,

Table 1. Minimal RNA rings and corresponding peptide sequences according to regular translations. In column 1, r indicates secondary structures resembling more rRNAs, remaining are more tRNA-like, capitals indicate the presumed cognate amino acid of the RNA ring if considered as a proto-tRNA according to its predicted anticodon. * indicates stop codons. Columns 4 and 5 indicate Pearson correlation coefficients $r \times 100$ between codon-amino acid affinity [35] and amino acid hydrophobicity estimated from distribution equilibria between vapor and aqueous solution (R1, $\log(RH_{vap}/RH_{water})$, R2, hydration potential, [68])

RNA	AA	CDs	Peptide	R1	R2
1	F	<u>AATTCATGCCAGACTGGTATGA</u>	NSCQTGMKFM PDWY EIHARLV*	-13.0	-13.7
2	M	<u>CATGCCAGAAATCTGGTATGA</u>	HARNSGMT CQKFWYDM PEILV*	-10.7	-11.8
3	S	<u>ATGGTGCCACTATCAAGATGA</u>	MVPLFKMNGATI QDEWCH YSR*	-20.9	-15.0
4	Pyl	<u>ATGCTATTCACCAAGATGGTGA</u>	MLFTKMNVAIHQD GECYSP RW*	-13.4	-8.3
5	V	<u>ATGGTGCTACCATTCAAGATGA</u>	MVLPFKMNGATI QDEWCY HSR*	-16.8	-9.7
6	N	<u>ATGGCCTATTCACAAGATGTGA</u>	MAYSQDVNGLFTR CEWPIH KM*	-19.1	-16.8
7	D	<u>ATGCCACTGGTATCAAGATGA</u>	MPLVFKMNATGI QDECHW YSR*	-19.8	-14.1
8	R	<u>ATGTGGCCTACATTCAAGATGA</u>	MWPTFKMNVAYI QDECGL HSR*	-22.6	-18.6
9	Sec	<u>ATGCCAAGATGGTATCACTGA</u>	MPRWYSLNAKMV FTECQD GIH*	-19.8	-14.1
10	L	<u>GCAATGTTTATGGAGACCATAA</u>	AMFMETISNVYGD HKQCLWR P*	-25.8	-19.8
11	P	<u>TATGTTTGGAGACCAAGCATAA</u>	YVWRPSMICLET KHNMF GDQA*	-24.4	-18.9
12	E	<u>TTCATGCCAGAAACTGGTATGA</u>	FMPETGMIHARN WYDSCQ KLV*	-24.7	-19.7
13	G	<u>ATGGTACTGCCATTCAAGATGA</u>	MVLPFKMNGTAI QDEWYCH SR*	-23.6	-16.0
14	I	<u>GCAGAATGTTTATGGACCATAA</u>	AECLWTISR MFMDHKQ NVYGP*	-29.0	-24.2
15	Q	<u>AATATGTTTGGACCAAGCATAG</u>	NMFGPSIEYV WTKHRICL DQA*	-35.7	-34.5
16	L	<u>TATGTTTGGAAAGCCAGACATAA</u>	YVWKPDII CLEARHNMF GSQT*	-20.8	-22.7
17	K	<u>TACATTTGGAAAGCCAGATGTAA</u>	YIWKPDVMH LEARCNTF GSQM*	-27.1	-27.0
18	S	<u>ACAATGTTTATGGAAGCCATAG</u>	TMFMEAIDNVY GSHRQCL WKP*	-32.9	-37.8
19	A	<u>ATGGAAGCCATTTACAATGTAG</u>	MEAIYNVDG SHLQCRW KPFTM*	<u>-38.4</u>	<u>-38.3</u>
20	W	<u>TACAGATGGAAGCCATTGTAA</u>	YRWKPFVMQ MEICNTD GSHL*	-27.1	-27.0
21	C	<u>AACATGCCAGATTCTGGTATGA</u>	NMPDSGMKHAR FWYETC QILV*	-24.8	-17.7
22	H	<u>TGCCAGAAACATCTGGTATGA</u>	CQKHS GMPETFWYD ARNILV*	-21.8	-17.7
23	T	<u>TATGGTTCTGCAAGAACCATGA</u>	YGSARTMIWF CKNHDM VLQEP*	-28.2	-17.2
24	Y	<u>TACCATTCTGCAAGAATGGTGA</u>	YHSARMVMP FCKNGD TILQEW*	-21.7	-13.5
25	G	<u>ATGGTGCCATTCAAGACTATGA</u>	MVPFKTMNGAI QDYEWCH SRL*	-19.9	-14.5

which is still higher than $P = 0.00001$). The overall tendency is weaker when considering other hydrophobicity scales, notably those obtained according to similar equilibria between solvents, but other than water, the natural solvent of biological systems. These are not reported in the following as results were systematically strongest for the scales presented in [72].

This analysis also produces statistically significant correlations at specific codons in RNA rings. All six

statistically significant correlations are negative as expected from modern mRNAs and proteins and are located at the transition between each translation round, for amino acids coded by the 6th, 7th, 8th, 12th, 15th, and 19th codons in RNA rings. This would suggest that peptides interact with their template RNA ring at each new translation round. This pattern emerging from theoretical RNA rings is not part of their design, and might reflect properties of RNA-templated, pre-tRNA translation.

Table 2. Pearson correlation coefficients r between codon affinity and cognate amino acid hydrophobicity (as for Table 1, [68], Log is for log (RHvap/RHwater) and Pot for the hydration potential used as hydrophobicity estimates) for each of the 21 amino acid-coding codons in the 25 theoretical minimal RNA rings from Table 1. Analyses are across the 25 RNA rings, keeping constant the codon position. Asterisks show $P < 0.05$. P values are two tailed

Position	Log	Pot
Round 1		
1	0.100	0.101
2	-0.070	-0.002
3	-0.292	-0.231
4	-0.290	-0.265
5	-0.508*	-0.331
6	-0.618*	-0.602*
7	-0.697*	-0.697*
Round 2		
8	-0.535*	-0.612*
9	-0.333	-0.321
10	-0.074	-0.043
11	0.006	-0.067
12	-0.607*	-0.544*
13	-0.599	-0.137
14	-0.063	-0.260
Round 3		
15	-0.775*	-0.459*
16	-0.116	-0.116
17	-0.055	-0.047
18	-0.375	-0.333
19	-0.380	-0.433
20	-0.028	-0.019
21	-0.210	-0.199
22-stop	n.d.	n.d.

Correlations obtained for hydrophobicity scales estimated from distribution equilibria between neutral solution and cyclohexane at 25 °C and 100 °C [76] produce less clear patterns. Patterns are slightly stronger at 100 °C than 25 °C, putatively suggesting that the codon-amino acid associations we observe in modern organisms evolved originally at high tem-

peratures, in line with some previous considerations on origins of life and the genetic code [75, 77–80].

3.2. Codon affinity-amino acid hydrophobicity correlations for specific RNA rings

Our second analysis tests for correlations between codon affinity and amino acid hydrophobicity (distribution equilibria between the dilute aqueous solution and the vapor phase at room temperature [72]) across codons, for each specific RNA ring (Table 1). All 25 correlations are negative ($P < 0.05$ for RNA ring 19), a statistically significant majority of cases ($P = 0.00000003$, two tailed sign test). This means that the association between codon affinity and amino acid hydrophobicity for the 25 RNA rings resembles that observed in modern genes.

3.3. Evolution of associations between codon affinity and amino acid hydrophobicity

Each of the 25 theoretical minimal RNA rings, because of their similarity with consensus tRNA sequences, can be assigned to a cognate amino acid according to their predicted anticodon sequence, assuming a tRNA-like function. Each of these tRNA-cognate amino acids is assigned a rank of inclusion in the genetic code, according to various hypotheses on genetic code inclusion history. Interestingly, and despite that these numerous hypotheses have very different backgrounds, from mathematical (e.g. the natural circular code, [14], the algebraic model [81]), physicochemical (e.g. amino acid yields in experiments mimicking early earth condition [82], amino acid size-complexity [83]) to physiological (e.g. the metabolic coevolution hypothesis [84], the self-referential model [85]), they tend to predict overall congruent histories of genetic code assignments of amino acids (40 hypothetical inclusion orders reviewed in [26, 27]). Some hypotheses are derived from aminoacylating rules of small RNA helices [86, 87] and tRNAs [88], affecting protein folding [9, 89], enabling to define a primordial code in tRNA acceptor stems and an associated order of integration of amino acids in the genetic code.

The inclusion ranks of the RNA ring predicted cognate amino acids are assumed to reflect the history of




Figure 2. Observed bias for contacts between codon and cognate amino acid in the ribosome crystal structure (Johnson and Lee 2010) as a function of linear sum of single nucleotide affinities for cognate amino acid averaged across all synonymous codons (strong affinities correspond to low values on the x -axis). The correlation ($r = -0.605$, one tailed $P = 0.003$) justifies the use of the sums of single nucleotide affinities to approximate codon affinities.

the theoretical organic system based on these 25 theoretical RNA rings and their cognate peptides (cognate here is used in the sense of the peptide translated from the RNA ring as an ancient template for protein synthesis). We examined the associations of the Pearson correlation coefficients between codon affinity and amino acid hydrophobicity for the 25 RNA rings and the inclusion ranks of their cognate amino acid (cognate is used here in the sense of the amino acid presumably aminoacylated to the RNA ring in relation to its presumed tRNA-like function). The RNA rings with predicted anticodons matching cognate amino acids selenocysteine and pyrrolysine were assigned penultimate and last genetic code inclusion ranks.

Scenario 1 assumes that correlations between codon affinity and amino acid hydrophobicity reflect spontaneous associations that were the physico-chemical basis for tRNA- and ribosome-free translation pre-existing modern translation. Along that sce-

nario, codon affinity-amino acid hydrophobicity associations observed today would be remnants of this earlier translation mechanism. Hence Pearson correlation coefficients r between codon affinity and amino acid hydrophobicity of RNA rings are expected to increase (become more positive) with the inclusion rank of the amino acid assigned as proto-tRNA cognate for that RNA ring.

Scenario 2 expects that correlations evolved towards becoming more similar to what is known for extant genes and proteins, which is generally a negative correlation. Hence Pearson correlation coefficients r between codon affinity and amino acid hydrophobicity of RNA rings are expected to decrease (become more negative) with the inclusion rank of the amino acid assigned as proto-tRNA cognate for that RNA ring.

Correlation coefficients r obtained for associations between codon affinities and hydrophobicity estimated from distribution equilibria between va-




Figure 3. Pearson correlation coefficient of association between amino acid polar requirement and codon affinity for that amino acid for 25 theoretical minimal RNA rings of 22 nucleotides coding after three translation rounds for a start and stop codon and one codon per amino acid, as a function of the genetic code inclusion order of the presumed cognate amino acid of that RNA ring if considered as a proto-tRNA. The tRNA cognate amino acid is determined by the nucleotide triplet at the predicted anticodon position. The genetic code inclusion order is according to Hartman's primordial one nucleotide code, assuming G and C were first [90–94].

por and aqueous solution fit scenario 1 for 38 among 40 hypotheses for genetic code evolution of amino acid inclusion orders, a significant majority ($P = 8 \times 10^{-8}$, two tailed sign test). The correlation with amino acid inclusion order is statistically significant at $P < 0.05$ for eight specific hypotheses for amino acid inclusion orders. The strongest positive association between the presumed inclusion order of the cognate amino acid of the RNA ring and the codon affinity-hydrophobicity correlation r is for Hartman's hypothesis that assumes that the genetic code started as a one-letter code, then became a two- and then three-letter code, with G and C as primordial nucleotides ($r = -0.43$, one tailed $P = 0.008$, Figure 3) [90–93]. Hence results are in line with scenario 1 that assumes that codon-amino acid hydrophobicity associations in modern gene-protein pairs are rem-

nants of an ancient tRNA- and ribosome-free translation mechanism based on these physicochemical interactions [95], which *de facto* would have produced the genetic code codon-amino acid assignments [96].

4. Conclusions

Results of analyses of peptides coded by RNA rings and RNA ring codons indicate that codon-amino acid affinity-amino acid hydrophobicity associations were strongest/most negative for RNA rings assumed most ancient. Accordingly, such associations observed in modern genes and their proteins would be remnants of a more ancient translation mechanism lacking tRNAs and ribosome, based on direct physicochemical interactions between codon and amino acid. Codon-amino acid assignments in the genetic

code would have resulted from these direct codon-amino acid interactions [96,97], which would also explain biases for codon-amino acid contacts in the ribosome ([29], and herein Figure 2).

It is unclear how the design of theoretical minimal RNA rings implies complex evolutionary properties such as associations between codon affinity and cognate amino acid hydrophobicity. However, this system actually mimicks unexpectedly numerous and complex properties of the genetic code [96, 98], tRNAs [43], protein coding genes [99, 100], their proteins [101] and of replication origins [102]. This RNA ring system enables to recover probable evolutionary scenarios of tRNAs and rRNAs [101, 103–108], and the assignment of AUG as universal initiation codon [109]. The design of RNA rings implies that the genetic code pre-existed the emergence of RNA rings. Hence because RNA rings resemble proto-tRNAs, the genetic code had to exist before tRNAs and proto-tRNAs. The association of codon affinity with hydrophobicity for RNA rings indicates that translation according to the genetic code pre-existed tRNA-mediated translation, potentially along a different codon structure where nucleotides at 1st and 2nd codon positions were switched [110, 111], but also evolved towards the situation observed in modern genes during the evolution of this RNA ring-based primordial RNA system [112–114]. Theoretical minimal RNA rings are realistic models for the RNA world during the transition from translation based on codon affinity for amino acids and lacking tRNA-like adaptors to tRNA-like mediated translation.

Acknowledgments

Thanks to Charles W. Jr Carter for his important advice on hydrophobicity estimates.

References

- [1] B. Kumar, S. Saini, "Analysis of the optimality of the standard genetic code", *Mol. Biosys.* **12** (2016), p. 2642-2651.
- [2] C. R. Woese, D. H. Dugre, S. A. Dugre, M. Kondo, W. C. Saxinger, "On the fundamental nature and evolution of the genetic code", *Cold Spring Harb. Symp. Quant. Biol.* **31** (1966), p. 723-736.
- [3] S. R. Pelc, M. G. E. Welton, "Stereochemical relationship between coding triplets and amino-acids", *Nature* **209** (1966), p. 868-870.

- [4] H. Seligmann, G. N. Amzallag, "Chemical interactions between amino acid and RNA: multiplicity of the levels of specificity explains origin of the genetic code", *Die Naturwissenschaften* **89** (2002), p. 542-551.
- [5] D. H. Ardell, "On error minimization in a sequential origin of the standard genetic code", *J. Mol. Evol.* **47** (1998), p. 1-13.
- [6] S. J. Freeland, L. D. Hurst, "The genetic code is one in a million", *J. Mol. Evol.* **47** (1998), p. 238-248.
- [7] A. Guilloux, J. L. Jestin, "The genetic code and its optimization for kinetic energy conservation in polypeptide chains", *Biosystems* **109** (2012), p. 141-144.
- [8] B. Caudron, J. L. Jestin, "Sequence criteria for the antiparallel character of protein beta-strands", *J. Theor. Biol.* **315** (2012), p. 146-149.
- [9] H. Seligmann, G. Warthi, "Genetic code optimization for cotranslational protein folding: codon directional asymmetry correlates with antiparallel betasheets, tRNA synthetase classes", *Comput. Struct. Biotechnol. J.* **15** (2017), p. 414-424.
- [10] H. Seligmann, "Do anticodons of misacylated tRNAs preferentially mismatch codons coding for the misloaded amino acid?", *BMC Mol. Biol.* **11** (2010), p. 41.
- [11] H. Seligmann, "Error compensation of tRNA misacylation by codon-anticodon mismatch prevents translational amino acid misinsertion", *Comput. Biol. Chem.* **35** (2011), p. 81-95.
- [12] H. Seligmann, "Coding constraints modulate chemically spontaneous mutational replication gradients in mitochondrial genomes", *Curr. Genomics* **13** (2012), p. 37-54.
- [13] R. M. Barthélémy, H. Seligmann, "Cryptic tRNAs in chaetognath mitochondrial genomes", *Comput. Biol. Chem.* **95** (2016), p. 119-132.
- [14] D. G. Arquès, C. J. Michel, "A complementary circular code in the protein coding genes", *J. Theor. Biol.* **182** (1996), p. 45-58.
- [15] A. Ahmed, G. Frey, C. J. Michel, "Frameshift signals in genes associated with the circular code", *Silico Biol.* **7** (2007), p. 155-168.
- [16] N. El Houmami, H. Seligmann, "Evolution of nucleotide punctuation marks: from structural to linear signals", *Front. Genet.* **8** (2017), p. 36.
- [17] G. Warthi, H. Seligmann, "Transcripts with systematic nucleotide deletion of 1–12 nucleotide in human mitochondrion suggest potential non-canonical transcription", *PLoS ONE* **14** (2019), e0217356.
- [18] H. Seligmann, D. D. Pollock, "The ambush hypothesis: hidden stop codons prevent off-frame gene reading", *DNA Cell. Biol.* **23** (2003), p. 701-705.
- [19] H. Seligmann, "Cost minimization of ribosomal frameshifts", *J. Theor. Biol.* **249** (2007), p. 162-167.
- [20] H. Seligmann, "The ambush hypothesis at the whole-organism level: off frame, 'hidden' stops in vertebrate mitochondrial genes increase developmental stability", *Comput. Biol. Chem.* **34** (2010), p. 80-85.
- [21] H. Seligmann, "Localized context-dependent effects of the 'ambush' hypothesis: more off-frame stop codons downstream of shifty codons", *DNA Cell. Biol.* **38** (2019), p. 786-795.
- [22] X. Wang, Q. Dong, G. Chen, J. Zhang, Y. Liu, J. Zhao, H. Peng, Y. Wang, Y. Cai, X. Wang, C. Yang, M. Lynch, 2016 "The universal genetic code, protein coding genes and genomes of

- all species were optimized for frameshift tolerance”, bioRxiv, posted Aug 25, 2016, <http://dx.doi.org/10.1101/067736>.
- [23] R. Geyer, M. A. Madany, “On the efficiency of the genetic code after frameshift mutations”, *PeerJ* **6** (2018), e4825.
- [24] G. K. Philip, S. J. Freeland, “Did evolution select a nonrandom “alphabet” of amino acids?”, *Astrobiology* **11** (2011), p. 235-240.
- [25] M. Ilardo, M. Meringer, S. Freeland, B. Rasulev, H. J. Cleaves II, “Extraordinarily adaptive properties of the genetically encoded amino acids”, *Sci. Rep.* **5** (2015), p. 9414.
- [26] E. N. Trifonov, “Consensus temporal order of amino acids and evolution of the triplet code”, *Gene* **261** (2000), p. 139-151.
- [27] E. N. Trifonov, “The triplet code from first principles”, *J. Biomol. Struct. Dyn.* **22** (2004), p. 1-11.
- [28] H. Seligmann, “Protein sequences recapitulate genetic code evolution”, *Comput. Struct. Biotech. J.* **16** (2018), p. 177-199.
- [29] D. B. F. Johnson, L. Wang, “Imprints of the genetic code in the ribosome”, *Proc. Natl. Acad. Sci. USA* **107** (2010), p. 8298-8303.
- [30] R. Root-Bernstein, “Simultaneous origin of homochirality, the genetic code and its directionality”, *Bioessays* **29** (2007), p. 689-698.
- [31] X. Han Da, H. Y. Wang, Z. L. Ji, A. F. Hu, Y. F. Zhao, “Amino acid homochirality may be linked to the origin of phosphate-based life”, *J. Mol. Evol.* **70** (2010), p. 572-582.
- [32] D. G. Blackmond, “The origin of biological homochirality”, *Phil. Trans. R. Soc. Lond. B* **366** (2011), p. 2878-2884.
- [33] R. Breslow, “The origin of homochirality in amino acids and sugars on prebiotic earth”, *Tetrahedron Lett.* **52** (2011), p. 4228-4232.
- [34] C. J. Michel, H. Seligmann, “Bijective transformation circular codes and nucleotide exchanging RNA transcription”, *Biosystems* **118** (2014), p. 39-50.
- [35] A. A. Polyansky, B. Zagrovic, “Evidence of direct complementarity interactions between messenger RNAs and their cognate proteins”, *Nucl. Acids Res.* **41** (2013), p. 8434-8443.
- [36] L. Bartonek, B. Zagrovic, “mRNA/protein sequence complementarity and its determinants: the impact of affinity scales”, *PLoS Comput. Biol.* **13** (2017), e1005648.
- [37] B. Zagrovic, L. Bartonek, A. A. Polyansky, “RNA-protein interactions in an unstructured context”, *FEBS Lett.* **592** (2018), p. 2901-2916.
- [38] N. Maizels, A. M. Weiner, “Phylogeny from function: evidence from the molecular fossil record that tRNA originated in replication, not translation”, *Proc. Natl. Acad. Sci. USA* **91** (1994), p. 6729-6734.
- [39] H. Seligmann, “Swinger RNA self-hybridization and mitochondrial non-canonical swinger transcription, transcription systematically exchanging nucleotides”, *J. Theor. Biol.* **399** (2016), p. 84-91.
- [40] P. R. Wills, C. W. Carter Jr, “Insurmountable problems of the genetic code initially emerging in an RNA world”, *Biosystems* **165** (2018), p. 155-166.
- [41] J. Demongeot, “Sur la possibilité de considérer le code génétique comme un code à enchaînement”, *Rev. Biomaths* **62** (1978), p. 61-66.
- [42] J. Demongeot, J. Besson, “Genetic-code and cyclic codes”, *C. R. Acad. Sci. III* **296** (1983), p. 807-810.
- [43] J. Demongeot, J. Besson, “Genetic code and cyclic codes II”, *C. R. Acad. Sci. III* **319** (1996), p. 520-528.
- [44] J. Demongeot, A. Elena, G. Weil, “Potential-Hamiltonian decomposition of cellular automata. Application to degeneracy of genetic code and cyclic codes III”, *C. R. Biol.* **329** (2006), p. 953-962.
- [45] J. Demongeot, A. Moreira, “A possible circular RNA at the origin of life”, *J. Theor. Biol.* **249** (2007), p. 314-324.
- [46] H. Seligmann, D. Raoult, “Unifying view of stem-loop hairpin RNA as origin of current and ancient parasitic and non-parasitic RNAs, including in giant viruses”, *Curr. Opin. Microbiol.* **31** (2016), p. 1-8.
- [47] H. Seligmann, D. Raoult, “Stem-loop RNA hairpins in giant viruses: invading rRNA-like repeats and a template free RNA”, *Front. Microbiol.* **9** (2018), p. 101.
- [48] G. Soslau, “Circular RNA (circRNA) was an important bridge in the switch from the RNA world to the DNA world”, *J. Theor. Biol.* **447** (2018), p. 32-40.
- [49] J. Salzman, C. Gawad, P. L. Wang, N. Lacayo, P. O. Brown, “Circular RNAs are the predominant transcript isoform from hundreds of human genes in diverse cell types”, *PLoS ONE* **7** (2012), e30733.
- [50] E. Lasda, R. Parker, “Circular RNAs: diversity of form and function”, *RNA* **20** (2014), p. 1829-1842.
- [51] S. P. Barrett, J. Salzman, “Circular RNAs: analysis, expression and potential functions”, *Development* **143** (2016), p. 1838-1847.
- [52] Y. Zhang, W. Xue, X. Li, J. Zhang, S. Chen, J. L. Zhang, L. Yang, L. L. Chen, “The biogenesis of nascent circular RNAs”, *Cell Rep.* **15** (2016), p. 611-624.
- [53] S. Huang, B. Yang, B. J. Chen, N. Bliim, U. Ueberham, T. Arendt, M. Janitz, “The emerging role of circular RNAs in transcriptome regulation”, *Genomics* **109** (2017), p. 401-407.
- [54] Y. Zhong, Y. Du, X. Yang, Y. Mo, C. Fan, F. Xiong, D. Ren, X. Ye, C. Li, Y. Wang, F. Wei, C. Guo, X. Wu, X. Li, Y. Li, G. Li, Z. Zeng, W. Xiong, “Circular RNAs function as ceRNAs to regulate and control human cancer progression”, *Mol. Cancer* **17** (2018), p. 79.
- [55] B. P. Nicolet, S. Engels, F. Agliarolo, E. van den Akker, M. von Lindern, M. C. Wolkers, “Circular RNA expression in human hematopoietic cells is widespread and cell-type specific”, *Nucl. Acids Res.* **46** (2018), p. 8168-8180.
- [56] N. R. Pamudurti, O. Bartok, M. Jens, R. Ashwal-Fluss, C. Stottmeister, L. Ruhe, M. Hanan, E. Wyler, D. Perez-Hernandez, E. Ramberger, S. Shenzi, M. Samson, G. Dittmar, M. Landthaler, M. Chekulaeva, N. Rajewsky, S. Kadener, “Translation of circRNA”, *Mol. Cell* **66** (2017), p. 9-21.
- [57] I. Legnini, G. Di Timoteo, F. Rossi, M. Morlando, F. Briganti, O. Sthandler, T. Santini, A. Andronache, M. Wade, P. Laneve, N. Rajewsky, I. Bozzoni, “Circ-ZNF609 is a circular RNA that can be translated and functions in myogenesis”, *Mol. Cell* **66** (2017), p. 22-37.
- [58] V. Opuu, M. Silvert, T. Simonson, “Computational design of fully overlapping coding schemes for protein pairs and triplets”, *Sci. Rep.* **7** (2017), p. 15873.
- [59] E. Faure, L. Delaye, S. Tribolo, A. Levasseur, H. Seligmann, R. M. Barthélémy, “Probable presence of an ubiquitous cryp-

- tic mitochondrial gene on the antisense strand of the cytochrome oxidase I gene”, *Biol. Direct* **6** (2011), p. 56.
- [60] H. Seligmann, “Two genetic codes, one genome: frameshifted primate mitochondrial genes code for additional proteins in presence of antisense antitermination tRNAs”, *Biosystems* **105** (2011), p. 271-285.
- [61] H. Seligmann, “An overlapping genetic code for frameshifted overlapping genes in *Drosophila* mitochondria: antisense antitermination tRNAs UAR insert serine”, *J. Theor. Biol.* **298** (2012), p. 51-76.
- [62] H. Seligmann, “Overlapping genetic code for overlapping frameshifted genes in Testudines, and *Lepidochelys olivacea* as special case”, *Comput. Biol. Chem.* **41** (2012), p. 18-34.
- [63] H. Seligmann, “Directed mutations recode mitochondrial genes: from regular to stopless genetic codes”, in *Mitochondrial DNA-New Insights* (H. Seligmann, G. Warthi, eds.), In-techOpen, 2018.
- [64] S. Itzkovitz, U. Alon, “The genetic code is nearly optimal for allowing information within protein-coding sequences”, *Genome Res.* **17** (2007), p. 405-412.
- [65] X. Wang, X. Wang, G. Chen, J. Zhang, Y. Liu, C. Yang, “The shiftability of protein coding genes: the genetic code was optimized for frameshift tolerating”, *PeerJ Preprints* **3** (2015), e806v1.
- [66] R. Geyer, M. A. Madany, “On the efficiency of the genetic code after frameshift mutations”, *PeerJ* **6** (2018), e4825.
- [67] C. R. Woese, “Evolution of the genetic code”, *Die Naturwissenschaften* **60** (1973), p. 447-459.
- [68] R. Wolfenden, L. Andersson, P. M. Cullis, C. C. B. Southgate, “Affinities of amino acid side chains for solvent water”, *Biochemistry* **20** (1981), p. 849-855.
- [69] D. C. Mathew, Z. Luthey-Schulten, “On the physical basis of the amino acid polar requirement”, *J. Mol. Evol.* **66** (2008), p. 519-528.
- [70] C. Chothia, “The nature of accessible and buried surfaces in proteins”, *J. Mol. Biol.* **105** (1976), p. 1-12.
- [71] J. Janin, “Surface and inside volumes in globular proteins”, *Nature* **277** (1979), p. 491-492.
- [72] R. Wolfenden, P. M. Cullis, C. C. B. Southgate, “Water, protein folding, and the genetic code”, *Science* **206** (1979), p. 575-577.
- [73] A. Radzicka, R. Wolfenden, “Comparing polarities of the amino acids: side-chain distribution coefficients between the vapor phase, cyclohexane, 1-octanol, and neutral aqueous solution”, *Biochemistry* **27** (1988), p. 1664-1670.
- [74] R. Wolfenden, “Experimental measures of amino acid hydrophobicity and the topology of transmembrane and globular proteins”, *J. Gen. Physiol.* **129** (2007), p. 357-362.
- [75] M. Di Giulio, “The late stage of genetic code structuring took place at a high temperature”, *Gene* **261** (2000), p. 189-195.
- [76] R. Wolfenden, C. A. Lewis, Y. Yuan, C. W. Carter Jr, “Temperature dependence of amino acid hydrophobicities”, *Proc. Natl. Acad. Sci. USA* **112** (2015), p. 7484-7488.
- [77] M. Di Giulio, “The universal ancestor was a thermophile or a hyperthermophile: tests and further evidence”, *J. Theor. Biol.* **221** (2003), p. 425-436.
- [78] M. Di Giulio, “The universal ancestor and the ancestor of bacteria were hyperthermophiles”, *J. Mol. Evol.* **57** (2003), p. 721-730.
- [79] S. Akanuma, Y. Nakajima, S. Yokobori, M. Kimura, N. Nemoto, T. Mase, K. Miyazono, M. Tanokura, A. Yamagishi, “Experimental evidence for the thermophilicity of ancestral life”, *Proc. Natl. Acad. Sci. USA* **110** (2013), p. 11067-11072.
- [80] N. H. Sleep, “Geological and geochemical constraints on the origin and evolution of life”, *Astrobiology* **18** (2018), p. 1199-1219.
- [81] J. E. M. Hornos, Y. M. M. Hornos, “Algebraic model for the evolution of the genetic code”, *Phys. Rev. Lett.* **71** (1993), p. 4401-4404.
- [82] S. L. Miller, “Production of amino acids under possible primitive earth conditions”, *Science* **117** (1953), p. 528-529.
- [83] M. J. Dufton, “Genetic code synonym quotas and amino acid complexity: cutting the cost of proteins?”, *J. Theor. Biol.* **187** (1997), p. 165-173.
- [84] J. T. F. Wong, “A co-evolution theory of the genetic code”, *Proc. Natl. Acad. Sci. USA* **72** (1975), p. 1909-1912.
- [85] R. C. Guimarães, “Self-referential encoding on modules of anticodon pairs - roots of the biological flow system”, *Life (Basel)* **7** (2017), p. 16.
- [86] C. Francklyn, K. Musier-Forsyth, P. Schimmel, “Small RNA helices as substrates for aminoacylation and their relationship to charging of transfer RNAs”, *Euro. J. Biochem.* **206** (1992), p. 315-321.
- [87] C. W. Carter Jr, R. Wolfenden, “tRNA acceptor-stem and anticodon bases form independent codes related to protein folding”, *Proc. Natl. Acad. Sci. USA* **112** (2015), p. 7489-7494.
- [88] C. W. Carter Jr, P. R. Wills, “Hierarchical groove discrimination by Class I and II aminoacyl-tRNA synthetases reveals a palimpsest of the operational RNA code in the tRNA acceptor-stem bases”, *Nucl. Acids Res.* **46** (2018), p. 9667-9683.
- [89] C. W. Carter Jr, R. Wolfenden, “Acceptor-stem and anticodon bases embed amino acid chemistry into tRNA”, *RNA Biol.* **13** (2016), p. 145-151.
- [90] H. Hartman, “Speculations on the evolution of the genetic code”, *Orig. Life Biosph.* **6** (1975), p. 423-427.
- [91] H. Hartman, “Speculations on the origin and evolution of metabolism”, *J. Mol. Evol.* **4** (1975), p. 359-370.
- [92] H. Hartman, “Speculations on the evolution of the genetic code II”, *Orig. Life Biosph.* **9** (1978), p. 133-136.
- [93] H. Hartman, “Speculations on the origin of the genetic code”, *J. Mol. Evol.* **40** (1995), p. 541-544.
- [94] E. N. Trifonov, “Elucidating sequence codes: three codes for evolution”, *Ann. N.Y. Acad. Sci.* **870** (1999), p. 330-338.
- [95] M. Yarus, “The genetic code and RNA-amino acid affinities”, *Life (Basel)* **7** (2017), p. 13.
- [96] J. Demongeot, H. Seligmann, “Spontaneous evolution of circular codes in theoretical minimal RNA rings”, *Gene* **705** (2019), p. 95-102.
- [97] H. Seligmann, “First arrived, first served: competition between codons for codon-amino acid stereochemical interactions determined early genetic code assignments”, *Naturwissenschaften* **107** (2020), p. 20.
- [98] J. Demongeot, H. Seligmann, “Pentamers with non-

- redundant frames: bias for natural circular code codons”, *J. Mol. Evol.* **88** (2020), p. 194-201.
- [99] J. Demongeot, H. Seligmann, “More pieces of ancient than recent theoretical minimal proto-tRNA-like RNA rings in genes coding for tRNA synthetases”, *J. Mol. Evol.* **87** (2019), p. 1-23.
- [100] J. Demongeot, H. Seligmann, “Bias for 3'-dominant codon directional asymmetry in theoretical minimal RNA rings”, *J. Comput. Biol.* (2019), doi:10.1089/cmb.2018.0256.
- [101] J. Demongeot, H. Seligmann, “Theoretical minimal RNA rings recapitulate the order of the genetic code's codon-amino acid assignments”, *J. Theor. Biol.* **471** (2019), p. 108-116.
- [102] J. Demongeot, H. Seligmann, “Theoretical minimal RNA rings designed according to coding constraints mimic deamination gradients”, *Naturwissenschaften* **106** (2019), p. 44.
- [103] J. Demongeot, H. Seligmann, “The uroboros theory of life's origin: 22-nucleotide theoretical minimal RNA rings reflect evolution of genetic code and tRNA-rRNA translation machineries”, *Acta Biotheor.* **67** (2019), p. 273-297.
- [104] J. Demongeot, H. Seligmann, “Evolution of tRNA into rRNA secondary structures”, *Gene Rep.* (2019), 100483.
- [105] J. Demongeot, H. Seligmann, “Accretion history of large ribosomal subunits deduced from theoretical minimal RNA rings is congruent with histories derived from phylogenetic and structural methods”, *Gene* **738** (2020), 144436.
- [106] J. Demongeot, H. Seligmann, “Comparisons between small ribosomal RNA and theoretical minimal RNA ring secondary structures confirm phylogenetic and structural accretion histories”, *Sci. Rep.* **10** (2020), p. 7693.
- [107] J. Demongeot, H. Seligmann, “RNA rings strengthen hairpin accretion hypotheses for tRNA evolution: a reply to commentaries by Z. F. Burton and M. Di Giulio”, *J. Mol. Evol.* **88** (2020), p. 243-252.
- [108] J. Demongeot, H. Seligmann, “The primordial tRNA acceptor stem code from theoretical minimal RNA ring clusters”, *BMC Genet.* **21** (2020), p. 7.
- [109] J. Demongeot, H. Seligmann, “Why is AUG the start codon?: theoretical minimal RNA rings: maximizing coded information biases 1st codon for the universal initiation codon AUG”, *Bioessays* (2020), e1900201.
- [110] H. Seligmann, J. Demongeot, “Codon directional asymmetry suggests swapped prebiotic 1st and 2nd codon positions”, *Int. J. Mol. Sci.* **21** (2020), e347.
- [111] J. Demongeot, H. Seligmann, “Deamination gradients within codons after 1 ↔ 2 position swap predict amino acid hydrophobicity and parallel β-sheet conformational preference”, *Biosystems* **191–192** (2020), 104116.
- [112] A. Soma, A. Onodera, J. Sugahara, A. Kanai, N. Yachie, M. Tomita, F. Kawamura, Y. Sekine, “Permuted tRNA genes expressed via a circular RNA intermediate in *Cyanidioschyzon merolae*”, *Science* **318** (2007), p. 450-453.
- [113] T. Pan, R. R. Gutell, O. C. Uhlenbeck, “Folding of circularly permuted transfer RNAs”, *Science* **254** (1991), p. 1361-1364.
- [114] A. Gutfraind, A. Kempf, “Error-reducing structure of the genetic code indicates code origin in non-thermophile organisms”, *Orig. Life Evol. Biosph.* **38** (2008), p. 75-85.