



ELSEVIER

Contents lists available at SciVerse ScienceDirect

Comptes Rendus Chimie

www.sciencedirect.com



Full paper/Mémoire

Coinage metal complexes with N'NN' and SeNSe ligands – Pincer vs bridging coordination pattern

Adriana Ilie^a, Antonio Laguna^b, Cristian Silvestru^{a,*}^a Department of Inorganic Chemistry, Faculty of Chemistry & Chemical Engineering, Babes-Bolyai University, RO-400028 Cluj-Napoca, Romania^b Departamento de Química Inorgánica, Instituto Síntesis Química y Catálisis Homogénea (ISQCH), Universidad de Zaragoza-CSIC, 50009 Zaragoza, Spain

ARTICLE INFO

Article history:

Received 28 February 2012

Accepted after revision 14 May 2012

Available online 15 June 2012

Keywords:

Ag(I) complexes

Au(III) complex

Cu(I) complexes

N'NN' ligand

SeNSe ligand

X-ray diffraction

ABSTRACT

The reactions of the ligands 2,6-(Me₂NCH₂)₂C₅H₃N (N'NN') (**1**) and 2,6-(PhSeCH₂)₂C₅H₃N (SeNSe) (**4**) with different coinage metal starting materials gave 1:1, 2:1 or 1:2 metal-to-ligand species, i.e. [Ag(N'NN'){O(O)CCF₃}] (**2**), {[Ag(PPh₃)₂(N'NN')](OTf)₂} (**3**), [Au(SeNSe)Cl]Cl₂ (**5**), [Ag(PPh₃)(SeNSe)](OTf) (**6**), [Cu(MeCN)(SeNSe)](PF₆) (**7**) or [Cu(SeNSe)₂](PF₆) (**8**). The new compounds were investigated by IR, multinuclear NMR spectroscopies as well as mass spectrometry. In most cases, the ligands **1** and **4** act as pincer ligands. An attempt to grow single crystals of **2** gave an unexpected result. The crystal investigated by X-ray diffraction proved to be a polynuclear species, [Ag₄(N'NN'){O(O)CCF₃}₄(EtOH)]_n (**2a**), which contains an unusual, bimetallic triconnective coordination pattern of the N'NN' ligand. Two tetranuclear [Ag₄(N'NN'){O(O)CCF₃}₄(EtOH)] units form centrosymmetric dimers further associated into a polymer which contains four different coordination environments around silver atoms. The complex **3**, in which the ligand also exhibits a bimetallic triconnective pattern, shows an intense, long-lived luminescence in the solid state with emission energies in the green region of the visible spectrum.

© 2012 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

La réaction de ligands 2,6-(Me₂NCH₂)₂C₅H₃N (N'NN') (**1**) et 2,6-(PhSeCH₂)₂C₅H₃N (SeNSe) (**4**) avec les métaux de l'XI-ème groupe a donné des différentes espèces métal-ligand 1:1, 2:1 ou 1:2, i.e. [Ag(N'NN'){O(O)CCF₃}] (**2**), {[Ag(PPh₃)₂(N'NN')](OTf)₂} (**3**), [Au(SeNSe)Cl]Cl₂ (**5**), [Ag(PPh₃)(SeNSe)](OTf) (**6**), [Cu(MeCN)(SeNSe)](PF₆) (**7**) ou [Cu(SeNSe)₂](PF₆) (**8**). Les nouveaux composés ont été investigués par IR, spectroscopie RMN multinucléaire et par la spectrométrie de masse. Dans la plupart des cas, les ligands **1** et **4** agissent comme une pince ligand. La tentative d'obtenir de monocristaux du **2** a donné un résultat inattendu. Le cristal investigué par diffraction de rayons X a relevé la formation d'une espèce polynucléaire [Ag₄(N'NN'){O(O)CCF₃}₄(EtOH)]_n (**2a**), qui contient d'une manière surprenante un modèle de coordination bimétallique triconnective du ligand N'NN'. Deux unités tétranucléaires [Ag₄(N'NN'){O(O)CCF₃}₄(EtOH)] forment des dimères centrosymétriques associés ultérieurement dans un polymère qui contient quatre modalités de coordination différentes au niveau des atomes d'argent. Le complexe **3**, dans lequel le ligand présente également un modèle bimétallique triconnective, montre une intense, longue durée de vie de luminescence à l'état solide, avec des énergies d'émission dans la région verte du spectre visible.

© 2012 Académie des sciences. Publié par Elsevier Masson SAS. Tous droits réservés.

* Corresponding author.

E-mail address: cristi@chem.ubbcluj.ro (C. Silvestru).

1. Introduction

The coordination chemistry of 2,6-(Me₂NCH₂)₂C₅H₃N (**1**, N'NN') and related ligands was investigated mainly in relation to the potential application in catalysis of the corresponding metal complexes [1]. Several metal ions were used to investigate the coordination ability of **1**, e.g. Fe^{II} [1f,1g], Ru^{II} [1a–d,g,2], Ru^{III} [1d], Ru^{IV} [3], Pd^{II} [4], Cu^I [5b], Cu^{II} [5], U^{VI} [6], Zn^{II} [7]. By contrast, the ligand 2,6-(PhSeCH₂)₂C₅H₃N (**4**, SeNSe) (Scheme 1) was recently reported [8a] and only few of its Pd^{II} and Ru^{II} complexes were described as highly active catalysts for Heck coupling (C–C) reaction [8a] and oxidation of primary and secondary alcohols [8b], respectively. In most cases both ligands **1** and **4** were found to act as monometallic triconnective (“pincer”, Scheme 1, A) units, i.e. with both the sp²-hybridized nitrogen atom of the central pyridine ring and the donor atoms from two flexible pendant arms coordinated to the same metal atom. In only one case a different coordination pattern was described, i.e. the cation of the copper(I) complex [Cu(N'NN')₂][CuCl₂] contains both ligand units chelating the metal atom through the pyridine nitrogen atom and the nitrogen of one pendant arm, while the other pendant arm is not coordinated to a metal centre (Scheme 1, B). This suggests that in addition to a pincer pattern such ligands might act as a bridge, the “free” pendant arm being involved in coordination to a second metal atom (Scheme 1, C) and thus providing the possibility for metal–metal interactions. For coinage metals such metalophilic interactions can influence the properties of the complexes, e.g. the optical properties [9].

With respect to other nitrogen donor ligands related to **1**, including macrocycles containing the fragment 2,6-substituted pyridine, in addition to metals as Mn^{II} [1g,10], Fe^{II} [1g,11], Ru^{II}, Co^{II} [1g], Cu^{II} [5b,11,12], Zn^{II} [11,13,14] or Pb^{II} [13b], several Ag^I complexes were investigated and their structure reported [10–13,15].

We describe herein the synthesis and spectroscopic characterization of some new coinage metal complexes


containing the potential tridentate ligands N'NN' (**1**) and SeNSe (**4**) as well as the structure of the tetranuclear silver(I) species [Ag₄(N'NN'){O(O)CCF₃}₄(EtOH)] which shows intermetallic contacts and a bimetallic triconnective N'NN' ligand.

2. Results and discussion


The potential tridentate ligand 2,6-(Me₂NCH₂)₂C₅H₃N (**1**, N'NN') was prepared according to a published method [4b], while for the synthesis of the related selenium-containing ligand 2,6-(PhSeCH₂)₂C₅H₃N (**4**, SeNSe) a slightly modified procedure [8a] was used. Reactions of **1** with [AgO(O)CCF₃] (1:1 molar ratio) and [Ag(PPh₃)(OTf)] (1:2 molar ratio) in anhydrous CH₂Cl₂ afforded the isolation of compounds [Ag(N'NN'){O(O)CCF₃}] (**2**) and [Ag(PPh₃)₂(N'NN')](OTf)₂ (**3**) as light yellow or white-cream solids, respectively (Scheme 2). Both complexes were stable to air and moisture for long periods of time.

The organoselenium(II) derivative **4** was reacted with H[AuCl₄].3H₂O, [Ag(PPh₃)(OTf)] or [Cu(MeCN)₄]PF₆ to yield 1:1 complexes [Au(SeNSe)Cl]Cl₂ (**5**), [Ag(PPh₃)(SeNSe)](OTf) (**6**) and [Cu(MeCN)(SeNSe)](PF₆) (**7**) in good yields. When **4** was treated with [Cu(MeCN)₄]PF₆ in 2:1 molar ratio the complex [Cu(SeNSe)₂](PF₆) (**8**) was isolated (Scheme 3). All complexes containing **4** as a ligand were isolated as stable yellow (**5**, **7** and **8**) or white solids (**6**). Analytical and spectroscopic data (IR and multinuclear NMR spectroscopies, mass spectrometry) were consistent with the anticipated formulas for the new complexes. The assignment of the ¹H and ¹³C resonances in the room temperature NMR spectra (recorded in CDCl₃ solutions) was based on the 2D experiments.

The molecular ions for complexes **2**, **3** and **5–8** were not observed in the MALDI(+) or ESI(+) mass spectra, thus suggesting easy splitting of the corresponding anionic group. The MALDI(+) spectra of **2** and **5** show the fragment [Ag(N'NN')⁺] (*m/z* 301) and [Au(SeNSe)⁺] (*m/z* 616) as base peaks. For the silver complexes **3** and **6** the base peak



Scheme 1. Ligands **1** and **4** and potential coordination patterns.

Scheme 2. Synthesis of silver complexes **2** and **3**.


corresponds to $[\text{Ag}(\text{PPh}_3)_2]^+$ (m/z 631), but for both complexes the spectra contain also the peaks assigned to $[\text{Ag}(\text{PPh}_3)(\text{N}'\text{NN}')^+]$ (m/z 562) and $[\text{Ag}(\text{PPh}_3)(\text{SeNSe})^+]$ (m/z 788), respectively. The ESI(+) mass spectrum of the copper complex **8** exhibit the metal-containing fragments $[\text{Cu}(\text{SeNSe})_2]^+$ (m/z 901) and $[\text{Cu}(\text{SeNSe})]^+$ (m/z 482), the later one being the base peak for the complex **7** too.

The IR spectrum of **2** shows strong to medium bands at 1679 , 1666 cm^{-1} [$\nu_{\text{as}}(\text{CO}_2)$] and 1457 cm^{-1} [$\nu_{\text{s}}(\text{CO}_2)$]. Although there is no clear difference between the Δ values [$\Delta = \nu_{\text{as}}(\text{CO}_2) - \nu_{\text{s}}(\text{CO}_2)$] calculated for **3** (av. 215 cm^{-1}) vs. the ionic $\text{NH}_4(\text{O}_2\text{CCF}_3)$ [1667 cm^{-1} for $\nu_{\text{asym}}(\text{CO}_2)$ and 1465 cm^{-1} for $\nu_{\text{sym}}(\text{CO}_2)$; $\Delta = 202\text{ cm}^{-1}$] [16], it is difficult to draw structural conclusions based only on the IR data. Apparently, the IR data seem to be consistent with a triflate anion covalently bound and ionic in nature in the silver complexes **3** and **6**, respectively, in solid state [17]. However, in the absence of a single-crystal X-ray diffraction study it is difficult to clearly decide between these possibilities.

The ^1H and ^{13}C NMR spectra for complexes **2** and **3** are very similar, with respect to the aliphatic and aromatic resonances, to those of the free ligand **1**. The sharp singlet resonances observed for the protons and carbon atoms of

the methylene and methyl groups suggest the equivalence of the pendant arms of the N'NN' ligand unit, at room temperature, on the NMR time scale. For complex **3** this suggests a dynamic behavior, the silver atoms of a cation unit being alternatively coordinated by two nitrogen atoms. This dynamic process is also supported by the presence of only one set of broader ^{13}C resonances for the phenyl groups of the PPh_3 ligands (phosphorus-carbon coupling observed, see Experimental Section). In addition to the expected ^{13}C resonances for the neutral ligands, the spectra of both compounds **2** and **3** contain resonances for the anions, with a quartet pattern due to carbon-fluorine coupling, i.e. δ 117.53 [$^1J(\text{CF})$ 291.3 Hz, CF_3] and 162.36 [$^2J(\text{CF})$ 34.8 Hz, $-\text{C}(\text{O})\text{O}-$] for **2**, and δ 120.57 [q, $^1J(\text{CF})$ 320.8 Hz, CF_3] for **3**. The ^{19}F NMR spectrum of **3** exhibits only one singlet resonance (δ -77.9 ppm), consistent with the presence of only one species in solution.

Similar considerations can be drawn out from the ^1H and ^{13}C NMR spectra for metal complexes of the selenium-containing ligand **4**, although the ^{77}Se resonance could be observed only for the free ligand. Only one set of ^1H and ^{13}C resonances was observed for the two pendant arms (e.g. singlet resonances for all methylene protons or carbons). Also the protons in positions 3 and 5 of the pyridine

Scheme 3. Synthesis of metal complexes **5–8**.

ring gave one doublet resonance in the aromatic region. This suggests either a symmetrical (Se,N,Se-monometallic triconnective) coordination in the case of complexes **5–7** or a fast dynamic process, at room temperature, for compound **8**, with the copper atom being coordinated by the nitrogen of the pyridine ring and alternating selenium atom from pendant arms [Cu(NSe)₂ core]. For complexes **6** and **7** the ¹H and ¹³C NMR spectra exhibit, in addition to the resonances for ligand **4**, further signals for the PPh₃ and MeCN molecules coordinated to the metal atom. The presence of the OTf[−] and PF₆[−] anions was proved by ¹⁹F and ³¹P NMR spectra; the presence of only one resonance with expected multiplicity pattern being consistent with the presence of only one species in solution.

In case of complexes **3** and **6**, which both contain PPh₃ coordinated to the metal atom, the ³¹P NMR spectroscopy was a very useful tool for the investigation of the coordination environment of the silver. For compound **3**, the ³¹P NMR spectrum, recorded at room temperature, exhibits two broad resonances at δ 11.9 and 16.0 ppm. This might be due to a dissociative process involving cleavage of N-Ag(PPh₃), as mentioned above. Such dissociative processes are well-documented in the case of phosphine complexes of coinage metals [18]. At −55 °C the ³¹P NMR spectrum of **3** shows two resolved resonances, each of them with a pattern of doublet of doublets due to the couplings with ¹⁰⁷Ag and ¹⁰⁹Ag isotopes, thus indicating the presence of two different environments for silver-phosphine cores. The coordination number (CN) around a silver atom in a silver-phosphine species can be estimated on the basis of the magnitude of the silver-phosphorus coupling constant. As mentioned in literature, the ¹J(¹⁰⁷Ag,P) coupling constants are inversely proportional to the coordination number of the silver atom [19] and increase with decreasing of Ag-P distances [20]. Thus, using the magnitude of the observed coupling constants the resonance at δ 11.0 [¹J(P,¹⁰⁷Ag) 511.3 Hz] was assigned to a Ag(PPh₃) fragment coordinated by two nitrogen atoms [N₂Ag(PPh₃) core, CN = 3], while the resonance at δ 14.4 [¹J(P,¹⁰⁷Ag) 627.6 Hz] was assigned to a Ag(PPh₃) fragment coordinated by one nitrogen atom [NAg(PPh₃) core, CN = 2], respectively. On the basis of similar considerations, it can be assumed that at −55 °C the ³¹P NMR spectrum of **6** shows resonances, with doublet of doublet pattern, which can be assigned to two different silver-containing species in solution. The resonance at δ 7.9 [¹J(P,¹⁰⁷Ag) = 398.0 Hz] can be assigned to a tetracoordinated species in which the Ag(PPh₃) fragment is coordinated by all donor atoms of a SeNSe ligand, while the less resolved resonance at δ 11.5 [¹J(P,¹⁰⁷Ag) = 551.6 Hz] might be due to a tricoordinated species in which a silver-selenium bond was dissociated.

The complex **3** exhibits an intense, long-lived luminescence in the solid state with emission energies in the green region of the visible spectrum. It is emissive in the solid state both at room temperature and in frozen solution (an intermediate situation between solid and solution state), as shown in Fig. 1. Emissions in the solid state appear near 488 nm, while excitation near 425 nm. These maxima appear shifted to higher energies in dichloromethane




Fig. 1. Emission and excitation spectra for compound **3** in solid state at room temperature.

frozen solutions, i.e. 525 nm for emission and 475 nm for excitation, respectively. In all cases only one band is observed.

In an attempt to grow single crystals suitable for X-ray diffraction studies compound **2** was dissolved in CH₂Cl₂ and hexane was slowly added (1:5, v/v). The X-ray diffraction analysis of one of the few colorless crystals obtained in this way revealed the molecular structure of an unexpected polynuclear silver species, [Ag₄(N'NN') {O(O)CCF₃}₄(EtOH)]_n (**2a**). The presence of the ethanol molecule can be explained by presence of this solvent in the methylene chloride used to grow the crystals. No attempts to prepare this polymer in a higher yield were carried out so far. Selected interatomic distances and angles are listed in Table 1.

The crystal of **2a** contains a polymer arrangement built from repeating tetranuclear units of the type depicted in Fig. 2. In such a tetranuclear unit the N'NN' ligand act as a bimetallic triconnective unit, bridging Ag1 [Ag(1)–N(1) 2.334(7) Å, Ag(1)–N(2) 2.416(7) Å] and Ag3 atoms [Ag(3)–N(3) 2.300(7) Å]. The Ag–N bond distances are slightly different, nevertheless the average of these bond distances compare well with those found in the literature [10–13,15]. The distance between the silver atoms bridged by the N'NN' ligand is too long [Ag(1)···Ag(3) 3.51(4) Å; cf. sums of the corresponding van der Waals radii, Σr_{vdw}(Ag,Ag) 3.4 Å [21]] to suggest any metal-metal interaction. However, silver-silver contacts are established with a third metal atom, Ag2 [Ag(1)–Ag(2) 3.075(5) Å, Ag(2)–Ag(3) 3.009(4) Å]. The metal-metal interactions in the resulting angular Ag₃ system are supported by three bridging trifluoroacetate units: two of them between Ag2 and Ag3 atoms, while the third one bridge Ag2 and Ag1 atom (coordinated by two nitrogen atoms of the N'NN' ligand). The oxygen atoms O2 and O5, from two trifluoroacetate ligands bridging different silver atoms of the Ag₃ system, are involved in coordination to the fourth metal atom of the tetranuclear fragment [Ag(4)–O(2) 2.684(8) Å, Ag(4)–O(5) 2.496(7) Å]. As a result these trifluoroacetate ligands exhibit each a trimetallic triconnective pattern. The Ag₂O₂ four-membered ring thus formed is not planar, but folded [dihedral angle O(2)Ag(2)O(5)/O(2)Ag(4)O(5) 31.5°].

Table 1
Selected bond distances [Å] and angles [°] for compound **2a**.^a

Ag(1)–O(1)	2.235(7)	Ag(3)–O(4)	2.16(2)
Ag(1)–N(1)	2.334(7)	Ag(3)–O(6)	2.693(6)
Ag(1)–N(2)	2.416(7)	Ag(3)–O(6a') ^a	2.595(6)
Ag(1)–Ag(2)	3.075(5)	Ag(3)–N(3)	2.300(7)
Ag(2)–O(2)	2.346(7)	Ag(4)–O(2)	2.684(8)
Ag(2)–O(3)	2.23(2)	Ag(4)–O(5)	2.496(7)
Ag(2)–O(5)	2.473(7)	Ag(4)–O(7)	2.234(7)
Ag(2)–O(9)	2.485(10)	Ag(4)–O(8)	2.240(8)
Ag(2)–Ag(3)	3.009(4)	Ag(4)–Ag(4a) ^a	2.889(4)
C(13)–O(1)	1.279(11)	C(17)–O(5)	1.254(9)
C(13)–O(2)	1.264(10)	C(17)–O(6)	1.243(8)
C(15)–O(3)	1.22(2)	C(19)–O(7)	1.212(10)
C(15)–O(4)	1.32(2)	C(19)–O(8a) ^a	1.252(10)
C(21)–O(9)	1.371(15)		
N(1)–Ag(1)–N(2)	76.3(3)	N(3)–Ag(3)–O(6a')	101.0(3)
O(1)–Ag(1)–N(1)	129.3(2)	O(6a')–Ag(3)–O(4)	78.8(6)
O(1)–Ag(1)–N(2)	129.0(3)	O(4)–Ag(3)–Ag(2)	75.4(6)
Ag(2)–Ag(1)–N(1)	116.8(2)	Ag(2)–Ag(3)–N(3)	107.70(19)
Ag(2)–Ag(1)–N(2)	130.6(2)	N(3)–Ag(3)–O(4)	173.2(7)
Ag(2)–Ag(1)–O(1)	81.5(2)	O(6a')–Ag(3)–Ag(2)	143.27(14)
		O(6)–Ag(3)–N(3)	99.3(2)
		O(6)–Ag(3)–O(4)	87.5(7)
		O(6)–Ag(3)–O(6a')	78.4(2)
		O(6)–Ag(3)–Ag(2)	74.8(1)
O(9)–Ag(2)–Ag(3)	173.7(2)	O(7)–Ag(4)–O(8)	162.8(2)
O(9)–Ag(2)–O(2)	82.2(3)	O(7)–Ag(4)–O(2)	95.9(3)
O(9)–Ag(2)–O(3)	95.7(7)	O(7)–Ag(4)–O(5)	98.7(2)
O(9)–Ag(2)–O(5)	100.5(3)	O(7)–Ag(4)–Ag(4a)	83.34(18)
Ag(3)–Ag(2)–O(2)	100.08(19)	O(8)–Ag(4)–O(2)	95.4(2)
Ag(3)–Ag(2)–O(3)	79.2(7)	O(8)–Ag(4)–O(5)	96.6(2)
Ag(3)–Ag(2)–O(5)	85.61(19)	O(8)–Ag(4)–Ag(4a)	79.51(18)
O(2)–Ag(2)–O(3)	146.1(7)	O(2)–Ag(4)–O(5)	76.9(2)
O(2)–Ag(2)–O(5)	83.9(2)	O(2)–Ag(4)–Ag(4a)	125.03(15)
O(3)–Ag(2)–O(5)	129.3(6)	O(5)–Ag(4)–Ag(4a)	157.88(14)
O(2)–Ag(2)–Ag(1)	73.69(18)		
O(3)–Ag(2)–Ag(1)	74.2(7)		
Ag(3)–Ag(2)–Ag(1)	70.63(13)		
O(5)–Ag(2)–Ag(1)	143.35(14)		
O(9)–Ag(2)–Ag(1)	104.7(3)		
C(2)–N(1)–Ag(1)	111.8(5)	C(6)–N(1)–Ag(1)	125.6(5)
C(7)–N(2)–Ag(1)	102.0(5)	C(10)–N(3)–Ag(3)	113.0(4)
C(8)–N(2)–Ag(1)	115.2(6)	C(11)–N(3)–Ag(3)	108.9(5)
C(9)–N(2)–Ag(1)	111.1(6)	C(12)–N(3)–Ag(3)	108.3(5)
C(13)–O(1)–Ag(1)	117.4(6)	C(17)–O(5)–Ag(2)	121.1(5)
C(13)–O(2)–Ag(2)	120.9(6)	C(17)–O(5)–Ag(4)	131.9(5)
		C(17)–O(6)–Ag(3)	123.5(6)
		C(17)–O(6)–Ag(3a')	134.8(5)
C(15)–O(3)–Ag(2)	120.3(15)	C(19)–O(7)–Ag(4)	122.2(6)
C(15)–O(4)–Ag(3)	128.0(13)	C(19)–O(8a)–Ag(4)	125.9(6)
C(21)–O(9)–Ag(2)	132.3(11)		
Ag(2)–O(2)–Ag(4)	93.8(2)	Ag(3)–O(6)–Ag(3a')	101.6(2)
Ag(2)–O(5)–Ag(4)	95.6(2)		
O(1)–C(13)–O(2)	130.5(8)	O(5)–C(17)–O(6)	127.2(8)
O(3)–C(15)–O(4)	123.0(16)	O(7)–C(19)–O(8a)	128.9(9)

^a Symmetry equivalent atoms (1 – x, 2 – y, –z) and (2 – x, 1 – y, –z) are given by "prime" and "a", respectively.

However this folding is not enough to bring in contact Ag2 and Ag4 atoms [Ag(2)⋯Ag(4) 3.679(5) Å].

This Ag4 atom is part of the central [Ag₂{O(O)CCF₃}]₂ system of the centrosymmetric dimer built from two




Fig. 2. ORTEP representation at 20% probability and atom numbering scheme for the tetranuclear unit in **2a** (hydrogen and fluorine atoms are omitted for clarity).

tetranuclear fragments (Fig. 3). The transannular Ag(4)–Ag(4a) of 2.889(4) Å, established between the metal atoms of the central Ag₂O₄C₂ eight-membered ring, suggests a strong silver–silver interaction. This distance is considerably shorter than observed in the similar Ag₂O₄C₂ eight-membered rings of the polymer [Ag₂{O(O)CCF₃}]_n [Ag–Ag 2.973(1) Å] [22].

An ethanol molecule is coordinated through its oxygen to the Ag2 atom, the vector of this bond being placed in *trans* to the Ag3 atom [O(9)–Ag(2)–Ag(3) 173.7(2)°]. A hydrogen bond between the hydroxyl group of the alcohol molecule and the O7 atom is suggested by the distance O(9)⋯O(7) 2.93 Å.

The dimer units are doubly connected into a polymer through Ag(3)–O(6a') [2.595(6) Å] interactions, the resulting Ag₂O₂ four-membered ring being in this case planar (Fig. 4).

Taking into account all silver–nitrogen [range 2.300(7)–2.416(7) Å], silver–oxygen [range 2.16(2)–2.693(6) Å] and silver–silver [range 2.889(4)–3.075(5) Å] bonds and interactions *per* metal atom, four different overall coordination geometry around the silver atoms can be considered:

- distorted tetrahedral Ag[N₂OAg] core for Ag1;
- distorted capped-trigonal bipyramidal Ag[O₄Ag₂] core for Ag2, with Ag1 capping the face described by Ag3, O2 and O3 atoms;
- distorted square pyramidal Ag[NO₃Ag] core for Ag3, with O6 in apical position;
- distorted trigonal bipyramidal Ag[O₄Ag] core for Ag4, with O7 and O8 atoms in axial positions [O(7)–Ag(4)–O(8) 162.8(2)°].




Fig. 3. View of a centrosymmetric dimer built from tetranuclear units through bridging trifluoroacetate anions in the crystal of **2a** (hydrogen atoms are omitted for clarity) [symmetry equivalent atoms (2-x, 1-y, -z) are given by "a"].




Fig. 4. View of the polymer arrangement built from dimer units through Ag...O interactions in the crystal of **2a** (hydrogen atoms are omitted for clarity) [symmetry equivalent atoms (2-x, 1-y, -z) and (1-x, 2-y, -z) are given by "a" and "prime", respectively].

3. Conclusions

As expected for most of the new coinage metal complexes reported here 2,6-(Me₂NCH₂)₂C₅H₃N (N'NN') (**1**) and 2,6-(PhSeCH₂)₂C₅H₃N (SeNSe) (**4**) act as pincer (monometallic triconnective) ligands. However, when an appropriate metal-to-ligand ratio was used a bridging (e.g. bimetallic triconnective) pattern can be achieved as in complex [(Ag(PPH₃)₂(N'NN'))(OTf)₂] (**3**). This compound was found to exhibit an intense luminescence in the solid state which might be due to potential metal-metal interaction. On the other hand in the copper(I) complex [Cu(SeNSe)₂](PF₆)₂ (**8**) the ligand appears to have a free donor atom still available for coordination to a metal centre. In this case a reaction of **8** with an appropriate coordinatively unsaturated metal-containing moiety can provide new homo- or heterometallic polynuclear species. The ability of such ligands to act as bridging units between metal atoms was also supported by the unexpected

polynuclear species, [Ag₄(N'NN'){O(O)CCF₃}]₄(EtOH)_n (**2a**), which contains metal-metal contacts and four different coordination environments around silver atoms.

4. Experimental

4.1. General considerations

All manipulations of air- and moisture-sensitive compounds were carried out in argon atmosphere using standard Schlenk techniques. Solvents were dried using standard procedures and freshly distilled under argon prior to use. Chemicals of commercial grade (Aldrich or Merck) were used as received. The ligand 2,6-(Me₂NCH₂)₂C₅H₃N (**1**, N'NN') was prepared according to a published procedure [4b]. The ¹H, ¹³C, ¹⁹F and ³¹P NMR spectra were recorded at room temperature, in CDCl₃, on Bruker Avance 300 (for **4**) or Avance 400 spectrometers. The ¹H and ¹³C chemical shifts are reported in ppm relative

to the residual peak of the solvent as internal standard [ref. CHCl_3 : $\delta(^1\text{H})$ 7.26, $\delta(^{13}\text{C})$ 77.0 ppm]. The resonance position in the ^{19}F and ^{31}P NMR spectra were referenced to external CCl_3F and H_3PO_4 (85%), respectively. The ^{77}Se spectra were obtained using diphenyl diselenide as external standard. Chemical shifts are reported relative to dimethyl selenide (δ 0 ppm) by assuming that the resonance of the standard is at δ 461 ppm [23]. The NMR spectra were processed using the *MestReNova* software [24]. The ESI(+) mass spectra were recorded using an Esquire 3000 ion-trap mass spectrometer (Bruker Daltonic GmbH) equipped with a standard ESI/APCI source. The MALDI(+) mass spectra were recorded from CHCl_3 solutions on a MALDI-TOF Microflex (Bruker) instrument (DCTB as matrix). Mass spectra were processed using the *MASPEC II* software [25]. Infrared spectra (4000–250 cm^{-1}) were recorded with a Perkin-Elmer Spectrum One IR instrument in attenuated total reflection method (ATR).

4.2. Syntheses of compounds

4.2.1. $[\text{Ag}\{2,6-(\text{Me}_2\text{NCH}_2)_2\text{C}_5\text{H}_3\text{N}\}\{\text{O}(\text{O})\text{CCF}_3\}] (2)$

Silver trifluoroacetate, $[\text{AgO}(\text{O})\text{CCF}_3]$ (0.055 g, 0.25 mmol), was added to a solution of **1** (0.048 g, 0.25 mmol) in anhydrous CH_2Cl_2 (20 mL). The reaction mixture was stirred for 1 h at room temperature. The removal of the solvent in vacuum gave **2** as a light yellow solid, which was washed with hexane. Yield: 0.167 g (91%). ^1H NMR (400 MHz, δ , ppm): 2.38 (s, 12H, NMe), 3.57 (s, 4H, CH_2), 7.18 [d, $^3J(\text{HH}) = 7.7$ Hz, 2H, $\text{H}_{3,5}$], 7.74 [t, $^3J(\text{HH}) = 7.7$ Hz, 1H, H_4]. ^{13}C NMR (100.6 MHz, δ , ppm): 46.73 (s, NMe), 64.99 (s, CH_2), 117.53 [q, $^1J(\text{CF}) = 291.3$ Hz, CF_3], 123.46 (s, $\text{C}_{3,5}$), 138.71 (s, C_4), 156.96 (s, $\text{C}_{2,6}$), 162.36 [q, $^2J(\text{CF}) = 34.8$ Hz, $-\text{C}(\text{O})\text{O}-$]. MS [MALDI(+), m/z , %]: 300.1 (100) $[\text{Ag}(\text{N}'\text{NN}')^+]$ $[\text{N}'\text{NN}' = 2,6-(\text{Me}_2\text{NCH}_2)_2\text{C}_5\text{H}_3\text{N}]$. IR (ATR, cm^{-1}): 1679 (s), 1666 (s) $[\nu_{\text{as}}(\text{CO}_2)]$, 1457 (m) $[\nu_{\text{s}}(\text{CO}_2)]$, 423 (w), 404 (m) $[\nu(\text{AgN})]$.

4.2.2. $\{[\text{Ag}(\text{PPh}_3)]_2\{2,6-(\text{Me}_2\text{NCH}_2)_2\text{C}_5\text{H}_3\text{N}\}\}(\text{OTf})_2 (3)$

Silver(triphenylphosphine) triflate $[\text{Ag}(\text{PPh}_3)(\text{OTf})]$ (0.531 g, 1.02 mmol) was added to a solution of **1** (0.099 g, 0.512 mmol) in anhydrous CH_2Cl_2 (20 mL). The reaction mixture was stirred for 1 h at room temperature. After removal of the solvent in vacuum, the solid was washed with hexane to give **3** as a white-cream compound. Yield: 0.51 g (92%). ^1H NMR (400 MHz, δ , ppm): 2.17 (s, 12H, NMe), 3.70 (s, 4H, CH_2), 7.32 [d, $^3J(\text{HH}) = 7.7$ Hz, 2H, $\text{H}_{3,5}$], 7.44 (m, 30H, PPh), 7.81 [t, $^3J(\text{HH}) = 7.7$ Hz, 1H, H_4]. ^{13}C NMR (100.6 MHz, δ , ppm): 46.38 (s, NMe), 65.41 (s, CH_2), 120.57 [q, $^1J(\text{CF}) = 320.8$ Hz, CF_3], 123.61 (s, $\text{C}_{3,5}$), 129.10 [d, $^2J(\text{CP}) = 6.4$ Hz, PPh-ortho], 130.56 [d, $^1J(\text{CP}) = 34.3$ Hz, PPh-*ipso*], 130.88 (s, PPh-*para*), 133.57 [d, $^3J(\text{CP}) = 14.3$ Hz, PPh-*meta*], 139.58 (s, C_4), 157.17 (s, $\text{C}_{2,6}$). ^{19}F NMR (376.5 MHz, δ , ppm): -77.9 (s, 6F, CF_3). ^{31}P NMR (162 MHz, δ , ppm, r.t.): 11.9 (s,br), 16.0 (s,br). ^{31}P NMR (121 MHz, δ , ppm, -55 °C): 11.0 [dd, $^1J(\text{P},^{107}\text{Ag}) = 511.3$, $^1J(\text{P},^{109}\text{Ag}) = 562.8$ Hz], 14.4 [dd, $^1J(\text{P},^{107}\text{Ag}) = 627.6$, $^1J(\text{P},^{109}\text{Ag}) = 724.1$ Hz]. MS [MALDI(+), m/z , %]: 631.3 (100) $[\text{Ag}(\text{PPh}_3)_2^+]$, 562.3 (53) $[\text{Ag}(\text{PPh}_3)(\text{N}'\text{NN}')^+]$ $[\text{N}'\text{NN}' = 2,6-(\text{Me}_2\text{NCH}_2)_2\text{C}_5\text{H}_3\text{N}]$. IR (ATR, cm^{-1}): 1600 (vw), 1583 (vw) $[\nu(\text{C}=\text{C})]$, 1479 (w), 1434 (m) $[\nu(\text{C}=\text{N})]$, 1265 (s), 1242 (m) $[\nu_{\text{as}}(\text{SO}_3)]$, 1221 (m)

$[\nu_{\text{s}}(\text{CF}_3)]$, 1150 (m) $[\nu_{\text{as}}(\text{CF}_3)]$, 1024 (s) $[\nu_{\text{s}}(\text{SO}_3)]$, 432 (w), 400 (w) $[\nu(\text{AgN})]$.

4.2.3. $2,6-(\text{PhSeCH}_2)_2\text{C}_5\text{H}_3\text{N} (4)$

Compound **4** was obtained from Ph_2Se_2 (1 g, 3.2 mmol), NaBH_4 (0.242 g, 6.4 mmol) and 2,6-bis(bromomethyl)pyridine (0.42 g, 3.2 mmol) in ethanol (40 mL) using a slightly modified procedure [8a]. The solvent was completely removed in vacuum from the reaction mixture and the remaining white solid was washed with hexane. Recrystallization from ethanol gave the title compound **4** as white crystals. Yield: 1.176 g (85%). ^1H NMR (300 MHz, δ , ppm): 4.19 [s, $^2J(\text{HSe}) = 12.4$ Hz, 4H, CH_2], 6.92 [d, $^3J(\text{HH}) = 7.7$ Hz, 2H, $\text{H}_{3,5}$], 7.23 (m, 6H, SePh-*meta* + *para*), 7.38 [t, $^3J(\text{HH}) = 7.7$ Hz, 1H, H_4], 7.23 (m, 4H, SePh-*ortho*) = 6.25 Hz, ^{13}C NMR (100.6 MHz, δ , ppm): 33.67 [s, $^1J(\text{CSe}) = 62.5$ Hz, CH_2], 120.98 (s, $\text{C}_{3,5}$), 127.27 (s, SePh-*para*), 128.96 (s, SePh-*meta*), 130.01 (s, SePh-*ipso*), 133.49 (s, SePh-*ortho*), 136.74 (s, C_4), 158.36 (s, $\text{C}_{2,6}$). ^{77}Se NMR (100 MHz, δ , ppm): 368.3 (s). MS [MALDI(+), m/z , %]: 419.9 (20) $[\text{M}^+ + \text{H}]$. IR (ATR, cm^{-1}): 1576 (m), 1568 (m) $[\nu(\text{C}=\text{C})]$, 1475 (m), 1436 (m) $[\nu(\text{C}=\text{N})]$.

4.2.4. $[\text{Au}\{2,6-(\text{PhSeCH}_2)_2\text{C}_5\text{H}_3\text{N}\}\text{Cl}]_2 (5)$

$\text{HAuCl}_4 \cdot 3\text{H}_2\text{O}$ (0.011 g, 0.028 mmol) was added to a solution of **4** (0.012 g, 0.028 mmol) in anhydrous diethyl ether (20 mL). The reaction mixture was stirred for 30 minutes at room temperature, followed by removal of the solvent to dryness. The residue was solved in dry THF (20 mL), treated with NaHCO_3 (0.047 g, 0.056 mmol) and then stirred for 24 h at room temperature. The solution was concentrated to a volume of 3 mL and hexane (5 mL) was added. A yellow oil separated and it was treated with Et_2O and EtOH to give the title compound **5** as a light yellow solid. Yield: 0.020 g (96%). ^1H NMR (400 MHz, δ , ppm): 4.64 [s, 4H, CH_2], 6.92 [d, $^3J(\text{HH}) = 7.7$ Hz, 2H, $\text{H}_{3,5}$], 7.36 [t, $^3J(\text{HH}) = 7.4$ Hz, 4H, SePh-*meta*], 7.44 [m, 3H, H_4 + SePh-*para*], 7.66 [d, $^3J(\text{HH}) = 7.6$ Hz, 4H, SePh-*ortho*]. MS [MALDI(+), m/z , %]: 616.0 (100) $[\text{Au}(\text{SeNSe})^+]$, 419.9 (67) $[\text{SeNSe}^+ + \text{H}]$ $[\text{SeNSe} = 2,6-(\text{PhSeCH}_2)_2\text{C}_5\text{H}_3\text{N}]$. IR (ATR, cm^{-1}): 463 (m) $[\nu(\text{AuN})]$, 330 (m) $[\nu(\text{AuCl})]$.

4.2.5. $[\text{Ag}(\text{PPh}_3)\{2,6-(\text{PhSeCH}_2)_2\text{C}_5\text{H}_3\text{N}\}](\text{OTf}) (6)$

$[\text{Ag}(\text{PPh}_3)(\text{OTf})]$ (0.034 g, 0.06 mmol) was added to a solution of **4** (0.028 g, 0.06 mmol) in anhydrous CH_2Cl_2 (15 mL). The reaction mixture was stirred for 1 h at room temperature. After removal of the solvent in vacuum, the solid was washed with hexane to give **6** as a white solid. Yield: 0.051 g (82%). ^1H NMR (400 MHz, δ , ppm): 4.20 [s, $^2J(\text{HSe}) = 13.8$ Hz, 4H, CH_2], 7.05 [d, $^3J(\text{HH}) = 7.8$ Hz, 2H, $\text{H}_{3,5}$], 7.10 (m, 8H, SePh-*ortho* + *meta*), 7.20 (m, 2H, SePh-*para*), 7.30 (m, 12H, PPh-*ortho* + *meta*), 7.42 (m, 3H, PPh-*para*), 7.52 [t, $^3J(\text{HH}) = 7.7$ Hz, 1H, H_4]. ^{13}C NMR (100.6 MHz, δ , ppm): 35.24 (s, CH_2), 123.28 (s, $\text{C}_{3,5}$), 127.55 (s, SePh-*ipso*), 128.02 (s, SePh-*para*), 129.16 [d, $^2J(\text{CP}) = 9.9$ Hz, PPh-*ortho*], 129.37 (s, SePh-*meta*), 130.79 (s, PPh-*para*), 130.95 [d, $^1J(\text{CP}) = 32.4$ Hz, PPh-*ipso*], 132.68 (s, SePh-*ortho*), 133.63 [d, $^3J(\text{CP}) = 15.8$ Hz, PPh-*meta*], 138.56 (s, C_4), 157.01 (s, $\text{C}_{2,6}$). ^{19}F NMR (376.5 MHz, δ , ppm): -77.9 (s, 3F, CF_3). ^{31}P NMR (121.5 MHz, δ , ppm, -55 °C): 7.9 [dd, $^1J(\text{P},^{107}\text{Ag}) = 398.0$, $^1J(\text{P},^{109}\text{Ag}) = 446.6$ Hz], 11.5 [dd,br,

$^1J(\text{P}, ^{107}\text{Ag}) = 551.6$, $^1J(\text{P}, ^{109}\text{Ag}) = 597.0$ Hz]. MS [ESI(+), m/z , %]: 787.7 [Ag(PPh₃)(SeNSe)⁺], 631.0 [Ag(PPh₃)₂⁺ as base peak] [SeNSe = 2,6-(PhSeCH₂)₂C₅H₃N]. IR (ATR, cm⁻¹): 1592 (w), 1568 (m) [$\nu(\text{C}=\text{C})$], 1478 (m), 1436 (m) [$\nu(\text{C}=\text{N})$], 1268 (m) [$\nu_{\text{as}}(\text{SO}_3)$], 1147 (m) [$\nu_{\text{s}}(\text{CF}_3)$], 1093 (m) [$\nu_{\text{as}}(\text{CF}_3)$], 1022 (m) [$\nu_{\text{s}}(\text{SO}_3)$], 395 (m) [$\nu(\text{AgN})$].

4.2.6. [Cu(MeCN){2,6-(PhSeCH₂)₂C₅H₃N}](PF₆) (7)

To a solution of **4** (0.082 g, 0.2 mmol) in anhydrous CH₂Cl₂ (40 mL) [Cu(MeCN)₄]PF₆ (0.072 g, 0.2 mmol) was added in one portion. The reaction mixture was stirred at room temperature, under argon, for 2 h. Evaporation of the solvent in vacuum resulted in a yellow solid which was washed with hexane to give the title compound **7**. Yield: 0.106 g (87%). ¹H NMR (400 MHz, δ , ppm): 2.22 (s, 3H, Me_{acetone}nitrile), 4.42 (s, br, 4H, CH₂), 7.20 (m, 12H, H_{3,5} + SePh), 7.61 [t, ³J(HH) 7.3 Hz, 1H, H₄]. ¹³C NMR (100.6 MHz, δ , ppm): 1.98 (s, Me_{acetone}nitrile), 38.44 (s, CH₂), 117.12 (s, CN_{acetone}nitrile), 123.46 (s, C_{3,5}), 127.01 (s, SePh-*ipso*), 128.73 (s, SePh-*para*), 129.68 (s, SePh-*meta*), 132.35 (s, SePh-*ortho*), 139.01 (s, C₄), 155.86 (s, C_{2,6}). ¹⁹F NMR (376.5 MHz, δ , ppm): -72.9 [s, ¹J(FP) = 712.4 Hz, 6F, PF₆]. ³¹P NMR (162 MHz, δ , ppm): -144.2 [hept, ¹J(FP) = 712.3 Hz, 1P, PF₆]. MS [ESI(+), m/z , %]: 481.8 [Cu(SeNSe)⁺ as base peak], 419.9 [SeNSe⁺ + H] [SeNSe = 2,6-(PhSeCH₂)₂C₅H₃N]. IR (ATR, cm⁻¹): 1594 (w), 1568 (w) [$\nu(\text{C}=\text{C})$], 1478 (w), 1457 (m) [$\nu(\text{C}=\text{N})$], 828 (vs) [$\nu(\text{PF}_6)$].

4.2.7. [Cu{2,6-(PhSeCH₂)₂C₅H₃N}2](PF₆) (8)

[Cu(MeCN)₄]PF₆ (0.036 g, 0.1 mmol) was added in one portion to a solution of **4** (0.082 g, 0.2 mmol) in anhydrous CH₂Cl₂ (40 mL). The reaction mixture was stirred at room temperature, under argon, for 2 h. The solvent was removed in vacuum. The remaining solid was washed with hexane to give the title compound **8** as a yellow solid. Yield: 0.094 g (92%). ¹H NMR (400 MHz, δ , ppm): 4.37 [s, br, ²J(HSe) = 13.1 Hz, 8H, CH₂], 7.22 (m, 10H, H_{3,5} + SePh-*ortho* + *meta*), 7.31 (m, 2H, SePh-*para*), 7.73 [m, ³J(HH) 7.7 Hz, 1H, H₄]. ¹³C NMR (100.6 MHz, δ , ppm): 36.93 [s, ¹J(CSe) = 61.7 Hz, CH₂], 123.94 (s, C_{3,5}), 128.23 (s, SePh-*ipso*), 128.53 (s, SePh-*para*), 129.61 (s, SePh-*meta*), 132.43 (s, SePh-*ortho*), 139.28 (s, C₄), 156.91 (s, C_{2,6}). ¹⁹F NMR (376.5 MHz, δ , ppm): -73.0 [s, ¹J(FP) = 712.7 Hz, 6F, PF₆]. ³¹P NMR (162 MHz, δ , ppm): -144.2 [hept, ¹J(FP) = 712.7 Hz, 1P, PF₆]. MS [ESI(+), m/z , %]: 900.8 [Cu(SeNSe)₂⁺], 481.8 [Cu(SeNSe)⁺], 419.9 [SeNSe⁺ + H, as base peak] [SeNSe = 2,6-(PhSeCH₂)₂C₅H₃N]. IR (ATR, cm⁻¹): 1593 (w), 1566 (m) [$\nu(\text{C}=\text{C})$], 1477 (m), 1455 (m) [$\nu(\text{C}=\text{N})$], 828 (vs) [$\nu(\text{PF}_6)$].

4.3. X-ray structure determination

The details of the crystal structure determination and refinement for compound **2a** are given in Table 2.

Data were collected at room temperature on Bruker SMART APEX diffractometer, using graphite-monochromated Mo-K α radiation ($\lambda = 0.71073$ Å). For this purpose the crystals were mounted on a cryoloop. The structures were refined with anisotropic thermal parameters. The hydrogen atoms were refined with a riding model and a

Table 2

X-ray crystal data and structure refinement for compound **2a**.

Empirical formula	C ₂₁ H ₂₄ Ag ₄ F ₁₂ N ₃ O ₉
Formula weight	1121.91
T (K)	297(2)
Crystal system	Triclinic
Space group	P-1
a (Å)	10.83(2)
b (Å)	13.90(3)
c (Å)	13.90(3)
α (°)	79.27(2)
β (°)	70.73(2)
γ (°)	70.73(2)
Volume (Å ³)	1859(6)
Z	2
ρ_{calc} (g cm ⁻³)	2.004
Absorption coefficient (mm ⁻¹)	2.180
F(000)	1078
Crystal size (mm ³)	0.31 × 0.26 × 0.21
Reflections collected	13132
Independent reflections	6486 [$R_{\text{int}} = 0.0343$]
R_1 ($I > 2\sigma(I)$)	0.0616
wR ₂	0.1256
Goodness-of-fit on F ²	1.092

mutual isotropic thermal parameter, except the one from the ethanol group which was not determined. The O(3) and O(4) atoms from a OOCF₃ fragment and the four CF₃ groups are disordered over two positions and were refined with the following occupancies: 38:62, 37:63, 40:60, 60:40 and 39:61, respectively. For structure solving and refinement the software package SHELX-97 was used [26]. The drawings were created with the Diamond program [27].

Crystallographic data for the structural analysis of compound **2a** have been deposited with the Cambridge Crystallographic Data Centre CCDC No. 865938. These data can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB2 1EZ, UK; tel.; +44 1223 336 408; fax: +44 1223 336 033; deposit@ccdc.cam.ac.uk).

Acknowledgements

Financial support from the National University Research Council (CNCSIS, Romania; Research Project TD66/2007) and Ministerio de Economía y Competitividad (Spain, MEC/FEDER CTQ2010-20500-C02-01) is gratefully acknowledged. We also thank the National Center for X-ray Diffraction (Babes-Bolyai University, Cluj-Napoca, Romania) for support in the solid state structure determination.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.crci.2012.05.012>.

References

- [1] (a) R.A.T.M. Abbenhuis, J. Boersma, G. van Koten, J. Org. Chem. 63 (1998) 4282; (b) I. del Rio, G. van Koten, Tetrahedron Lett. 40 (1999) 1401;

- (c) I. del Rio, R.A. Gossage, M.S. Hannu, M. Lutz, A.L. Spek, G. van Koten, *Organometallics* 18 (1999) 1097;
- (d) R.I. del Rio, G. van Koten, M. Lutz, A.L. Spek, *Organometallics* 19 (2000) 361;
- (e) B. Göbelt, K. Matyjaszewski, *Macromol. Chem. Phys.* 201 (2000) 1619;
- (f) R.K. O'Reilly, V.C. Gibson, A.J.P. White, D.J. Williams, *Polyhedron* 23 (2004) 2921;
- (g) G.J.P. Britovsek, J. England, S.K. Spitzmesser, A.J.P. White, D.J. Williams, *Dalton Trans.* (2005) 945.
- [2] (a) R.A.T.M. Abbenhuis, I. del Rio, M.M. Bergshoef, J. Boersma, N. Veldman, A.L. Spek, G. van Koten, *Inorg. Chem.* 37 (1998) 1749;
- (b) I. del Rio, R.A. Gossage, M. Lutz, A.L. Spek, G. van Koten, *Inorg. Chim. Acta* 287 (1999) 113;
- (c) I. del Rio, R.A. Gossage, M. Lutz, A.L. Spek, G. van Koten, *J. Organomet. Chem.* 583 (1999) 69;
- (d) I. del Rio, S. Back, M.S. Hannu, G. Rheinwald, H. Lang, G. van Koten, *Inorg. Chim. Acta* 300 (2000) 1094;
- (e) S. Back, G. Rheinwald, I. del Rio, G. van Koten, H. Lang, *Acta Crystallogr. Sect. E* 57 (2001) m444;
- (f) R. Packheiser, P. Ecorchard, B. Walfort, H. Lang, *J. Organomet. Chem.* 693 (2008) 933.
- [3] T.W. Welch, S.A. Cifan, P.S. White, H.H. Thorp, *Inorg. Chem.* 36 (1997) 4812.
- [4] (a) B.A. Markies, P. Wijkens, J. Boersma, A.L. Spek, G. van Koten, *Rec. Trav. Chim. Pays-Bas* 110 (1991) 133;
- (b) B.A. Markies, P. Wijkens, J. Boersma, H. Kooijman, N. Veldman, A.L. Spek, G. van Koten, *Organometallics* 13 (1994) 3244;
- (c) B.A. Markies, P. Wijkens, A. Dedieu, J. Boersma, A.L. Spek, G. van Koten, *Organometallics* 14 (1995) 5628.
- [5] (a) V.W. Day, M.F. Fredrich, S. Dabestani, P.S. Bryan, *ACA Ser.* 25 (1977) 23;
- (b) A.N. Vedernikov, P. Wu, J.C. Huffman, K.G. Caulton, *Inorg. Chim. Acta* 330 (2002) 103.
- [6] B. Masci, P. Thuery, *Acta Crystallogr. Sect. C* 60 (2004) m584.
- [7] (a) P. Molenveld, W.M.G. Stikvoort, H. Kooijman, A.L. Spek, J.F.J. Engbersen, D.N. Reinhoudt, *J. Org. Chem.* 64 (1999) 3896;
- (b) I. del Rio, R.A. Gossage, M.S. Hannu, M. Lutz, A.L. Spek, G. van Koten, *Can. J. Chem.* 78 (2000) 1620.
- [8] (a) D. Das, G.K. Rao, A.K. Singh, *Organometallics* 28 (2009) 6054;
- (b) D. Das, P. Singh, O. Prakash, A.K. Singh, *Inorg. Chem. Commun.* 13 (2010) 1370.
- [9] (a) E.J. Fernández, A. Laguna, J.M. López-de-Luzuriaga, *Coord. Chem. Rev.* 249 (2005) 1423;
- (b) E.J. Fernández, A. Laguna, J.M. López-de-Luzuriaga, *Dalton Trans.* (2007) 1969.
- [10] J. Heine, H. Westemeier, S. Dehnen, Z. Anorg. Allg. Chem. 636 (2010) 996.
- [11] E.C. Constable, G. Zhang, C.E. Housecroft, M. Neuburger, J.A. Zampese, *Eur. J. Inorg. Chem.* (2010) 2000.
- [12] J.D. Crowley, P.H. Bandoen, L.R. Hanton, *Polyhedron* 29 (2010) 70.
- [13] (a) S. Gonzalez, L. Valencia, R. Bastida, D.E. Fenton, A. Macias, A. Rodriguez, *J. Chem. Soc., Dalton Trans.* (2002) 3551;
- (b) H. Adams, B.A. Najera, *J. Coord. Chem.* 56 (2003) 1215.
- [14] R. Cacciapaglia, A. Casnati, L. Mandolini, D.N. Reinhoudt, R. Salvio, A. Sartori, R. Ungaro, *J. Org. Chem.* 70 (2005) 624.
- [15] (a) H. Takemura, N. Kon, M. Yasutake, T. Shinmyozu, *Tetrahedron* 59 (2003) 427;
- (b) M.C. Fernandez-Fernandez, R. Bastida, A. Macias, P. Perez-Lourido, L. Valencia, *Inorg. Chem.* 45 (2006) 2266;
- (c) Y. Ma, B. Liu, C. Xue, *Acta Crystallogr. E* 66 (2010) m1117;
- (d) E.C. Constable, G. Zhang, C.E. Housecroft, J.A. Zampese, *Cryst. Eng. Commun.* 12 (2010) 3724.
- [16] G.B. Deacon, R.J. Phillips, *Coord. Chem. Rev.* 33 (1980) 227.
- [17] S.J. Angus-Dunne, L.E.O. Lee Chin, R.C. Burns, G.A. Lawrance, *Trans. Met. Chem.* 31 (2006) 268.
- [18] (a) E.L. Muetterties, C.W. Aleganti, *J. Am. Chem. Soc.* 94 (1972) 6386;
- (b) P.F. Barron, J.C. Dyason, P.C. Healy, L.M. Engelhardt, B.W. Skelton, A.H. White, *J. Chem. Soc. Dalton Trans.* (1986) 1965;
- (c) A. Caballero, A. Guerrero, F.A. Jalón, B.R. Manzano, R.M. Claramunt, M.D. Santa Maria, C. Escolástico, J. Elguero, *Inorg. Chim. Acta* 347 (2003) 168.
- [19] (a) S.M. Socol, J.G. Verkade, *Inorg. Chem.* 23 (1984) 3487;
- (b) R.E. Bachman, D.F. Andretta, *Inorg. Chem.* 37 (1998) 5657;
- (c) A. Cingolani, Efendi, F. Marchetti, C. Pettinari, R. Pettinari, B.W. Skelton, A.H. White, *Inorg. Chim. Acta* 329 (2002) 100;
- (d) E.J. Sekabunga, M.L. Smith, T.R. Webb, W.E. Hill, *Inorg. Chem.* 41 (2002) 1205;
- (e) A. Ilie, C.I. Rat, S. Scheutzow, K. Lux, T.M. Klapötke, C. Silvestru, K. Karaghiosoff, *Inorg. Chem.* 50 (2011) 2675.
- [20] M. Barrow, H.-B. Bürgi, M. Camalli, F. Caruso, E. Fischer, L.M. Venanzi, L. Zambonelli, *Inorg. Chem.* 22 (1983) 2356.
- [21] J. Emsley, *Die Elemente*, Walter de Gruyter, Berlin, 1994.
- [22] (a) R.G. Griffin, J.D. Ellett Jr., M. Mehring, J.G. Bullitt, J.S. Waugh, *J. Chem. Phys.* 57 (1972) 2147;
- (b) E.V. Karpova, A.I. Boltalin, Yu.M. Korenev, S.I. Troyanov, *Koord. Khim.* 25 (1999) 70.
- [23] P.N. Jayaram, G. Roy, G. Mugesh, *J. Chem. Sci.* 120 (2008) 143.
- [24] MestReC and MestReNova, Mestrelab Research S.L., A Coruña 15706, Santiago de Compostela.
- [25] MASPEC II Data System, Mass Spectrometry Services Ltd., M25 9WB Manchester.
- [26] G.M. Sheldrick, *Acta Crystallogr. Sect. A* 64 (2008) 112.
- [27] K. Brandenburg, DIAMOND–Visual Crystal Structure Information System, Release 3.1d, Crystal Impact GbR, Bonn, Germany, 2006.