



Full paper/Mémoire

# A facile and efficient [bmim]N<sub>3</sub> catalyzed direct oxidative esterification of arylaldehydes with alcohols

Hassan Valizadeh <sup>\*</sup>, Mina Ahmadi

Department of Chemistry, Faculty of Sciences, Azarbaijan Shahid Madani University, 53714-161 Tabriz, Iran

## ARTICLE INFO

## Article history:

Received 29 April 2012

Accepted after revision 10 August 2012

Available online 18 September 2012

## Keywords:

Oxidative esterification

Direct esterification of aldehydes

[bmim]N<sub>3</sub>

Task-specific ionic liquid

## ABSTRACT

Task-specific ionic liquid, [bmim]N<sub>3</sub> was used as an effective catalyst and reaction medium for the direct oxidative esterification of arylaldehydes with alcohols. The oxidative esterification reaction of a variety of arylaldehydes took place smoothly with some primary and secondary alcohols in [bmim]N<sub>3</sub>. Satisfactory results were obtained with arylaldehydes containing electron withdrawing groups. Tertiary alcohols did not react under these conditions.

© 2012 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

## 1. Introduction

The stoichiometric activation of the parent acid as an acyl halide, anhydride or activated ester (in situ or otherwise) amenable to subsequent nucleophilic substitution is the most common strategy for the synthesis of carboxylic acid ester derivatives under mild conditions [1,2]. An interesting and potentially valuable alternative transformation, in which there has been a recent resurgence in interest, is the catalytic oxidative esterification of aldehydes under mild conditions [2–6]. Few oxidative esterification reactions have been reported until now [4,7–9].

Ionic liquids are produced from the weak coordination of organic cations such as 1-butyl-3-methylimidazolium, *N*-alkylpyridinium or tetraalkylammonium with inorganic anions such as Cl<sup>−</sup>, BF<sub>4</sub><sup>−</sup> or HSO<sub>4</sub><sup>−</sup>. These compounds have attracted the chemist's interest due to their interesting physical and chemical properties [10,11]. Many of these compounds are able to play multiple roles as catalyst and/or reagent and solvent have been reported in the literature.


These types of ionic liquids were known as task-specific ionic liquids (TSILs). TSILs contain functional groups, which are covalently bonded to a cation or anion in these compounds, and have been increasingly used in synthetic organic chemistry [12–16]. In continuation of our interest to use ionic liquids (IL's), water or solventless systems as green reaction media [17–21], in this report, we wish to highlight our results on using [bmim]N<sub>3</sub> as an efficient catalyst in the direct oxidative esterification of arylaldehydes with alcohols (Scheme 1). This new oxidative esterification method is expected to be an attractive procedure directly giving esters from aldehydes and alcohols from the viewpoint of green chemistry.

## 2. Results and discussions

Ionic liquid [bmim]N<sub>3</sub> was readily prepared from [bmim]Cl via the exchange of Cl<sup>−</sup> by N<sub>3</sub><sup>−</sup> in the presence of NaN<sub>3</sub> at room temperature in 91% yield and purified by drying in a vacuum at 60 °C to remove the residual starting materials and reagents. Esterification of *p*-nitrobenzaldehyde with methanol was chosen as a model reaction. The activity of ionic liquid [bmim]Cl as catalyst was examined for the model reaction in the absence of any solvent and oxidant at room temperature. The experimental results

\* Corresponding author.

E-mail address: h-valizadeh@azaruniv.edu (H. Valizadeh).



Scheme 1. [Bmim] $N_3$  catalyzed solvent-free direct oxidative esterification of arylaldehydes.

revealed that a negligible amount of methyl *p*-nitrobenzoate was obtained over 10 h at room temperature and also at higher temperatures. Several ionic liquids such as [bmim]Cl, [bmim]Br, [bmim]BF<sub>4</sub> and [bmim]N<sub>3</sub> were examined in this procedure. The experimental results using [bmim]Br are the same as using [bmim]Cl under different conditions. Examining the typical reaction in [bmim]BF<sub>4</sub> at different temperatures showed that this ionic liquid is also not the effective catalyst for the oxidative esterification reaction and no ester product was identified under these conditions. The best results were found using 1-butyl-3-methylimidazolium azide in this procedure and methyl *p*-nitrobenzoate was isolated in 55% yield at room temperature. The reaction conditions was optimized using different molar ratios of the [bmim]N<sub>3</sub>, *p*-nitrobenzaldehyde and MeOH at different temperatures. The best ratio was found to be 3:1:3 at 58 °C. [Bmim]N<sub>3</sub> was examined for the oxidative esterification of other arylaldehydes. As can be seen from Table 1, aromatic aldehydes, containing electron withdrawing groups react faster and afford the related esters in higher yields in comparison with those containing electron donating groups such as *o*-hydroxybenzaldehyde, *o*-methoxybenzaldehyde, *p*-methoxybenzaldehyde and  $\beta$ -resorcinolaldehyde under the same conditions. The reaction of *o*-nitrobenzaldehyde and 3-nitrobenzaldehyde with Me-OH, Et-OH or *pr*-OH to yield the related

esters is more sluggish, as compared to the corresponding *p*-substituted substrates. Aromatic aldehydes substituted with electron-donating groups at the *p*-position such as *p*-hydroxybenzaldehyde and *p*-methoxybenzaldehyde react slowly. At most 10% conversions to methyl *p*-hydroxybenzoate and methyl *p*-methoxybenzoate were observed in 10 h under the reaction conditions. We also tested the reaction of aryl aldehydes with electron-donating groups while bubbling air and oxygen through the reaction mixture and no significant effect on the oxidation was observed.

We used the different primary alcohols such as Et-OH, *Pr*-OH and Bu-OH in reaction with some arylaldehydes under optimized conditions. The results were gathered in Table 1.

As it can be seen from Table 1, good to high yields of the related esters were prepared from the reaction of mentioned alcohols with different arylaldehydes. *p*-Nitrobenzaldehyde was reacted with isopropylalcohol in the presence of [bmim]N<sub>3</sub> and afforded to isopropyl-*p*-nitrobenzoate in 90% yield at 60 °C. Testing the reaction of *t*-butylalcohol with arylaldehydes containing different electron releasing and electron withdrawing groups in [bmim]N<sub>3</sub> showed that no any ester products were formed under different conditions. All of products were characterized by comparing their (<sup>1</sup>H-NMR and IR) spectroscopic data and melting points with literature values. For comparison, the reaction times and yields of some products of the present method and other reported methods were gathered in Table 2.

In summary, a variety of arylaldehydes were directly transformed to the related alkylesters using task-specific azide ionic liquid, [bmim]N<sub>3</sub> via the oxidative esterification under mild conditions. It was found that [bmim]N<sub>3</sub> is a highly effective, mild and convenient catalyst for the direct oxidative esterification of arylaldehydes with alcohols. Azide ionic liquid plays dual role as catalyst and reaction

Table 1  
[Bmim]N<sub>3</sub> catalyzed solvent-free direct oxidative esterification of arylaldehydes with alcohols.

Entry	R <sup>1</sup>	R <sup>2</sup>	Product number	Yield <sup>a</sup> (%)	M.P. (lit [Ref.]) (°C)	
					Found	Reported [Ref.]
1	4-CN	Me	<b>2a</b>	81	61–62	62 [22]
2	4-CN	Et	<b>2b</b>	85	50–51	52–54 [23]
3	4-CN	Pr	<b>2c</b>	70	35–36	–
4	4-CN	Bu	<b>2d</b>	76	51–52	54–55 [24]
5	<i>p</i> -NO <sub>2</sub>	Me	<b>2e</b>	80	90–91	95 [25]
6	<i>p</i> -NO <sub>2</sub>	Et	<b>2f</b>	82	54–55	56–57 [25]
7	<i>p</i> -NO <sub>2</sub>	Pr	<b>2g</b>	85	30–31	33–34 [25]
8	<i>p</i> -NO <sub>2</sub>	<i>i</i> -pr	<b>2h</b>	90	103–105	–
9	<i>p</i> -NO <sub>2</sub>	Bu	<b>2i</b>	75	30–31.5	35–37 [25]
10	3-NO <sub>2</sub>	Me	<b>2j<sup>b</sup></b>	50	75–76	78 [26]
11	2-OH	Me	<b>2k</b>	–	–	–
12	2OMe	Me	<b>2l</b>	–	–	–
13	2,4-Dihydroxy	Me	<b>2m</b>	–	–	–
14	4-OMe	Me	<b>2n</b>	–	–	–

<sup>a</sup> Isolated yields were determined based on the starting aldehyde.

<sup>b</sup> The yields of 3-nitrobenzaldehyde in reaction with other alcohols were very low.

Table 2  
Comparison of the times and yields of some products of the present method with reported methods.

Product number	Time (h)		Yield (%)	
	Found	Reported [Ref.]	Found	Reported [Ref.]
<b>2d</b>	3	16	76	42 [22]
<b>2e</b>	3	12	80	78 [8]
<b>2a</b>	3	–	81	79 [3]
<b>2f</b>	3	5	82	75 [27]

medium in this procedure. Although the literature enumerates a number of procedures for conversion of aldehydes to esters, the simplicity, solvent-free conditions, and inexpensiveness of our procedure makes it a practical alternative.

### 3. Selected spectroscopic-data

**4-Cyanobenzoic acid methyl ester (2a):** white solid; IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 2229 (CN), 1718 (C=O), 1600 (C=C);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.16–8.13 (d, 2H,  $J = 7.2$  Hz), 7.76–7.74 (d, 2H,  $J = 7.2$  Hz), 3.96 (s, 3H).

**4-Cyanobenzoic acid ethyl ester (2b):** white solid; IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 2230 (CN), 1719 (C=O), 1602 (C=C);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.16–8.13 (d, 2H,  $J = 7.1$  Hz), 7.76–7.74 (d, 2H,  $J = 7.1$  Hz), 4.46–4.39 (q, 2H), 1.44–1.40 (t, 3H).

**4-Cyanobenzoic acid n-propyl ester (2c):** yellow solid; IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 2227 (CN), 1725 (C=O), 1609 (C=C);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.17–8.14 (d, 2H), 7.77–7.74 (d, 2H), 4.35–4.3 (t, 2H,  $J = 6.8$  Hz), 1.85–1.78 (m, 2H), 1.07–1.02 (t, 3H,  $J = 7.2$  Hz).

**4-Cyanobenzoic acid n-butyl ester (2d):** light yellow solid, IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 2232 (CN), 1719 (C=O), 1611 (C=C);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.16–8.13 (d, 2H,  $J = 7.3$  Hz), 7.77–7.74 (d, 2H,  $J = 7.3$  Hz), 4.39–4.35 (t, 2H,  $J = 6.8$  Hz), 1.80–1.73 (m, 2H), 1.58–1.45 (m, 2H), 1.02–0.98 (t, 3H,  $J = 7.6$  Hz).

**4-Nitro-benzoic acid methyl ester (2e):** light yellow solid; IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 1718 (C=O), 1339 and 1519 ( $\text{NO}_2$ ), 1608 (C=C);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.33–8.31 (d, 2H,  $J = 7.2$  Hz), 8.23–8.20 (d, 2H,  $J = 7.2$  Hz), 3.98 (s, 3H).

**4-Nitro-benzoic acid ethyl ester (2f):** light yellow solid; IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 1719 (C=O), 1339 and 1519 ( $\text{NO}_2$ ), 1610 (C=C);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.31–8.28 (d, 2H,  $J = 7.1$  Hz), 8.22–8.21 (d, 2H,  $J = 7.1$  Hz), 4.46–4.41 (q, 2H), 1.46–1.41 (t, 3H,  $J = 7.2$  Hz).

**4-Nitro-benzoic acid n-propyl ester (2g):** yellow solid; IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 1719 (C=O), 1341 and 1520 ( $\text{NO}_2$ ), 1600 (C=C);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.31–8.28 (d, 2H,  $J = 7.2$  Hz), 8.23–8.20 (d, 2H,  $J = 7.2$  Hz), 4.36–4.32 (t, 2H,  $J = 6.8$  Hz), 1.86–1.79 (m, 2H), 1.07–1.02 (t, 3H,  $J = 7.4$  Hz).

**4-Nitro-benzoic acid isopropyl ester (2h):** solid; IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 1719 (C=O), 1345 and 1525 ( $\text{NO}_2$ ), 1604 (C=C);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.30–8.27 (d, 2H,  $J = 7.4$  Hz), 8.22–8.19 (d, 2H,  $J = 7.4$  Hz), 4.36–4.32 (t, 2H,  $J = 6.8$  Hz), 1.86–1.79 (m, 2H), 1.55 (m, 1H), 1.42–1.140 (d, 6H,  $J = 7.2$  Hz).

**4-Nitro-benzoic acid n-butyl ester (2i):** light yellow solid; IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 1719 (C=O), 1343 and 1529 ( $\text{NO}_2$ ), 1601 (C=C);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.30–8.27 (d, 2H,  $J = 7.3$  Hz), 8.22–8.19 (d, 2H,  $J = 7.2$  Hz), 4.41–4.36 (t, 2H,  $J = 7.6$  Hz), 1.81–1.74 (m, 2H), 1.53–1.45 (m, 2H), 1.02–0.97 (t, 3H,  $J = 7.8$  Hz).

**3-Nitro-benzoic acid methyl ester (2j):** light yellow solid; IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 1718 (C=O), 1340 and 1524 ( $\text{NO}_2$ ), 1608 (C=C);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.28–8.24 (d, 2H,  $J = 7.1$  Hz), 8.22–8.18 (d, 2H,  $J = 7.1$  Hz), 4.00 (s, 3H).

## 4. Experimental

### 4.1. General information

All reagents were purchased from Merck Company and used without further purification. Infrared spectra were recorded in KBr and were determined on a Perkin Elmer FT-IR spectrometer.  $^1\text{H}$  NMR spectra were recorded on a Bruker Avance AC-300 MHz using TMS as internal standard. All melting points measured in open glass-capillaries using Stuart melting point apparatus.

### 4.2. Preparation of 1-butyl-3-methylimidazolium azide

1-Butyl-3-methylimidazolium chloride was prepared from the reaction of *N*-methylimidazole with *n*-butylchloride at 80 °C under neat conditions. Sodium azide (46 mmol) was added to a solution of this freshly prepared ionic liquid (40 mmol) in dichloromethane (18 mL) and stirred for 20 h at room temperature. The suspension was filtered to remove the precipitated sodium chloride salt and the organic layer was washed with water (5 × 8 mL) until no precipitation of AgCl occurred in aqueous phase on addition of a concentrated  $\text{AgNO}_3$  solution. The solvent and other volatile materials were removed from organic layer in vacuo and the resulting ionic liquid was stirred with activated charcoal for 12 h, after which the ionic liquid was passed through a short alumina column(s) (acidic and/or neutral) to give a colorless ionic liquid, which was dried at 60 °C in vacuo for 24 h or until no visible signs of water present in the IR spectrum. Yields generally 80–85%. IR (KBr): 2120, 1649, 1613, 1529, 1353, 1200  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.15 (3H, t,  $-\text{CH}_3$ ), 1.48 (2H, m,  $\text{CH}_2$ ), 1.80 (2H, m,  $\text{CH}_2$ ), 3.85 (3H, s,  $\text{N}-\text{CH}_3$ ), 4.01 (2H, t,  $-\text{N}-\text{CH}_2$ ), 7.24 (1H, d, Ar-H), 7.45 (1H, d, Ar-H), 9.21 (1H, broad, Ar-H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ): 12.30, 19.1, 24.4, 36.4, 38.7, 124.1, 127.0, 142.8. Anal. Calcd (%) for  $\text{C}_8\text{H}_{15}\text{N}_5$ : C, 53.02; H, 8.34; N, 38.64. Found (%): C, 53.12; H, 8.36; N, 38.61.

### 4.3. Oxidative esterification of arylaldehydes using [bmim]N<sub>3</sub>, general procedure

Arylaldehyde (25 mmol) and [bmim]N<sub>3</sub> (75 mmol) and alcohol (75 mmol) were mixed thoroughly and stirred at 50–60 °C. The completion of reaction was monitored by TLC using (EtOAc/petroleum ether 1:7) as eluent and the product was isolated via preparative chromatography from EtOAc/petroleum ether as eluent.

## Acknowledgement

The partial financial assistance from the Research Vice Chancellor of Azarbaijan Shahid Madani University is gratefully acknowledged.

## References

- [1] J. Otera, Esterification: Methods, Reactions and Applications, Wiley, New York, 2003.
- [2] R.C. Larock, Comprehensive Organic Transformations, VCH, New York, 1989.
- [3] R. Lerebours, C. Wolf, J. Am. Chem. Soc. 128 (2006) 13052.

- [4] W.-J. Yoo, C.-J. Li, *J. Org. Chem.* 71 (2006) 6266.
- [5] N. Mori, H. Togo, *Tetrahedron* 61 (2005) 5915.
- [6] R. Gopinath, B.K. Patel, *Org. Lett.* 2 (2000) 577.
- [7] C. Noonan, L. Baragwanath, S.J. Connon, *Tetrahedron Lett.* 49 (2008) 4003.
- [8] S.I. Kiyooka, Y. Wada, M. Ueno, T. Yokoyama, R. Yokoyama, *Tetrahedron* 63 (2007) 12695.
- [9] Y. Diaoa, R. Yana, S. Zhanga, P. Yanga, Z. Li, L. Wang, H. Dong, *J. Mol. Catal. A: Chem* 303 (2009) 35.
- [10] K.N. Marsh, J.A. Boxall, R. Lichtenthaler, *Fluid Phase Equilib.* 219 (2004) 93.
- [11] S. Keskin, D. Kayrak-Talay, U. Akman, Ö. Hortacsu, *J. Supercrit. Fluids* 43 (2007) 150.
- [12] H. Zhao, N. Yu, Y. Ding, R. Tan, C. Liu, D. Yin, H. Qiu, D. Yin, *Micropor. Mesopor. Mat.* 136 (2010) 10.
- [13] L. Wang, H.P. Li, *Tetrahedron* 65 (2009) 364.
- [14] W. Bi, M. Tian, J. Zhou, K.H. Row, *J. Chromatogr. B* 878 (2010) 2243.
- [15] L.D.S. Yadav, R. Patel, V.K. Rai, V.P. Srivastava, *Tetrahedron Lett.* 48 (2007) 7793.
- [16] J. Sun, W. Cheng, W. Fan, Y. Wang, Zh. Meng, S. Zhang, *Catal. Today* 148 (2009) 361.
- [17] H. Valizadeh, A. Fakhari, *Molecules* 15 (2010) 2972.
- [18] H. Valizadeh, A. Shomali, *Dyes Pigments* 92 (2012) 1138.
- [19] H. Valizadeh, M. Amiri, A. Shomali, *C. R. Chimie* 14 (2011) 1103.
- [20] H. Valizadeh, H. Gholipour, *C. R. Chimie* 14 (2011) 963.
- [21] H. Valizadeh, M. Amiri, E. Khalili, *Mol. Divers* 16 (2012) 319.
- [22] M. Haake, B. Schümmelfeder, *Synthesis* 9 (1991) 753.
- [23] <http://www.lookchem.com/>.
- [24] A. Schoenberg, I. Bartoletti, R.F. Heck, *J. Org. Chem.* 39 (1974) 3318.
- [25] R. Tank, U. Pathak, M. Vimal, Sh. Bhattacharyya, L. Kumar Pandey, *Green Chem.* 13 (2011) 3350.
- [26] L. Williamson, Kenneth "Macroscale and Microscale Organic Experiments" (Textbook), fourth ed., Houghton Mifflin, 2003.
- [27] G.A. Heropoulos, C. Villalonga-Barber, *Tetrahedron Lett.* 52 (2011) 5319.