

# Lead abundances in bony fish from the world oceans: significance for contaminated marine habitats and environmental policies

Alain Véron<sup>a,b</sup>, Isabelle Bernier<sup>c</sup> and Bruno Hamelin<sup>a</sup>

<sup>a</sup> CEREGE, Aix-Marseille Univ., CNRS, IRD, Coll. De France, Europôle Méditerranéen de l'Arbois, BP80, 13545 Aix en Provence cedex 4, France.

<sup>b</sup> GEOTOP, Université du Québec à Montréal, CP8888 Centre-Ville, Montréal, QC H3C3P8, Canada.

<sup>c</sup> Rectorat Aix-Marseille, DRAES, Place Lucien Paye, 13621 Aix en Provence cedex 1, France.

Sup. Table S1: Lead concentrations in muscles and livers from the main oceanic basins with corresponding references and FAO sampling areas (detailed captions are provided in the excel file).

Le	mean fish length (cm) range is provided when mean is not available "mixed" is used for juvenile-adult composites or several species "juv" is used for juveniles (when size not available)
sd	standard deviation
M	Pb concentration in muscle (ppm wet weight)
L	Pb concentration in muscle (ppm wet weight)
<b>0.000</b>	wet concentration converted from dry concentration
sd	standard deviation
n	number of analyzed fish tissues
DATE	date of sampling
Fa	FAO area
Fsa	FAO subarea
Fd	FAO division
GFCMsa	GFCM subarea
LOC	nearest regional sea and/or country
na	not available



## REF

- 1 Afandi, I., Talba, S., et al. (2018). Trace metal distribution in pelagic fish species from the north-west African coast (Morocco). *Int. Aquat. Res.*, 10:191-205.
- 2 Alfonso, C., Lourenco, H. M., Dias, A., Nunes, M. L., and Castro, M. (2007). Contaminant metals in black seabream fish (*Aphanius carbo*) caught off Madeira and the Azores. *Food Chem.*, 101, 120-125.
- 3 Bosch, A. C. (2015). Status of mercury and other heavy metals in South African marine fish species. Thesis, Faculty of AgriSciences, Stellenbosch Univ., South Africa. 149p
- 4 Brown, F. M. J., and Balls, P. W. (1997). In: Trace metals in fish and shellfish from Scottish waters, Scottish Fisheries Research Report number 60, The Scottish Office Agriculture, Environment and Fisheries Dept. 36p.
- 5 Burger, J., and Goedkoop, P. W. (2005). Heavy metals in commercial fish in New Jersey. *Environ. Res.*, 99:403-412.
- 6 Celik, U., Cakli, S., and Oschlenschlaeger, J. (2004). Determination of the lead and cadmium burden in some eastern Atlantic and Mediterranean fish species by DPSAV. *Eur. Food Res. Technol.*, 218:298-305.
- 7 Cifuentes, A., and Diaz, J. M. (2004). Trace elements in fish from the eastern sea of Mexico: identification of sources of contamination. *Food Chem.*, 147:357-360.
- 8 Coops, D., Auger, D., Averyt, B., Lutze, M., Masse, P., and Hall, J. (1992). Flounder (*Platichthys flesus*) muscles as an indicator of metal and organochlorine contamination of French Atlantic coastal waters. *Ambio*, 21(2):176-182.
- 9 Cronin, M., Davies, I. M., Newton, A., Pirie, J. M., Topping, G., and Swan, S. (1998). Trace metal concentrations in deep-sea fish from the North Atlantic. *Mar. Environ. Res.*, 45(3):225-238.
- 10 Diop, M., Howsam, M., Diop, C., Cazier, F., Goossens, J. F., Diouf, A., and Amara, R. (2016). Spatial and seasonal variations of trace elements concentrations in liver and muscle of round Sardineille (*Sardinella aurita*) and Senegalese sole (*Solea senegalensis*) along the Senegalese coast. *Chemosphere*, 144:758-766.
- 11 Duquenne, S. (1992). Bioaccumulation métallique et métallophénolique chez trois espèces de poissons du littoral Nord-Pas-de-Calais. Thesis, Univ. Lille, France. 264p.
- 12 Guerin, T., Chebel, R., Vastel, C., Siroz, V., Volatier, J.-L., Leblanc, J.-C., and Noël, L. (2011). Determination of 20 trace elements in fish and other seafood from the French market. *Food Chem.*, 127:934-942.
- 13 Hanson, P. J. (1997). Response of hepatic trace element concentrations in fish exposed to elemental and organic contaminants. *Estuaries*, 20(4), 659-676.
- 14 Henry, F., Amara, R., Courcot, L., Lacouture, D., and Bertho, M.-L. (2004). Heavy metals in four fish species from the French coast of the Eastern English Channel and Southern Bight of the North Sea. *Environ. Int.*, 30: 675-683.
- 15 Julshamn K., Lundbyke, A. K., Hegstad, K., Berntsen, M., and M.G.M. Boe, (2015). Norwegian monitoring programme on the inorganic and organic contaminants in fish caught in the Barents Sea, Norwegian Sea and North Sea, 1994-2001. *Food Add. & Cont.*, 21(4):365-376.
- 16 Lozañez, G., Brito, A., Hardisson, A., Gutierrez, A., Gonzalez-Weller, D., Lozañez, I.J. (2009). Content of lead and cadmium in barrel hogfish, Bodianus scrofa, island grouper, Mycteroperca rubra, and Portuguese dog fish, Centroscymnus coelolepis, from Canary Islands, Spain. *Bull. Environ. Contam. Toxicol.*, 83, 591-594.
- 17 Olmedo, P., Pla, A., Hernandez, A. F., Barber, F., Ayuñil, L., and Gil, F. (2013). Determination of toxic elements in fish and sediments from the Barents Sea. Draft report. Bergen, Norway: Norwegian Institute for Marine Research.
- 18 Olmedo-Pla, A., Hernandez, A. F., Barber, F., Ayuñil, L., and Gil, F. (2013). Determination of toxic elements in fish and sediments from the Barents Sea. Draft report. Bergen, Norway: Norwegian Institute for Marine Research.
- 19 Massé, A., Stange, K., and Klungsoy, J. (1996). In: Trace elements in fish and sediments from the Barents Sea. Draft report. Bergen, Norway: Norwegian Institute for Marine Research.
- 20 Medeiros, R. J., dos Santos, L. M. G., Freire, A. S., Santelli, R. E., Braga, A. N. C. B., Krauss, T. M., and Jacob, S. D. C. (2012). Determination of inorganic trace elements in edible marine fish from Rio de Janeiro State, Brazil. *Food Control*, 23:535-541.
- 21 Morzano, M. A., Rabaneto, L. C., Milani, R. F., Miyazakura, L., and Balzan, S. C. (2011). Assessment of trace elements in fishes of Japanese foods marketed in São Paulo (Brazil). *Food Control*, 22:778-785.
- 22 Morheit, H. M., Belphiy, D., and Morheit El, A. (2013). Contamination métallique de *Papilinus acarne*, *Sardina pilchardus* et *Diododus vulgaris* de la côte Atlantique sud (Maroc). *Larviss* J., 14:131-148.
- 23 Mormede, S., and Davies, I. M. (2001a). Heavy metal concentrations in commercial deep-sea fish from the Rockall Trough. *Cont. Shelf Res.*, 21:899-916.
- 24 Mormede, S., and Davies, I. M. (2001b). Trace elements in deep-water fish species from the Rockall Trough. *Fisheries Res.*, 52:197-206.
- 25 Olmedo, P., Pla, A., Hernandez, A. F., Barber, F., Ayuñil, L., and Gil, F. (2013). Determination of toxic elements (mercury, cadmium, lead, tin and arsenic) in fish and shellfish samples. Risk assessment for the consumers. *Environ. Int.*, 59:63-72.
- 26 Olusola, J. O., and Festus, A. A. (2015). Assessment of heavy metals in some marine fish species relevant to their concentration in water and sediment from coastal waters of Ondo State, Nigeria. *Mar. Sci. res. Dev.*, 5(2):1-6.
- 27 Oehlenschläger, J. (2009). Trace element concentration in muscle tissue of the benthopelagic grenadier (*Coryphaenoides armatus*) from the Iberian deep-sea. *Inf. Fisherieforsch.*, 56:45-48.
- 28 Polak-Juszczak, L. (2009). Temporal trends in the bioaccumulation of trace metals in herring, sprat, and cod from the southern Baltic Sea in the 1994-2003 period. *Chemosphere*, 76:1334-1339.
- 29 Rodriguez, A., and Davies, I. M. (2001). Trace element concentrations in deep-water fish from the Rockall Trough. *Cont. Shelf Res.*, 21:899-916.
- 30 Stange, K., Massé, A., and Klungsoy, J. (1996). In: Contaminants in fish and sediments from the North Atlantic ocean (Tema Nord 1996-522). Copenhagen, Denmark: Nordic Council of Ministers. 79p.
- 31 Stoeppler, M., and Bandt, K. (1979). Comparative studies on trace metal levels in marine biota. II. Trace metals in krill, krill products, and fish from the Atlantic Scotia Sea. *Z. Lebensm. Unters. Forsch.*, 169:95-98.
- 32 Stoeppler, M., and Nürnberg, H. W. (1979). Comparative studies on trace metal levels in marine biota. III. Typical levels and accumulation of toxic trace metals in muscle tissue and organs of marine organisms from different European Seas. *Ecotoxicol. Environ. Safety*, 3:335-351.
- 33 Storelli, M., Normanno, G., Barone, G., Dambrasio, A., Errico, L., Garofalo, R., and Giacomelli-Stuffer, R. (2012). Toxic metals (Hg, Cd, and Pb) in fishery products imported into Italy: suitability for Human consumption. *J. Food Protec.*, 75(1):189-194.
- 34 Torres, P., Rodrigues, A., Soares, L., and Garcia, P. (2016). Metal concentrations in two commercial tuna species from an active volcanic region in the Mid-Atlantic Ocean. *Arch. Environ. Contam. Toxicol.*, 70:341-347.
- 35 Viera, C., Morais, S., Ramos, S., Delerue-Matos, C., and Oliveira, M. B. P. P. (2011). Mercury, cadmium, lead and arsenic levels in three pelagic fish species from the Atlantic Ocean: intra- and inter-specific variability and human health risks for consumption. *Food Chem. Toxicol.*, 49, 923-932.
- 36 Windom, H., Stein, D., Sheldon, R., and Smith, R. (1987). Comparison of trace metal concentrations in muscle tissue of a benthopelagic fish (*Coryphaenoides armatus*) from the Atlantic and Pacific Oceans. *Deep Sea Res.*, 34:213-220.




## REF


- 1 Agusa, T., Kunio, T., et al. (2007). Exposure assessment for trace elements from consumption of marine fish in Southeast Asia. *Environ. Poll.*, **145**:766-777.
- 2 Araujo, C. V. M., and Gómez-Macías, I. A. (2016). Heavy metals in yellowtail kingfish (*Seriola quinqueradiata*) and black jack mackerel (*Scomber japonicus*). *J. Environ. Sci.*, **77**(2):18-31.
- 3 Baek, J. H., Kim, S. H., et al. (2012). Heavy metal contents and chemical speciation in black mackerel (*Scomber japonicus*) muscle. *J. Food Sci.*, **77**(2):18-31.
- 4 Belotti, F. A., Shahzami-Ottman, M., and Madan, A. G. (2012). Evaluation of trace metal levels in tissues of two commercial fish species in Kasar and Morinti coastal waters, Peninsular Malaysia. *J. Environ. Pub. Health*, **ID352309**:1-14.
- 5 Bassari, A. (1994). A study on the trace element concentrations of *Thunnus thynnus*, *Thunnus obesus* and *Katsuwonus pelamis* in the Aegeans: location, age, size, and risk. *J. Toxicol. Environ. Health Part A*, **70**(22):1897-1911.
- 6 Burger, J., Goedfeld, M., et al. (2007). Heavy metals in Pacific cod (*Gadus macrocephalus*) from the Aleutians: location, age, size, and risk. *J. Toxicol. Environ. Health Part A*, **70**(22):1897-1911.
- 7 Frias-Experiencias, M. G., Zamora-Sarabia, F. K. G., et al. (2014). Cadmium, copper, lead, and zinc contents of fish marketed in NW Mexico. *The Sci. World J.*, **ID546897**:1-4.
- 8 Gibbs, P. J., and Madan, A. G. (1996). Trace element concentrations in fish from the northern Indian Ocean. *Mar. Poll. Bull.*, **30**(10):657-674.
- 9 Gu, Y.-G., Huang, Q., Li, H.-H., Wang, L.-G., Ning, J.-J., and Du, F.-Y. (2017) Heavy metals in fish tissue/stomach contents of four marine wild commercially valuable fish species from the western continental shelf of South China Sea. *Mar. Poll. Bull.*, **114**:1125-1129.
- 10 Han, B.-C., Jenq, W. L., Chen, R. Y., Fang, T. C., Hung, T. C., and Tseng, R. J. (1998). Estimation of target hazard quotients and potential risks for metals by consumption of seafood in Taiwan. *Arch. Environ. Contam. Toxicol.*, **35**:711-720.
- 11 Ikebe, K., Nishimura, T., and Tanaka, R. (1991). Contents of 17 metal elements in food determined by inductively coupled plasma atomic emission spectrometry-fish and shellfishes. *J. Food Hyg. Soc. Jpn.*, **32**(4):336-350.
- 12 Li, M., Wei, D., et al. (2020). Assessing lead sources in fishes of the northeast Pacific ocean. *Anthropocene*, **29**(D100234):1-12.
- 13 Madan, A. G., and Frias, M. G. (2007). Trace element concentrations in fish from the northern Indian Ocean. *Mar. Poll. Bull.*, **54**:417-420.
- 14 Madan, A. G., Frias, M. G., and Belotti, F. A. (1994). A comparison of the non-essential elements cadmium, mercury, and lead found in fish and sediment from Alaska and California. *Sci. Tot. Environ.*, **339**:189-205.
- 15 Merian, M., Warner, M., Chouvelon, T., Pedraza, F., Rodríguez y Barreiro, A. M., and Bustamante, P. (2013). Trace element bioaccumulation in reef fish from New Caledonia: influence of trophic groups and risk assessment for consumers. *Mar. Environ. Res.*, **87**:88-96.
- 16 Ono, M. C., Norazirah, A. Z., Noor, A. M. S., and Kamaluzaman, Y. (2018). Selected heavy metals content in commercial fishes at different season landed at fisheries development authority of Malaysia complex (LKIM), Kuala Terengganu, Malaysia. *J. Sustain. Sci. Manag.*, **13**(5):29-38.
- 17 Patterson, C. C., and Settle, D. (1977). Comparative distribution of alkaline earths and lead among main tissues of the tuna *Thunnus alalunga*. *Mar. Biol.*, **39**:289-295.
- 18 Powell, J. H., and Powell, R. E. (2001). Trace element in fish overlying substrate in the tropical West Pacific. *Water Air Soil Poll.*, **125**:81-88.
- 19 Powell, J. H., and Powell, R. E. (1988). Trace element concentrations in fish from the Maldives. *Arch. Environ. Poll.*, **16**:143-158.
- 20 Ruelas-Inzunza, J. (2001). Concentración y distribución de metales pesados en tejidos de organismos representativos de la flora y fauna del bío océano de California y áreas adyacentes. Thesis, Univ. Nacional Autónoma de México, Mexico. 1996.
- 21 Ruelas-Inzunza, J., Soto-Jiménez, M.F., Ruiz-Fernández, A.C., Bojórquez-Leiva, H., Pérez-Bernal, H., and Paiz-Osuna, F. (2012). 210Po activity and concentrations of selected trace elements (As, Cd, Cu, Hg, Pb, Zn) in the muscle tissue of tunas *Thunnus albacares* and *Katsuwonus pelamis* from the Eastern Pacific Ocean. *Biol. Trace Elem. Res.*, **149**:371-376.
- 22 Ruelas-Inzunza, J., Soto-Jiménez, M.F., Ruiz-Fernández, A.C., Ramos-Usuna, M., Mones-Saucedo, J., and Paiz-Osuna, F. (2014). 210Po, Cd and Pb distribution and biomagnification in the yellowfin tuna *Thunnus albacares* and skipjack tuna *Katsuwonus pelamis* from the Eastern Pacific. *Mar. Poll. Bull.*, **87**:98-103.
- 23 Sangalang, R. H., and Quinay, E. B. (2015). Lead content of round scad (*Decapterus macromelas*) from Batangas Bay, Philippines. *Asia Pac. J. Multi. Res.*, **3**(4):118-124.
- 24 Storelli, M., Normanno, G., Barone, G., Damascio, A., Errico, L., Garofalo, R., and Giacomelli-Stuffer, R. (2012). Toxic metals (Hg, Cd, and Pb) in fishery products imported into Italy: suitability for Human consumption. *J. Food Protec.*, **75**(1):189-194.
- 25 Windom, H., Stein, D., Sheldon, R. & Smith, R. (1967). Comparison of trace metal concentrations in muscle tissue of a benthopelagic fish (*Cymophyes armatus*) from the Atlantic and Pacific Oceans. *Deep Sea Res.*, **34**:213-220.



## REF

- 1 Ahmed, Q., Yousuf, F., Turkmen, M., Tabassum, S., and Khoshnood, R. (2013). Trace elemental contents in tissues of talang queenfish, *Scomberoides commersonianus*, from Karachi coast, Pakistan. *The Black Sea J. Sci.*, 3(8):68-76.
- 2 Ahmed, N. S., El-Deek, M. S. and Emara, H. I. (1996). Heavy metals in the muscle and bone of some fish species from the Red Sea. *J. KAU. Mar. Sci.*, 7:25-31.
- 3 Ahmed, Q., Bat, L., and Ali, Q. M. (2012). Heavy metal levels in *Thunnus albacares* (Bonnaterre, 1788) from Karashi fish harbour, Pakistan. *Pak. J. Mar. Sci.*, 21(1-2):13-21.
- 4 Ahmed, Q., Yousuf, F. et al. (2015). *Euthynnus affinis* (little tuna): fishery, biomics, seasonal elemental variations, health risk assessment and conservational management. *Front. Life Sci.*, 8(1):71-96.
- 5 Ahmed, Q., Benzer, S., and Yousuf, F. (2016). Distribution of heavy metals in different tissues of Indian mackerel from Karachi fish harbour, Karachi, Pakistan. *Ind. J. Anim. Res.*, 50(5):759-763.
- 6 Al-Shwafi, N. A. (2002). Heavy metals concentration levels in some fish species in the Red Sea and Gulf of Aden-Yemen. *Catar Univ. Sci.*, 22:171-176.
- 7 Al-Zaidan, A. S., Al-Sarawi, H. A., et al. (2015). Histopathology and contaminant concentrations in fish from Kuwait's marine environment. *Mar. Poll. Bull.*, 100:637-645.
- 8 Bastami, K. D., Afkhami, M., et al. (2015). Bioaccumulation and ecological risk assessment of heavy metals in the sediments and mullet *Liza lunuligera* in the northern part of the Persian Gulf. *Mar. Poll. Bull.*, 94:329-334.
- 9 Bosch, A. C. (2015). Status of mercury and other heavy metals in South African marine fish species. Thesis, Faculty of AgriSciences, Stellenbosch Univ., South Africa. 149p.
- 10 Bu-Olayan, A. H., and Subrahmanyam, M. N. V. (1996). Trace metals in fish from the Kuwait coast using microwave acid digestion technique. *Environ. Int.*, 22(6):753-758.
- 11 Deshpande, A., Bhendiger, S., Shirsiker, T., Dhaware, D., and Khandekar, R. N. (2009). Analysis of heavy metals in marine fish from Mumbai docks. *Environ. Monit. Assess.*, 159:493-500.
- 12 Dobardaran, S., Naddaf, K., Nazmara, S., and Ghaedi, H. (2010). Heavy metals (Cd, Cu, Ni and Pb) content in two species of Persian Gulf in Bushehr port, Iran. *Afr. J. Biotech.*, 9(37):6191-6193.
- 13 El-Bahr, S. M., and Abdelfahy A. (2015). Heavy metals and trace element contents in edible muscle of three commercial fish species, and assessment of possible risks associated with their human consumption in Saudi Arabia. *J. Adv. Vet. Anim. Res.*, 2(3):271-278.
- 14 El-Moselly, Kh. M., Othman, A. I., El-Azem, H. A., and El-Metwally, M. E. A. (2014). Bioaccumulation of heavy metals in some tissues of fish in the Red Sea, Egypt. *Egypt. J. Basic Appl. Sci.*, 1(2):97-105.
- 15 Ganjavi, M., Ezzatpanah, H., Givannat, M. H., and Shams, A. (2010). Effect of canned tuna fish processing steps on lead and cadmium contents of Iranian tuna fish. *Food Chem.*, 118:525-528.
- 16 Hanna, R. G. M. (1989). Levels of heavy metals in some Red Sea fish before Hot Brine pools' mining. *Mar. Poll. Bull.*, 20:631-635.
- 17 Hasimah, N. A. K., Noik, J. V., Teh, Y. Y., Lee, C. Y., and Ng, P. H. C. (2011). Assessment of cadmium (Cd) and lead (Pb) levels in commercial marine fish organs between wet markets and supermarkets in Klang Valley, Malaysia.
- 18 Honda, K., Sahrui, M., Hidaka, H., and Tatsukawa, R. (1983). Organ and tissue distribution of heavy metals, and their growth-related changes in Antarctic fish, *Pagothenia borchgrevinkii*. *Agicul. Biol. Chem.*, 47(11): 2521-2532.
- 19 Hosseini, M., Nabavi, S. M. B., Nabavi, S. N., and Pour, N. A. (2015). Heavy metals (Cd, Co, Cu, Ni, Pb, Fe, and Hg) content in four fish commonly consumed in Iran; risk assessment to the consumers. *Environ. Monit. Assess.*, 187(237):1-7.
- 20 Jaffar, M., and Ashraf, M. (1988). Selected trace metal concentrations in different tissues of fish from coastal waters of Pakistan (Arabian Sea). *Ind. J. Mar. Sci.*, 17:231-234.
- 21 Khoshnood, Z., Khoshnood, R., Mokhlesi, A., Ehsanpour, M., Afkhami, M., and Khazaali, A. (2012). Determination of Cd, Pb, Hg, Cu, Fe, Mn, Al, As, Ni and Zn in important commercial fish species in northern of Persian Gulf. *J. Cell. Anim. Biol.*, 6(1):1-9.
- 22 Kojadinicovic, J., Potier, M., Le Corre, M., Cosson, R., and Bustamante, P. (2007). Bioaccumulation of trace elements in pelagic fish from the Western Indian Ocean. *Environ. Poll.*, 145(2):548-566.
- 23 Krishnamurti, A. J., and Nair, V. R. (1999). Concentration of metals from Thane and Bassein creeks of Bombay. *Ind. J. Mar. Sci.*, 28:39-44.
- 24 Kureishi, T. W. (1993). Concentration of heavy metals in marine organisms around Qatar before and after the gulf war oil spill. *Mar. Poll. Bull.*, 27:183-186.
- 25 De Mora, S., Fowler, S. W., Wyse, E., and Azemard, S. (2004). Distribution of heavy metals in marine bivalves, fish and coastal sediments in the Gulf and Gulf of Oman. *Mar. Poll. Bull.*, 49:410-424.
- 26 Mortazavi, M. S., and Sharifian, S. (2012). Metal concentrations in two commercial fish from Persian Gulf, in relation to body length and sex. *Bull. Environ. Contam. Toxicol.*, 89:450-454.
- 27 Naji, A., Khan, F. R., and Hashemi, S. M. (2016). Potential human health risk assessment of trace metals via the consumption of marine fish in Persian Gulf. *Mar. Poll. Bull.*, 109:667-671.
- 28 Niri, A. S., Sharifian, S., and Ahmadi, R. (2015). Assessment of metal accumulation in two fish species (*Tenualosa ilisha* and *Otolithes ruber*), captured from the north of Persian Gulf. *Bull. Environ. Contam. Toxicol.*, 94:71-76.
- 29 Norouzi, M., Mansouri, B., Hamidian, A. H., Zarei, I., and Mansouri, A. (2012). Metal concentrations in tissues of two fish species from Qeshm Island, Iran. *Bull. Environ. Contam. Toxicol.*, 89:1004-1008.
- 30 Pourang, N., Nikouyan, A., and Dennis, J. H. (2005). Trace element concentrations in fish, surficial sediments and water from northern part of the Persian Gulf. *Environ. Monitor. Assess.*, 109:293-316.
- 31 Reddy, M. S., Mehta, B., et al. (2007). Bioaccumulation of heavy metals in some commercial fishes and crabs of the gulf of Cambay, India. *Current Sci.*, 92(11):1489-1491.
- 32 Rejomon, G., Nair, M., and Joseph, T. (2010). Trace metal dynamics in fishes from the southwest coast of India. *Environ. Monit. Assess.*, 67(1-4):243-255.
- 33 Saei-Dehkordi, S. S., and Fallah, A. A. (2011). Determination of copper, lead, cadmium and zinc content in commercially valuable fish species from the Persian Gulf using derivative potentiometric stripping analysis. *Microchem. J.*, 98:156-162.
- 34 Sharif, A. K. M., Mustafa, A. I., Hossain, M. A., Amin, M. N., and Saifullah, S. (1993). Lead and cadmium contents in ten species of tropical marine fish from the Bay of Bengal. *The Sci. Tot. Environ.*, 133:193-199.
- 35 Sivarupam, P., Sankar, T. V., and Nair, P. G. V. (2006). Heavy metal concentrations in fish, shellfish and fish products from internal markets of India via-à-vis international standards. *Food Chem.*, 102:612-620.
- 36 Tarig, J., Jaffar, M., Ashraf, M., and Moazzam, M. (1993). Heavy metal concentrations in fish, shrimp, seaweed, sediment, and water from the Arabian Sea, Pakistan. *Mar. Poll. Bull.*, 24(124):644-647.
- 37 Tarig, J., Ashraf, M., Jaffar, M., and Masud, K. (1998). Selected trace metal concentrations in seven fish species from the Arabian Sea, Pakistan. *J. Chem. Soc. Pak.*, 20(4):249-251.
- 38 Thiagarajan, D., Dhaneesh, K. V., Ajith Kumar, T. T., Kumaresan, S., and Balasubramanian, T. (2012). Metals in fish along the southeast coast of India. *Bull. Environ. Contam. Toxicol.*, 88:582-588.
- 39 Yasmeen, K., Mirza, M. A., Khan, N. A., Kausar, N., Rheman, A.-U., and Hanif, M. (2016). Trace metals health risk appraisal in fish species of Arabian Sea. *SpringerPlus*, 5(1):859.





## REF

- 1 Abdallah, M. A. H. (2008). Trace element levels in some commercially valuable fish species from coastal waters of Mediterranean Sea, Egypt. *J. Mar. Syst.*, 73:114-122.
- 2 Aksu, A., Balkis, N., Taskin, O. S., and Ersan, M. S. (2011). Toxic metal (Pb, Cd, As and Hg) and organochlorine pesticide residue levels in hake (*Merluccius merluccius*) from the Marmara Sea, Turkey. *Environ. Monit. Assess.*, 182:509-521.
- 3 Alkan, N., Aktas, M., and Gedik, K. (2012). Comparison of metal accumulation in fish species from the southeastern Black Sea. *Bull. Environ. Contam. Toxicol.*, 88:807-812.
- 4 Alkan, N., Alkan, A., Gedik, K., and Fisher, A. (2016). Assessment of metal concentrations in commercially important fish species in Black Sea. *Toxicol. Ind. Health*, 32(3):447-456.
- 5 Bilandzic, H., Dokic, M., and Sedak, M. (2011). Metal content determination in four fish species from the Adriatic Sea. *Food Chem.*, 124:1005-1010.
- 6 Canli, M., and Atli, G. (2003). The relationships between heavy metal (Cd, Cr, Cu, Fe, Pb, Zn) levels and the size of six Mediterranean fish species. *Environ. Poll.*, 121:129-136.
- 7 Celik, U., Cakici, S., and Oehlenschlager, J. (2004). Determination of the lead and cadmium burden in some northeastern Atlantic and Mediterranean fish species by DPSV. *Eur. Food Res. Technol.*, 218:298-305.
- 8 Chouvelon, T., Strady, E., et al. (2019). Patterns of trace metal bioaccumulation and trophic transfer in a phytoplankton-zooplankton-small pelagic fish marine food web. *Mar. Poll. Bull.*, 146:1013-1030.
- 9 Cogun, H. Y., Yuzeferoglu, T. A., Firat, O., Gok, G., and Kargin, F. (2006). *Environ. Monit. Assess.*, 121:431-438.
- 10 Cogun, H. Y., Yuzeferoglu, T. A., Firat, O., Gok, G., and Kargin, F. (2005). Seasonal variation and tissue distribution of heavy metals in shrimp and fish species from the Yumurtalik coast of Iskenderun Gulf, Mediterranean. *Environ. Contam. Toxicol.*, 75:707-715.
- 11 Copati, C., Bella, F., Castagna, M., Fallico, R., Sciacca, S., and Ferrante, M. (2012). Heavy metals concentrations in fish from Sicily (Mediterranean Sea) and evaluation of possible health risks to consumers. *Bull. Environ. Contam. Toxicol.*, 88:78-83.
- 12 Copati, C., Arena, G., et al. (2013). Heavy metals concentrations in fish and shellfish from eastern Mediterranean Sea: Consumption advisories. *Food Chem. Toxicol.*, 53:37-37.
- 13 Curcu, A. K., Topkaya, M., Erdogan, G., and Aboul Enein, H. n. (2018). Quantitative determination of heavy metal contamination in horse mackerel and whiting caught in the Sea of Marmara. *Bull. Environ. Contam. Toxicol.*, 102:498-503.
- 14 Engin, M. S. (2015). The assessment of trace metals at gill, muscle and liver tissues in Mugil cephalus. *Environ. Monit. Assess.*, 187(255):1-6.
- 15 Ergul, H. A., and Aksan, S. (2013). Evaluation of non-essential element and micronutrient concentrations in seafood from the Marmara and Black Seas. *J. Black Sea/Med. Environm.*, 19(3):312-330.
- 16 Erkan, N., Ozden, O., and Ulusoy, S. (2009). Levels of trace elements in commercially important fish, crustaceans and mollusks from Istanbul fish market. *Trace Environ. Bull.*, 18(7b):1307-1311.
- 17 Ersoy, B., and Celik, M. (2010). The essential and toxic elements in tissues of six demersal fish from Eastern Mediterranean Sea. *Food Chem. Toxicol.*, 48:1377-1382.
- 18 Ferrante, M., Pappalardo, A. M., et al. (2017). Bioaccumulation of metals and biomarkers of environmental stress in *Parablemmis sanguinolentus* (Pallas, 1814) sampled along the Italian coast. *Mar. Poll. Bull.*, 122:288-296.
- 19 Findik, D., and Cicek, E. (2011). Metal Concentrations in Two Bioindicator Fish Species, *Merlangius merlangus*, *Mullus barbatus*, Captured from the West Black Sea Coasts (Bartin) of Turkey. *Bull. Environ. Contam. Toxicol.*, 87:399-403.
- 20 Guner, S., Dincer, B., Alemdag, N., Colak, A., and Tufekci, M. (1998). Proximate composition and selected mineral content of commercially important fish species from the Black Sea. *J. Sci. Food Agric.*, 78:337-342.
- 21 Iamigeli, A. I., Ubaldi, A., et al. (2013). Metals in Mediterranean aquatic species. *Mar. Poll. Bull.*, 94:278-283.
- 22 Kalay, M., Ay, O., and Canli, M. (1999). Heavy metal concentrations in fish tissues from the Northeastern Mediterranean Sea. *Bull. Environ. Contam. Toxicol.*, 63:673-681.
- 23 Kalogeropoulos, N., Karavoltos, S., Sakellari, A., Avramidou, S., Dassanakis, M., Scoullos, M. (2012). Heavy metals in raw, fried and grilled Mediterranean finfish and shellfish. *Food Chem. Toxicol.*, 50:3702-3708.
- 24 Kargin, F. (1996). Seasonal changes in levels of heavy metals in tissues of *Mullus barbatus* and *Sparus aurata* collected from Iskenderun Gulf (Turkey). *Water Air Soil Poll.*, 90:557-562.
- 25 Keskin, Y., Baskaya, R., Ozyar, O., Yurdun, T., Luleci, N. E., Hayran, O. (2007). Cadmium, lead, mercury and copper in fish from the Marmara Sea, Turkey. *Bull. Environ. Contam. Toxicol.*, 78:258-261.
- 26 Khaled, A. (2004). Heavy metals concentrations in certain tissues of five commercially important fishes from El-Mex Bay, Alexandria, Egypt. *Egypt. J. Aquat. Biol. Fish.*, 8(1):51-64.
- 27 Kuruksezer, F., Altay, O., Uluturhan, E., and Konata, A. (2001). Trace metal and organochlorine residue levels in red mullet (*Mullus barbatus*) from the eastern Aegean, Turkey. *Wat. Res.*, 35(9):2327-2332.
- 28 Kuruksezer, F., Uluturhan, E., Konata, A., and Altay, O. (2002). Trace metal concentrations in edible fishes from Izmir Bay, Eastern Aegean. *Mar. Poll. Bull.*, 44:816-832.
- 29 Licata, P., Trombetta, D., et al. (2005). Heavy metals in liver and muscle of bluefin tuna (*Thunnus thynnus*) caught in the straits of Messina (Sicily, Italy). *Environ. Monit. Assess.*, 107:239-248.
- 30 Mendl, D., Demirkiz, Z., Tuzen, M., and Soylik, M. (2010). Seasonal investigation of trace element contents in commercially valuable fish species from the Black sea, Turkey. *Food Chem.*, 48(3):865-870.
- 31 Nesto, N., Romano, S., Moschino, V., Mauri, M., and Da Ros, L. (2007). Bioaccumulation and biomarker responses of trace metals and micro-organic pollutants in mussels and fish from the Lagoon of Venice, Italy. *Mar. Poll. Bull.*, 55:469-484.
- 32 Nisbet, C., Terzi, G., Pilgr, O., and Sarac, N. (2010). Determination of Heavy Metal Levels in Fish Samples Collected from the Middle Black Sea. *Kafkas Univ. Vet. Fak. Derg.*, 16(1):119-126.
- 33 Olmedo, P., Pla, A., Hernandez, A. F., Barber, F., Ayuso, L., and Gil, F. (2013). Determination of toxic elements (mercury, cadmium, lead, tin and arsenic) in fish and shellfish samples. Risk assessment for the consumers. *Environ. Int.*, 59:63-72.
- 34 Ourgaud, M. (2015). Influence des apports anthropiques sur les flux de carbone et de contaminants dans les réseaux trophiques de poissons à Posidonia oceanica. Thesis, Aix-Marseille Univ, France. 353p.
- 35 Pastor, A., Hernandez, F., Prat, M. A., Bellan, J., Sandio, J. V., and Castillo, M. T. (1994). Levels of heavy metals in some marine organisms from the Western Mediterranean area (Spain). *Mar. Poll. Bull.*, 28(1):50-53.
- 36 Pazi, I., Gonul, L. T., et al. (2017). Potential risk assessment of metals in edible fish species for human consumption from the Eastern Aegean Sea. *Mar. Poll. Bull.*, 120(1-2):409-413.
- 37 Perugini, G., Visiciano, P., Manera, M., Zaccaroni, A., Olivieri, V., and Amorena, M. (2014). Heavy metal (As, Cd, Hg, Pb, Cu, Zn, Se) concentrations in muscle and bone of four commercial fish caught in the central Adriatic Sea, Italy. *Environ. Monit. Assess.*, 186:2205-2213.
- 38 Plavan, G., Itar, O., Teodosiu, C., Nicara, M., Micu, D., and Strangaru, S. A. (2017). Toxic metals in tissues of fishes from the Black Sea and associated human risk exposure. *Environ. Sci. Poll. Res.*, 24(8):7776-7787.
- 39 Ramos-Miras, J. J., Sanchez-Muros, M. J., et al. (2019). Potentially toxic elements in commonly consumed fish species from the western Mediterranean Sea (Almeria Sea): Bioaccumulation in liver and muscle tissues in relation to biometric parameters. *Sci. Tot. Environ.*, 671:280-287.
- 40 Sofuoğlu, K., Kalanlı, I., et al. (2018). Metals and elements in sardine and anchovy: species specific differences and correlations with proximate composition and size. *Sci. Tot. Environ.*, 645:229-238.
- 41 Stamatios, N., Kamidis, N., Pigida, P., Stergiou, D., and Kalianniotis, A. (2019). Bioaccumulation levels and potential health risks of mercury, cadmium, and lead in abalone (*Thunus alalunga*; Bonnetier, 1788) from the Aegean Sea, Greece. *Environ. Res. Pub. Health.*, 16 (5):821.
- 42 Stoeppler, M., and Nürnberg, H. W. (1979). Comparative studies on trace metal levels in marine biota: III. Typical levels and accumulation of toxic trace metals in muscle tissue and organs of marine organisms from different European Seas. *Ecotox. Environ. Safety.*, 3:335-351.
- 43 Storelli, M. M. (2008). Potential human health risks from metals (Hg, Cd, and Pb) and polychlorinated biphenyls (PCBs) via seafood consumption: Estimation of target hazard quotients (THQs) and toxic equivalents (TEQs). *Food Chem. Toxicol.*, 46:2782-2788.
- 44 Storelli, M. M., Giacomelli-Stuffer, R., Storelli, A., and Marcorigiani, G. O. (2005). Accumulation of mercury, cadmium, lead and arsenic in swordfish and bluefin tuna from the Mediterranean Sea: A comparative study. *Mar. Poll. Bull.*, 50:993-1003.
- 45 Storelli, M., Barone, G., Cuttino, G., Giungato, D., and Garofalo, R. (2010). Occurrence of toxic metals (Hg, Cd and Pb) in fresh and canned tuna: Public health implications. *Food Chem. Toxicol.*, 48:3167-3170.
- 46 Tepe, Y., Turkmen, M., and Turkmen, A. (2008). Assessment of heavy metals in two commercial fish species of four Turkish seas. *Environ. Monit. Assess.*, 146:277-284.
- 47 Topcuoglu, S., Kirbasoglu, C., and Gungor, N. (2002). Heavy metals in organisms and sediments from Turkish Coast of the Black Sea, 1997-1998. *Environ. Int.*, 27:521-526.
- 48 Turan, C., Dural, M., Okuz, A., and Orturk, B. (2009). Levels of heavy metals in some commercial fish species captured from the Black Sea and Mediterranean coast of Turkey. *Bull. Environ. Contam. Toxicol.*, 82:601-604.
- 49 Turkmen, A., Turkmen, M., Tepe, Y., Mazlum, Y., and Oymael, S. (2006). Metal concentrations in blue crab (*Callinectes sapidus*) and mullet (*Mugil cephalus*) in Iskenderun Bay, Northern East Mediterranean, Turkey. *Bull. Environ. Contam. Toxicol.*, 77:186-193.
- 50 Turkmen, M., Turkmen, A., Tepe, Y., Ates, A., and Golkus, K. (2008). Determination of metal contaminations in sea foods from Marmara, Aegean and Mediterranean seas: Twelve fish species. *Food Chem.*, 108:794-800.
- 51 Turkmen, A., Tepe, Y., Turkmen, M., and Mutin, E. (2009). Heavy metal contaminations in tissues of the garfish, Belone belone L., 1761, and the bluefish, *Pomatomus saltatrix* L., 1766, from Turkey waters. *Bull. Environ. Contam. Toxicol.*, 82:70-74.
- 52 Turkmen, A., Turkmen, M., Tepe, Y., and Alykti I. (2005). Heavy metals in three commercially valuable fish species from Iskenderun Bay, Northern East Mediterranean Sea, Turkey. *Food Chem.*, 91:167-172.
- 53 Turkmen, M., Turkmen, A., and Tepe, Y. (2008). Metal contamination in five fish species from Black, Marmara, Aegean and Mediterranean Seas, Turkey. *J. Chil. Chem. Soc.*, 53(1):1425-1439.
- 54 Turkmen, M., Turkmen, A., Tepe, Y., Tore, Y., and Ates, A. (2009). Determination of metals in fish species from Aegean and Mediterranean Seas. *Food Chem.*, 113:233-237.
- 55 Tuzen, M. (2003). Determination of heavy metals in fish samples of the Middle Black sea (Turkey) by graphite furnace atomic absorption spectrometry. *Food Chem.*, 80:119-123.
- 56 Tuzen, M. (2009). Toxic and essential trace elemental contents in fish species from the Black Sea, Turkey. *Food Chem.*, 47(8):1785-1790.
- 57 Uluozlu, O. D., Tuzen, M., Mendl, D., and Soylak, M. (2007). Trace metal content in nine species of fish from the Black and Aegean Seas, Turkey. *Food Chem.*, 104:835-840.
- 58 Yabani, M., and Alparslan, Y. (2015). Potential health hazard assessment in terms of some heavy metals determined in demersal fishes caught in Eastern Aegean Sea. *Bull. Environ. Contam. Toxicol.*, 95:494-498.
- 59 Yilmaz, A. B., Sangun, M. K., Yaghoglu, D., and Turan, C. (2010). Metals (major, essential to non-essential) composition of the different tissues of three demersal fish species from Iskenderun Bay, Turkey. *Food Chem.*, 123:410-415.
- 60 Yilmaz, A. B. (2003). Levels of heavy metals (Fe, Cr, Ni, Cr, Pb, And Zn) in tissue of *Mugil cephalus* and T. *Mediterraneus* from Iskenderun Bay, Turkey. *Environ. Res.*, 92:277-281.
- 61 Yilmaz, A. B. (2005). Comparison of heavy metal levels of grey mullet (*Mugil cephalus* L.) and sea bream (*Sparus aurata* L.) caught in Iskenderun Bay (Turkey). *Turk. J. Vet. Anim. Sci.*, 29:257-262.
- 62 Yipel, M., and Yarsan, E. (2014). A risk assessment of heavy metal concentrations in fish and an invertibrate from the Gulf of Antalya. *Bull. Environ. Contam. Toxicol.*, 93:542-548.
- 63 Yuslu, V., Suelves, T., Ruiz-Atienza, L., Cervera, M. L., Benedito, V., and Pastor, A. (2008). Monitoring programme on cadmium, lead and mercury in fish and seafood from Valencia, Spain: levels and estimated weekly intake. *Food Add. Contam.*, 1(1):22-31.