



External Geophysics, Climate

Middle and Late Pennsylvanian cyclothem, American Midcontinent: Ice-age environmental changes and terrestrial biotic dynamics

C. Blaine Cecil^{a,b}, William A. DiMichele^{b,*}, Scott D. Elrick^c^a U.S. Geological Survey, 425 Brownsburg Turnpike, Rockbridge Baths, VA 20743, United States^b Department of Paleobiology MRC-121, NMNH Smithsonian Institution, Washington, DC 20560, United States^c Illinois State Geological Survey, University of Illinois, Champaign, IL 61820, United States

ARTICLE INFO

Article history:

Received 13 January 2014

Accepted after revision 13 March 2014

Available online 10 August 2014

Keywords:

Climate

Coal

Cyclothem

Paleoecology

Pennsylvanian

ABSTRACT

The Pennsylvanian portion of the Late Paleozoic Ice Age was characterized by stratigraphic repetition of chemical and siliciclastic rocks in the equatorial regions of the Pangean interior. Known as “cyclothem”, these stratigraphic successions are a 10^5 yr-record of glacial waxing and waning, superimposed on longer term, 10^6 yr intervals of global warming and cooling and a still longer term trend of increasing equatorial aridity. During periods of maximum ice–minimum sea level, the interior craton was widely exposed. Epicontinental landscapes were initially subjected to dry subhumid climate when first exposed, as sea level fell, but transitioned to humid climates and widespread wetlands during maximum lowstands. During interglacials (ice-minima) seasonally dry vegetation predominated. The wetland and seasonally dry biomes were compositionally distinct and had different ecological and evolutionary dynamics.

Published by Elsevier Masson SAS on behalf of Académie des sciences.

1. Introduction

The Pennsylvanian portion of the Late Paleozoic Ice Age (approx. 323–299 Ma) was characterized by regular waxing and waning of Southern Hemisphere continental glaciers (Fielding et al., 2008; Isbell et al., 2003a). In order to provide a precise definition of time scales for Earth's warming and cooling events, we use terminology as outlined in Cecil (2013). **Glacial stage** (shortest scale of a single glacial–interglacial cycle, 10^5 yr) pacing may have been on the rhythm of Milankovich-band orbital frequencies (Heckel, 2008). These individual glacial–interglacial cycles were superimposed on **glacial epoch** scale (longer-term, 10^6 yr) intervals of global warming and cooling (Birgenheier et al., 2009; Joeckel, 1999), further superimposed on a **period-scale** trend of long-term equatorial

drying and warming (10^7 yr) (Tabor and Poulsen, 2008). Regularity of climate and sea-level changes at the glacial-stage scale has permitted correlation across North America (Cecil et al., 2003a) and even across the Euramerican portions of Pangea in deposits of paralic (marine influenced) basins from the American Midcontinent to the Donets Basin (Eros et al., 2012; Heckel et al., 2007). Patterns of equatorial sea-level change strongly correlate with inferred polar ice volume on both stage and epoch temporal scales (Rygel et al., 2008).

Equatorial climate changed in concert with ice volume and sea level, most notably the patterns, durations and amounts of equatorial rainfall (Fig. 1B). These changes occurred on all time scales from glacial–interglacial stages (Cecil et al., 2003a; Horton et al., 2012), to epoch-scale intervals of global warming and cooling (e.g., Cecil, 1990; Rygel et al., 2008), to a period-scale, long-term trend of warming and drying (Cecil, 1990; Montañez and Poulsen, 2013; Tabor and Poulsen, 2008). Climate change has been

* Corresponding author.

E-mail address: dimichel@si.edu (W.A. DiMichele).




Fig. 1. (Color online.) Interior Pangean cyclothem. A. Stratigraphic architecture of a complete glacial–interglacial, stage-scale cycle. 1–paleosol; 2–coal; 3–estuarine gray-shale wedge; 4–ravinement surface; 5–marine black shale; 6–marine limestone; 7–deltaic and nearshore siliciclastics. B. Patterns of change in ice volume, sea level, equatorial climate. C. Equatorial floristic changes, tracking climate change. D. Pennsylvanian Pangea. E. Climate–Siliciclastic sediment yield relationship. F. Climate–Sediment type relationship.

integrated into the broader understanding of sedimentary and biological dynamics of the Late Paleozoic Ice Age (LPIA). Climate has a strong, direct effect on lithofacies patterns in both limnic (non-marine) and paralic basins. Lithologies such as coal (formerly peat—a Histosol) and mineral paleosols directly record climatic conditions. There are strong climatic effects on siliciclastic sediment availability and transport (Cecil et al., 2003b) (Fig. 1E), the extent to which environments of deposition are mixed and oxygenated, and the chemical conditions. Where siliciclastic input is absent or minimal, and where necessary chemical and physical conditions are met, climate becomes an important control on carbonate formation (Fig. 1F) (Cecil and Dulong, 2003; Cecil et al., 2003b).

Pangean plants and animals also strongly reflected and tracked environments (Fig. 1C), at all spatio-temporal

scales (DiMichele et al., 2009, 2010; Falcon-Lang and DiMichele, 2010; Opluštil et al., 2013). Thus, many paleontological patterns, including evolutionary dynamics, are direct reflections of regional climatic patterns, controlled by global-scale factors. Here we review the lithological signature and biological effects of LPIA glacial–interglacial cyclicity on landscapes of the vast, flat, central-western portions of the Pangean supercontinent, between the Central Pangean Mountains and Ancestral Rockies (Fig. 1D). These patterns are expressed primarily during the late Middle and Late Pennsylvanian in the Western Interior (Midcontinent) and Eastern Interior (Illinois) basins, USA, more so than in the Appalachian basin. It is there that the “cyclothem” concept developed (see Cecil et al., 2003a; Heckel, 1990; Langenheim and Nelson, 1992; Weller, 1931). This region is well suited for

revealing ice-age climate and sea level dynamics in equatorial latitudes: Paralic character and low elevations permitted far reaching sea-level fluctuations. Great distance from mountain ranges in the eastern and western interior areas greatly reduced habitat variation and effects of uplands on climate and biological patterns. Great flatness of the cratonic surface created widespread environmental spatio-temporal uniformity.

2. Record and dynamic drivers of glacial–interglacial cycles

2.1. *The Cyclothem*

Glacial–interglacial cyclicity has a distinctive, if geographically and environmentally variable, lithological signature. Such successions can be entirely marine (e.g., [Elrick and Scott, 2010](#)), entirely terrestrial (e.g., [Eble et al., 2006](#)), or mixed (e.g., [Heckel, 2008](#)), and subject to local tectonic, and climatic overprints. We focus on mixed terrestrial–marine cyclothem, as found in cratonic settings, particularly of the Eastern Interior basin USA, located between the more marine influenced Western Interior and more terrestrial Appalachian basins. Over time, the high relief, eroded surface of the Mid-Carboniferous unconformity ([Bristol and Howard, 1974](#)) was in-filled across the American Midcontinent. This lowered the relief of the cratonic platform over which glacial–interglacial cycles were expressed (e.g. [McKee and Crosby, 1975](#); [Watney et al., 1989](#)), resulting in alternation of marine and terrestrial environments in the Western and Eastern Interior regions of Pangea during each glacial–interglacial cycle ([Archer et al., 1994a](#)); here marine rocks generally dominate total stratigraphic thickness of a cycle ([Fig. 1A](#)).

Terrestrial facies include mineral paleosols ([Fig. 1A-1](#)), developed on previously deposited sediments, often succeeded by Histosols (peat, now coal) ([Fig. 1A-2](#)). Siliciclastic deposits with fluvial channel-form geometries may occur at the same stratigraphic level as the mineral paleosol, interpreted as drainages on the terrestrial landscape contemporaneous with the paleosol surface ([Davies and Gibling, 2013](#)). Plant fossils indicative of seasonally dry conditions may occur in these fluvial deposits, but usually are rare ([Feldman et al., 2005](#)). During peat development, some channels continued to be active, likely as blackwater rivers, transitioning to mud-filled estuaries during transgression ([Archer et al., 2014](#)). Estuarine tidal deposits, referred to as Gray Siltstone Wedges (GSW) ([Archer and Kvale, 1993](#)), may be present locally between the coal bed and overlying marine strata ([Fig. 1A-3](#)). GSWs are laterally continuous with deposits that backfill the courses of the former drainages through the peat swamp ([Elrick and Nelson, 2010](#)). Local gradational contacts between the coal bed and the base of the GSW are common and often include upright, fossil trees, and rich accumulations of plant compression fossils representative of wetland environments ([Archer et al., 1994b](#)).

The base of the marine portion of a cyclothem is marked by a transgressive erosional surface (a ravinement) above the coal bed and GSW ([Liu and Gastaldo, 1992](#)). This

surface may be overlain by a relatively thin “transgressive” marine limestone ([Heckel, 2008](#)), which is more common in Western Interior cycles than in the Eastern Interior or Appalachians. Variably developed, the transgressive limestone is often absent or represented only by shell hash. Where limestone is absent, the ravinement surface may be marked by local accumulations of phosphatic nodules or pyrite-permineralized plant remains. A marine, black, generally fissile shale typically lies in sharp, erosional contact with the underlying strata, immediately above the ravinement surface ([Fig. 1A-4](#)) and may be extremely widespread (e.g., [Cecil et al., 2003a](#); [James and Baker, 1972](#)). In the Western and Eastern Interior basins, black shale is overlain by open-marine limestone in conformable contact ([Fig. 1A-5](#)). The black shale and overlying limestone are generally the most widespread marine beds.

In many successions, coarsening upward, gray-shale overlies the open-marine limestone, ([Fig. 1A-6](#)). These deposits are likely of fluvial-deltaic origin. It is on this heterogeneous surface, following exposure during sea level regression and early lowstand, that the terrestrial part of the next cycle, the paleosol, is developed.

2.1.1. *The flat craton and water depth*

During the late Middle and Late Pennsylvanian, the Pangean interior was a low relief surface with a gradient possibly < 1 m/km, with a few areas, e.g. the Ozark Dome in Missouri, of tectonically created higher relief ([McKee and Crosby, 1975](#)). As a consequence of flat topography, small sea-level changes likely had large effects on coverage of the craton by marine waters. Low gradient and lack of significant topography would have allowed unimpeded airflow patterns over large areas. From a biological viewpoint, this regionally flat terrain suggests limited elevational effects on regional climate. Spatial disruption of lowstand ecosystems would have been widespread and extensive in response to drowning during times of rising sea level. Furthermore, due to covariation of climate and sea level with ice volume, ecosystems on the craton were affected by climate changes at the same time they were forced to move physical location by associated sea-level change.

2.1.2. *Terrestrial facies I: Paleosols*

Recent studies of coal underclays (e.g., [Cecil et al., 2003a](#); [Driese and Ober, 2005](#); [Rosenau et al., 2013](#)) indicate complex polygenetic histories, starting as mineral paleosols and transitioning to peat formation. In soil moisture terms ([Cecil, 2013](#), p. 23; [Soil Survey Staff, 1998](#)), as related to rainfall, these soils were first well drained, with Ustic soil moisture (seasonal dryness), induced by seasonal rainfall under subhumid climate. Increasing rainfall under moist subhumid to humid climate produced Udic soil moisture (evenly distributed annual rainfall equal to evapotranspiration). Finally, Aquic soil moisture (saturated) developed, rainfall exceeding evapotranspiration for most of the year (humid to perhumid climate). Paleosols also reflect the period-scale drying trend. Through Early and early Middle Pennsylvanian, most paleosols indicate initially Udic soil conditions, becoming Aquic as rainfall increased and seasonality decreased during the glacial

phase of a cycle (Driese and Ober, 2005). In late Middle and Late Pennsylvanian, however, paleosols with vertic features and calcium carbonate concentrations indicate initially Ustic soil moisture regimes (wet during the growing season but dry most of the year); these paleosols formed under seasonal, dry sub-humid to semi-arid climatic regimes. Carbonate concentrations reflect periods during which evapotranspiration strongly exceeded moisture throughput, resulting in pedogenic carbonate mineral deposition in the soil (Driese et al., 2005). In all of these polygenetic paleosols, later stages of development record translocation of clays, various minerals and iron downward through the soil, overprinting earlier soil profiles. This indicates increase in moisture throughput, reduction in seasonality and gradual but persistent increase in the consecutive number of months where rainfall exceeded evapotranspiration. Translocation of iron, in particular, known as gleying, indicates the existence of acidic (facilitating conversion of insoluble ferric to soluble ferrous iron), well drained (facilitating translocation of soluble iron) conditions. The acidity most likely results from organic matter buildup (a response to rainfall-induced increase in vegetative cover) in the soil O horizon, accompanied by formation of weak organic acids. As precipitation increased to humid or perhumid conditions, mineral substrates became permanently water logged, and clastic swamps, organic mucks, and shallow lakes developed on the mineral paleosol surface. In such wetland deposits characteristic plant fossils may be present. The deposits sometimes grade without break into coal beds, indicating a conversion of clastic swamps to peat swamps. Most often, however, the contact between the top of the paleosol and the bottom of the coal bed is relatively sharp. Variation in the paleosol-coal bed contact captures the meter-to-decimeter scale, topographic irregularity of the mineral-paleosol, pre-peat swamp landscape and indicates a rapid shift from mineral soil formation to widespread development of organic Histosols and peat formation (Fig. 1A-1).

Penecontemporaneous paleochannels, including extensive floodplains, are associated with the mineral paleosol interval. Rarely mapped in detail, a few examples (Allen et al., 2011; Bragonier et al., 2007; Feldman et al., 1995; Potter, 1963) indicate that floodplains were dynamic and that principle rivers sometimes deeply incised the craton.

2.1.3. Terrestrial facies II: Coal

Despite much study of coal, both as a rock and a former peat deposit, many misconceptions and misunderstandings remain. We address some of those here (Fig. 1A-2).

We differentiate peat from organic muck, and confine our discussion to those peats that could give rise to commercial grade coal beds. This means low mineral content of the original peat (< 20%), leading to low ash content (< 40%) of the resulting coal. Organic-rich deposits with mineral matter > 20% likely yielded black shales or organic-rich gray shales.

As a former peat deposit, coal beds are likely to have had highly complex histories of formation, including periods of drowning and decay, removal of peat by fires, and intervals of rapid organic accumulation (DiMichele and Phillips,

1994). In cases of repeated disruption of peat formation, multi-benched coals resulted. Cecil et al. (1985) summarized conditions of peat formation. For peat to form in warm equatorial regions precipitation must exceed evapotranspiration during most of the year or the water table will fall and organic destruction by fires and organism-mediated decay will occur. Equatorial rainfall must exceed evapotranspiration for > 10 months (humid to perhumid climates) for rates of organic accumulation to exceed destruction (Fig. 1F). Peat may form any place and any time these conditions are met, regardless of sea level, elevation, or some degree of topographic irregularity, including areas of free drainage, as evidenced by modern peat formation on windward slopes of the British Isles. Even under high, aseasonal rainfall, thickness and degree of decay of peat bodies will vary according to the local physical and chemical conditions (including no peat formation). Much evidence suggests that any given Pennsylvanian coal bed is time equivalent or nearly so throughout its extent: the peat formed as a blanket deposit over an entire landscape. Indications of this include such features as coal beds that extend across entire basins, regional-scale volcanic ash partings in a number of coal beds (Greb et al., 1999; Lyons et al., 1992; Opluštil et al., 2007; Wang et al., 2012), or widespread mineral partings that separate petrographically and palynologically distinctive coal benches (Eble et al., 2006; Greb et al., 2003). As a consequence, initiation and maintenance of peat-forming conditions call for a mechanism that can systematically raise water tables over a vast area of the craton effectively simultaneously. The most likely mechanism is humid to perhumid climate (see Cecil, 2003, p. 16), amplified by the exceedingly low relief and consequent poor drainage. Rivers draining Pennsylvanian peat swamps during the wettest phases of glacial-interglacial cycles appear to have carried low sediment loads and, at times, to have been “black water”, carrying most of their load as dissolved solids (Nelson et al., 2008). Such patterns reflect large reductions in sediment mobility caused by dense lowland vegetation cover. In the Pangean interior, this was combined with great distances to upland-region sediment sources. A modern parallel is the extremely low sediment load in Indonesian rivers under humid to perhumid climates, including those draining peat swamps (Cecil et al., 2003b). There, even in areas of steep elevational gradients, vegetation cover induced by high rainfall strongly limits erosion and sediment runoff into streams due to dense penetration and binding of the soil by plant roots (Cecil et al., 2003b).

2.1.4. Transitional terrestrial-to-marine facies

The most poorly studied lithofacies of a cyclothem is the Gray Siltstone Wedge (Fig. 1A-3). These deposits, of limited areal extent, back fill and occur lateral to peat-contemporaneous rivers that were converted to estuaries during sea-level transgression. Sediment comprising the GSWs may have been generated, at least in part, by tidal erosion during initial flooding of the craton. During these early phases of sea-level rise and onset of cratonic flooding, climate began a shift to increasingly seasonal, subhumid (Cecil et al., 2003a; Horton et al., 2012). Such climate change across the equatorial region reduced vegetation

density on the landscape, permitting significant increases in the sediment load of stream drainages (Cecil and Dulong, 2003). In combination with rising sea level, sediment was pushed inward at drowning river mouths, causing rapid aggradation, and the development of large mudflats flanking estuaries (Archer, 2004). Peat swamps were converted to short-lived clastic swamps. Evidence points to very high rates of sediment accumulation (Archer and Kvale, 1993; Archer et al., 2014), which account for the burial of upright trees and the abundant preservation of plant remains (e.g., DiMichele et al., 2009; King et al., 2011). Recent studies suggest that GSW deposits formed in a series of episodic meltwater pulses (Archer et al., 2014), during which sediment accumulation rates in and lateral to estuaries were rapid and high (Archer and Greb, 2012). Fossil evidence of in situ preserved plants indicates that the GSW was initially fresh-water. Salinities became brackish as sediment accumulated, indicated by invertebrates such as linguloids, myelinid pelecypods, and eurypterids. There is no evidence of normal marine salinities during GSW deposition.

2.1.5. Marine facies

Onset of marine conditions in a cyclothem is often marked by a widespread erosional or “ravinement” surface that scours the top of the coal bed and the GSW (Liu and Gastaldo, 1992). This can be deduced from the truncation of coal laminae and by erosional remnants of GSW sediments in areas more distal from channels (DeMaris et al., 1983). The ravinement surface is caused by tidal energy during the widespread flooding of epicontinental basins. Although a thin transgressive limestone, often a shell-hash lag, may be present immediately above the ravinement surface, more typically this surface is overlain directly by a fissile, marine black shale. Black shales had complex depositional and geochemical histories (Algeo et al., 2004; Schultz and Coveney, 1992) and anoxic to dysoxic conditions are indicated by high organic content and limited bioturbation. The physical conditions necessary to generate low oxygen bottom waters have been debated. Prevailing models estimate 70–100 m water depths (e.g., Heckel, 1977) and place these shales near marine high-stand (e.g. Algeo et al., 2004). Other studies (e.g., Cecil et al., 2003a; Coveney et al., 1991; Zangerl and Richardson, 1963), including those of Gondwanan ice volumes (Isbell et al., 2003b), suggest much shallower water. Low oxygen, shallow-depth bottom waters have been ascribed to limited mixing (Cecil et al., 2003a), reflecting low surface wind velocities and limited mixing during early transgression, when strong polar high-pressure fronts continued to restrict cross-equatorial migration of the Intertropical Convergence Zone (ITCZ), creating equatorial doldrums. Holterhoff and Cassady (2012) also ascribe black-shale deposition to climatic conditions, arguing that under wet, weakly seasonal climates, sea-bottom anoxia is created by a hyposaline cap from terrestrial runoff that causes nutrient loading of surface waters and high productivity. In all these models, despite slow, limited siliciclastic input, there is no carbonate production. Again, climate-driven scenarios offer solutions that water depth does not; the initiation

of carbonate formation begins with the onset of drier climatic conditions, and greater mixing of the water column caused by break down of the doldrums and increased surface winds (Cecil et al., 2003a) (Fig. 1A–4, 5, 6).

Equatorial climates became most strongly seasonal (dry sub-humid to semi-arid) during late transgression and into highstand. Modeling studies (Horton et al., 2012) indicate the highest absolute levels of annual rainfall during late transgression to early highstand phases of the cycle, but with strong seasonality. During this time period and as noted above, marine limestones became widespread across the craton. Subsequent aridity during late highstand and during regression reflects the largest seasonal excursions of the ITCZ during a glacial–interglacial cycle, a response to retreat of polar ice masses and associated Arctic high-pressure (Cecil et al., 2003a). This also can be visualized as a weakening of equatorial Hadley cell circulation (Peyser and Poulsen, 2008), resulting in increased seasonality and reduction of equatorial precipitation. Also, during interglacials more extensive seasonal migrations of the ITCZ create more intense equatorial winds, resulting in greater mixing and water column oxygenation, a suggested explanation for the shift from black-shale deposition to limestone formation (Cecil et al., 2003a).

Deltaic deposits of regional extent, emplaced during late highstand and early regression, often succeed open marine strata. With initiation of ice buildup, equatorial climates began to see increased precipitation, though of a seasonal nature, due to reductions in the amplitude of ITCZ annual migrations and general Hadley cell strengthening (Horton et al., 2012). The result was significant increases in siliciclastic sediment transport. The biota of these deposits varies widely from marine to brackish fauna; the plant component is largely wetland, reflecting the high water tables of the delta top, but rarely may reveal species typical of seasonally dry climates, reflecting the general background conditions (Pšenička et al., 2011).

It is partially upon these deltaic deposits that the terrestrial phase of the next glacial cycle began. As sea level receded (regression due to ice buildup), a terrestrial surface emerged across the basinal lowlands of the craton. Paleosols indicate that this initially occurred under a seasonally dry climate (Driese and Ober, 2005; Feldman et al., 2005; Joeckel, 1989; Rosenau et al., 2013), becoming more humid as ice volume increased and sea level fell.

2.2. A note on accommodation space and inter-basin correlations

There was great variation in the rates and amounts of accommodation created within different Pennsylvanian lowland basins. For example, the Lower and lower Middle Pennsylvanian of the Appalachian basin is nearly 1200 m thick, whereas it is only 200 m thick in the Illinois basin. Foreland basin subsidence may have been the dominant Appalachian basin tectonic control on accommodation space whereas sediment loading and other unknown tectonic forces controlled Eastern and Western Interior basin subsidence (McKee and Crosby, 1975). In contrast, in the Central Pangean Mountain regions of Central Europe,

there were highly variable rates of sediment accumulation, some basins being entirely intra-montane and potentially high altitude (Opluštil, 2005). Nonetheless, the effects of tectonics are sufficiently time averaged in large, paralic, equatorial basins to permit marine-organism-based correlations across wide areas for many time intervals of the Pennsylvanian; individual cycles and stacks of cycles have been correlated from the American Western Interior to as far east as the Donets basin of the Ukraine (Eros et al., 2012; Heckel et al., 2007; Rosscoe and Barrick, 2013), particularly in the late Middle and Late Pennsylvanian. Correlations among American basins based on marine organisms and coal palynology (Cecil et al., 2003a; Peppers, 1996) have made linkages between most of the major and many of the minor cycles.

3. Glacial–interglacial cyclicity and the equatorial terrestrial biota

Pennsylvanian glacial–interglacial cycles had major effects on both terrestrial and marine biotic dynamics in equatorial latitudes. Although we focus here on the terrestrial, there are parallels in the marine realm. Most importantly, there were environments that oscillated greatly in areal extent, particularly terrestrial peat swamps and marine black-mud ocean bottoms. For most of the time represented by a cyclothem, these occasionally widespread environmental-types either did not exist or were confined to small areas. At other times, and for relatively shorter periods, they were present over huge areas.

Is the Pleistocene to Recent ice age an analogue for the Pennsylvanian, and vice versa? Certainly, there are strong parallels in the fundamental allocyclicity of the system on large spatio-temporal scales. However, there are major differences in continental configuration, size and location of interior continental flatlands and positions of mountain ranges, which create distinct patterns of ocean and atmospheric circulation. In other words, glacial–interglacial cycling played out in different regions at these different times. Furthermore, and perhaps of even greater importance—flowering plants have dominated tropical ecosystems since the Late Cretaceous. Recent studies of Boyce and colleagues (Boyce et al., 2010) indicate angiosperms can pump water from the soil to the atmosphere via evapotranspiration at rates far exceeding any other plant group (though see Wilson et al., 2008, on medullosan water relations). Consequently, modern-type rainforests, where over 70% of rainfall is “recycled” via high altitude cooling and condensation of evapotranspirative moisture, may not have existed before the evolution of angiosperms. The implication is that the rainfall fueling Pennsylvanian peat swamps may have been generated largely by atmospheric convective precipitation, the plants contributing little to the creation of moisture clouds to support rainforest architectures. Thus, without flowering plants, interglacials of our modern terrestrial equatorial regions would be significantly more seasonally dry than at present.

One final caveat is needed when comparing and contrasting the Late Paleozoic and modern worlds—biodiversity. Late Paleozoic terrestrial ecosystems had many fewer species than those younger than Early

Cretaceous (e.g., Wing and DiMichele, 1995). In contrast, the number of dominant major phylogenetic groups (traditional Linnean classes) in the landscape was much greater than today (DiMichele and Phillips, 1996). The consequences of this are several and, as yet, only peripherally explored by paleobotanists as follows:

- strong niche conservatism. Resource partitioning appears to track phylogenetic relatedness—specific evolutionary lineages had distinct ecologies within terrestrial landscapes. This phenomenon has been labeled “phylogenetic niche partitioning” (e.g., Webb et al., 2002) and is present as far back into the land plant record as anyone has looked for it (Hotton et al., 2001);
- niche breadth of late Paleozoic species may have been greater than that found in most modern species, on average. There were fewer players in the game, but that game–resource competition and niche partitioning—was played by the same rules as today. At present, this is virtually unexplored;
- most kinds of plants relied on wind for some phase of their life cycle: pollination, seed dispersal or spore dispersal. A few relied on water for dispersal. Reliance on wind, or on water at certain life-cycle phases, created the potential for extensive gene flow. High gene flow can be expected to reduce rates of speciation and, thereby, rates of radical morphological or physiological innovation. As far as we know, only one study has addressed this (Raymond and Costanza, 2007), by examining rates of species turnover in pteridosperms (non-wind dispersed large seeds and pollen) vs. other Pennsylvanian seed-plant groups (wind dispersed);
- plant–animal interactions were much more limited in scope than in post-Paleozoic systems (Labandeira and Sepkoski, 1993). In the Paleozoic there were less herbivory and fewer kinds of herbivory, limiting the transfer of plant primary productivity to consumer portions of food webs. In addition, there is no definitive evidence for pollination of plants by animal vectors;
- Pennsylvanian equatorial terrestrial ecosystems were broadly of two types: those composed of wetland plants and those composed of plants adapted to seasonal drought (there probably were several kinds of these depending on the degree of moisture stress) (DiMichele et al., 2008). These can be considered unique biomes. There is little evidence to suggest large populations of species adapted to seasonal drought within areas of widespread wetlands; such occurrences as are known appear to represent allochthonous elements introduced by long-distance transport. In contrast, wetland elements are a nearly ubiquitous autochthonous to parautochthonous background in seasonally dry landscapes (e.g., Bashforth et al., 2014). This asymmetry is expected. Widespread wetlands call for great areal coverage of the continental interior by humid to perhumid climates. Under such conditions, drainage effects caused by local topographic irregularity are greatly suppressed, leaving few places where plants requiring seasonal drought might flourish. In contrast, seasonally dry landscapes, particularly those of sub-humid climates, are sprinkled with local areas of high water table, such as stream

margins, intra-channel bars, lakesides, and coastal lagoons, providing patches and corridors of wetlands for obligately wetland species.

There are two major issues to examine in the terrestrial equatorial regions when considering the biological consequences of the strongly linked oscillations in climate and sea level that accompanied glacial–interglacial cycles:

- what happened to the wetland species, and the ecosystems in which they thrived, during periods of widespread seasonal aridity in the equatorial regions?
- Conversely, where were seasonally dry taxa during periods of widespread wetland development? Empirical data suggest that these two kinds of ecosystem did not simply oscillate back and forth in space through time as climate changed. Rather, the two vegetation types and their component species pools had different dynamics between intervals of abundance in the lowlands (DiMichele et al., 2010).

3.1. Wetland Fate

Empirical evidence (e.g., Bashforth et al., 2014; Dolby et al., 2011) and climate models (Horton et al., 2012; Poulsen et al., 2007) suggest absence during interglacial periods of equatorial areas sufficiently wet to support large tracts of wetland-specialist vegetation. Consequently, it is most likely that populations of wetland plants broke up and survived in refugial areas: stream- and lake-sides, in braidplains, along coastlines (Falcon-Lang and DiMichele, 2010). Localized occurrences of wetland species within dryland assemblages, and occurrence of wetland palynomorphs within strata deposited during interglacial intervals (Falcon-Lang et al., 2009; Hawkins et al., 2013; Stephenson et al., 2008), indicate the presence of wetland-plant populations in seasonally dry landscapes. However, the likelihood of organic-matter preservation is greatly reduced in seasonally dry settings (Gastaldo and Demko, 2011) by:

- low probability of short-term burial below mean low water table accompanied by high rates of degradation;
- low likelihood of subsequent intermediate term burial by marine water (if such landscapes occur between highstand and early lowstand, during regression).

Based on these boundary conditions, wetland vegetation would have reassembled from refugia during each recurrence of widespread, equatorial humid climate. Despite these regular oscillations there is high compositional and dominance-diversity conservatism of wetland vegetation (e.g., DiMichele et al., 2009). The cause of this conservatism may be the dominance of wetland disassembly-reassembly cycles by lottery dynamics:

- during contraction into refugia, the dominant species of the preceding widespread wet period were most likely to dominate in refugia;
- these most numerically abundant species then reasserted proportional dominance as populations expanded

out of refugia during the next wet interval. The main limiting variable was dispersal capacity.

Because most wetland plants had wind-dispersed pollen or microspores (Phillips et al., 1985; Taylor and Millay, 1979), the possibility was high for maintaining genetic continuity among populations physically fragmented by widespread seasonal drought. The great exception to this generality is the medullosan pteridosperms, which had poorer pollen dispersal than other seed plants, due to the large size of the grains; consequently, they should have had more population isolation and higher rates of speciation (Raymond and Costanza, 2007). Nearly all the dominants have propagules that were either water (lycopsids–e.g., Phillips, 1979; pteridosperms–e.g., DiMichele et al., 2006) or wind (tree ferns–e.g., Willard et al., 2007; cordaitaleans–e.g., Souza and Iannuzzi, 2012) dispersed. Thus, no group in particular may have had a signal advantage during the onset of renewed wet climates, which would favor reassembly controlled largely by initial numbers. However, because of the periodic oscillations in population sizes, the entire system had a built-in hypersensitivity to extreme interglacial aridity, during which populations could be squeezed to extinction or regional extirpation. This apparently did cause occasional widespread ecosystem reorganization at environmental threshold points (DiMichele et al., 2010; Heckel, 1991; Phillips et al., 1974).

3.2. Seasonally dry vegetational dynamics

There appear to have been permanent, large populations of seasonally-dry-adapted plants in western equatorial Pangea and parts of the Central Pangean Mountains (e.g., Bashforth et al., 2014; Cridland and Morris, 1963; Falcon-Lang and Bashforth, 2005; Tidwell et al., 1992; Uhl et al., 2004). As a result, it would seem that these low diversity, nearly entirely wind pollinated and dispersed, largely seed-plant floras simply spread across the land surface in response to climate changes, following topographically constrained routes. They exhibit, as a consequence, long-term compositional homogeneity. And the change from cordaitalean to coniferalean dominated assemblages is long and drawn out over many millions of years, tracking a long-term trend in increasing equatorial aridity at all phases of glacial–interglacial cycles (e.g., Tabor and Poulsen, 2008).

As a consequence of these large standing populations, seasonally-dry-adapted vegetation appears to have experienced limited evolutionary innovation within any given species pool. Rather, innovations appear to have originated from the progressive colonization of environments with increasingly greater moisture and, possibly/probably, temperature extremes, and other environmental extremes, such as salinity (e.g., Poort and Kerp, 1990). This implies that survival of new body plans, physiologies, and reproductive strategies was most likely to occur where competition from large, standing populations of ancestral forms was lowest, at the ragged, gradational margins of environmental tolerance of the species (DiMichele and Aronson, 1992). It is in the deposits formed in these

progressively more extreme environments that progressively more phylogenetically advanced elements of respective clades are found (e.g., conifers–Kerp, 1996; Looy, 2007; gigantopterids–DiMichele et al., 2011; peltasperms–Poort and Kerp, 1990; Naugolnykh and Kerp, 1996; cycadophytes–DiMichele et al., 2001; corystosperms–Kerp et al., 2006).

Because of the poor preservation of seasonally dry floras during Pennsylvanian, and even Permian time, it is challenging to assemble a coherent picture of their evolutionary history. However, by documenting their biogeographic occurrences and linking them to indicators of climate, it is possible to develop an idea of the environmental factors affecting their evolutionary dynamics. These appear to have been different from those affecting the plants of the wetland biome and reflect mainly dry-season length or confinement to special edaphic conditions.

4. Conclusions

The Pennsylvanian and Early Permian glacial world has many parallels to the Pleistocene, Holocene and Recent. Cyclic waxing and waning of glaciers, and periodic intervals of warming with near disappearance of polar ice, created a dynamic world–geologically and biologically. The equatorial regions of this time hosted the well-known peat (coal) swamps. They also, however, were home to vast tracts of plants adapted to seasonal drought. Understanding the relationships of these distinct biomes relies upon an understanding of the climatic dynamics associated with glacial–interglacial cycles.

The evidence presented herein suggests strong linkages between ice volume, sea level, climate and patterns of lithological variation in the sedimentary rock record. This system is represented in the interior of Pangea by the repetitious spatio-temporal stratigraphic packages of so-called “cyclothems”. Models that link ice-volume, atmospheric circulation and sea-level change, placing maximum rainfall and maximum wetland development at lowstand, best explain the data. Minimum seasonality occurred during glacial maximum and at sea-level lowstand. During interglacials through early parts of succeeding glacial phases, seasonally dry climates prevailed and the exposed parts of basinal lowlands were colonized by seasonally dry vegetation. A full cycle may have represented about 100,000 years.

Vegetation and, most likely, associated terrestrial vertebrate and invertebrate faunas, closely followed the changes in climate state across the Pangean equatorial interior. These distinct biomes shared few taxa in common and had distinct ecological and evolutionary dynamics.

Acknowledgements

We thank Sylvie Bourquin for the invitation to submit this paper. Research supported by U.S. Geological Survey, Smithsonian Institution, and State of Illinois. We thank many colleagues for sharing their ideas and expertise.

References

- Algeo, T.J., Schwark, L., Hower, J.C., 2004. High-resolution geochemistry and sequence stratigraphy of the Hushpuckney Shale (Swope Formation, eastern Kansas): implications for climato-environmental dynamics of the Late Pennsylvanian Midcontinent Seaway. *Chem. Geol.* 206, 259–288.
- Allen, J.P., Fielding, C.R., Gibling, M.R., Rygel, M.C., 2011. Fluvial response to paleo-equatorial climate fluctuations during the Late Paleozoic ice age. *Geol. Soc. Amer. Bull.* 123, 1524–1538.
- Archer, A.W., 2004. Recurring assemblages of biogenic and physical sedimentary structures in modern and ancient extreme macrotidal estuaries. *J. Coastal Res.* 43, 4–22.
- Archer, A.W., Greb, S.F., 2012. Hypertidal facies from the Pennsylvanian Period: Eastern and Western Interior Coal Basins, USA. In: Davis, Jr., R.A., Dalrymple, R.W. (Eds.), *Principles of Tidal Sedimentology*. Springer, Netherlands, pp. 421–436.
- Archer, A.W., Kvale, E.P., 1993. Origin of gray-shale lithofacies (“clastic wedges”) in US midcontinental coal measures (Pennsylvanian): an alternative explanation. *Geological Society of America Special Papers* 286, 181–192.
- Archer, A.W., Feldman, H.R., Kvale, E.P., Lanier, W.P., 1994a. Comparison of drier to wetter-interval estuarine roof facies in the Eastern and Western Interior coal basins, USA. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 106, 171–185.
- Archer, A.W., Lanier, W.P., Feldman, H.R., 1994b. Stratigraphy and depositional history within incised-paleovalley fills and related facies, Douglas Group (Missourian–Virgilian; Upper Carboniferous) of Kansas, USA. *SEPM Special Publication* 51, 175–190.
- Archer, A.W., Elrick, S.D., Nelson, W.J., DiMichele, W.A., 2014. Catastrophic burial of Pennsylvanian Period coal swamps in the Illinois Basin: Hypertidal sedimentation during Gondwanan glacial melt-water pulses (in press).
- Bashforth, A.R., Cleal, C.J., Gibling, M.R., Falcon-Lang, H.J., Miller, R.F., 2014. Paleoeecology of Early Pennsylvanian vegetation on a seasonally dry tropical landscape (Tynemouth Creek Formation, New Brunswick, Canada). *Rev. Palaeobot. Palynol.* 200, 229–263.
- Birgenheier, L.P., Fielding, C.R., Rygel, M.C., Frank, T.D., Roberts, J., 2009. Evidence for dynamic climate change on sub-10⁶-year scales from the late Paleozoic glacial record, Tamworth Belt, New South Wales, Australia. *J. Sediment. Res.* 79, 56–82.
- Boyce, C.K., Lee, J.E., Feild, T.S., Brodribb, T.J., Zwieniecki, M.A., 2010. Angiosperms helped put the rain in the rainforests: The impact of plant physiological evolution on tropical biodiversity. *Ann. Mo. Bot. Garden* 97, 527–540.
- Bragonier, W.A., Bohan, M.P., Hawk, L.R., 2007. The monster channel complex of central-western Pennsylvania. *Pennsylvania Geology* 37, 2–13.
- Bristol, H.M., Howard, R.H., 1974. Sub-Pennsylvanian valleys in the Chesterian surface of the Illinois Basin and related Chesterian slump blocks. *Geological Society of America Special Paper* 148, 315–335.
- Cecil, C.B., 1990. Paleoclimate controls on stratigraphic repetition of chemical and siliciclastic rocks. *Geology* 18, 533–536.
- Cecil, C.B., 2003. The concept of autocyclic and allocyclic controls on sedimentation and stratigraphy, emphasizing the climatic variable. *SEPM Special Publication* 77, 13–20.
- Cecil, C.B., 2013. An overview and interpretation of autocyclic and allocyclic processes and the accumulation of strata during the Pennsylvanian–Permian transition in the central Appalachian Basin, USA. *Int. J. Coal Geology* 119, 21–31.
- Cecil, C.B., Dulong, F.T., 2003. Precipitation models for sediment supply in warm climates. *SEPM Special Publication* 77, 21–28.
- Cecil, C.B., Stanton, R.W., Neuzil, S.G., Dulong, F.T., Ruppert, L.F., Pierce, B.S., 1985. Paleoclimate controls on Late Paleozoic sedimentation and peat formation in the central Appalachian Basin (USA). *Int. J. Coal Geology* 5, 195–230.
- Cecil, C.B., Dulong, F.T., West, R.R., Stamm, R., Wardlaw, B.A., Edgar, N.T., 2003a. Climate controls on the stratigraphy of a Middle Pennsylvanian cyclothem in North America. *SEPM Special Publication* 77, 151–182.
- Cecil, C.B., Dulong, F.T., Harris, R.A., Cobb, J.C., Gluskoter, H.G., Hendro Nugroho, 2003b. Observations on Climate and Sediment Discharge in Selected Tropical Rivers. Indonesia *SEPM Special Publication* 77, 29–50.
- Coveney Jr., R.M., Watney, W.L., Maples, C.G., 1991. Contrasting depositional models for Pennsylvanian black shale discerned from molybdenum abundances. *Geology* 19, 147–150.
- Cridland, A.A., Morris, J.E., 1963. *Taeniopteris Walchia and Dichophyllum in the Pennsylvanian System of Kansas*. Univ. Kansas Sci. Bull. 44, 71–85.

- Davies, N.S., Gibling, M.R., 2013. The sedimentary record of Carboniferous rivers: Continuing influence of land plant evolution on alluvial processes and Palaeozoic ecosystems. *Earth Sci. Rev.* 120, 40–79.
- DeMaris, P.J., Bauer, R.A., Cahill, R.A., Damberger, H.H., 1983. Geologic investigation of roof and floor strata: longwall demonstration, Old Ben Mine No. 24. Prediction of coal balls in the Herrin Coal. Final technical report: part 2. Illinois State Geological Survey Contract-Grant Report 1983-2.
- DiMichele, W.A., Aronson, R.B., 1992. The Pennsylvanian–Permian vegetational transition: a terrestrial analogue to the onshore-offshore hypothesis. *Evolution* 46, 807–824.
- DiMichele, W.A., Phillips, T.L., 1994. Paleobotanical and paleoecological constraints on models of peat formation in the Late Carboniferous of Euramerica. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 106, 39–90.
- DiMichele, W.A., Phillips, T.L., 1996. Clades, ecological amplitudes, and ecomorphs: phylogenetic effects and persistence of primitive plant communities in the Pennsylvanian-age tropical wetlands. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 127, 83–105.
- DiMichele, W.A., Mamay, S.H., Chaney, D.S., Hook, R.W., Nelson, W.J., 2001. An Early Permian flora with Late Permian and Mesozoic affinities from north-central Texas. *J. Paleontology* 75, 449–460.
- DiMichele, W.A., Phillips, T.L., Pfefferkorn, H.W., 2006. Paleoecology of Late Paleozoic pteridosperms from tropical Euramerica. *J. Torrey Bot. Soc.* 133, 83–118.
- DiMichele, W.A., Kerp, H., Tabor, N.J., Looy, C.V., 2008. The so-called “Paleophytic–Mesophytic” transition in equatorial Pangea—Multiple biomes and vegetational tracking of climate change through geological time. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 268, 152–163.
- DiMichele, W.A., Montañez, I.P., Poulsen, C.J., Tabor, N.J., 2009. Climate and vegetational regime shifts in the Late Paleozoic ice age earth. *Geobiology* 7, 200–226.
- DiMichele, W.A., Cecil, C.B., Montañez, I.P., Falcon-Lang, H.J., 2010. Cyclic changes in Pennsylvanian paleoclimate and effects on floristic dynamics in tropical Pangaea. *Int. J. Coal Geology* 83, 329–344.
- DiMichele, W.A., Looy, C.V., Chaney, D.S., 2011. A new genus of gigantopterid from the Middle Permian of the United States and China and its relevance to the gigantopterid concept. *Int. J. Plant Sci.* 172, 107–119.
- Dolby, G., Falcon-Lang, H., Gibling, M.R., 2011. A conifer-dominated palynological assemblage from Pennsylvanian (Late Moscovian) alluvial drylands in Atlantic Canada: implications for the vegetation of tropical lowlands during glacial phases. *J. Geol. Soc.* 168, 571–584.
- Driese, S.G., Ober, E.G., 2005. Paleopedologic and paleohydrologic records of precipitation seasonality from Early Pennsylvanian “underclay” paleosols, USA. *J. Sediment. Res.* 75, 997–1010.
- Driese, S.G., Nordt, L.C., Lynn, W.C., Stiles, C.A., Mora, C.I., Wilding, L.P., 2005. Distinguishing climate in the soil record using chemical trends in a Vertisol climosequence from the Texas Coast Prairie, and application to interpreting Paleozoic paleosols in the Appalachian Basin, USA. *J. Sediment. Res.* 75, 339–349.
- Eble, C.F., Grady, W.C., Pierce, B.S., 2006. Compositional characteristics and inferred origin of three Late Pennsylvanian coal beds from the northern Appalachian Basin. *Geological Society of America Special Paper* 399, 197–222.
- Elrick, M., Scott, L.A., 2010. Carbon and oxygen isotope evidence for high-frequency (10^4 – 10^5 yr) and My-scale glacio-eustasy in Middle Pennsylvanian cyclic carbonates (Gray Mesa Formation), central New Mexico. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 285, 307–320.
- Elrick, S.D., Nelson, W.J., 2010. Facies relationships of the Middle Pennsylvanian Springfield coal and Dykersburg shale: constraints on sedimentation, development of coal splits and climate change during transgression. *Geological Society of America Abstracts with Programs* 42, 51.
- Eros, J.M., Montañez, I.P., Osleger, D.A., Davydov, V.I., Nemyrovska, T.I., Poletaev, V.I., Zhykalyak, M.V., 2012. Sequence stratigraphy and onlap history of the Donets Basin, Ukraine: insight into Carboniferous icehouse dynamics. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 313–314, 1–25.
- Falcon-Lang, H.J., Bashforth, A.R., 2005. Morphology, anatomy, and upland ecology of large cordaitalean trees from the Middle Pennsylvanian of Newfoundland. *Rev. Palaeobotan. Palynol.* 135, 223–243.
- Falcon-Lang, H.J., DiMichele, W.A., 2010. What happened to the coal forests during Pennsylvanian glacial phases? *Palaios* 25, 611–617.
- Falcon-Lang, H.J., Nelson, W.J., Elrick, S., Looy, C.V., Ames, P.R., DiMichele, W.A., 2009. Incised channel fills containing conifers indicate that seasonally dry vegetation dominated Pennsylvanian tropical lowlands. *Geology* 37, 923–926.
- Feldman, H.R., Gibling, M.R., Archer, A.W., Wightman, W.G., Lanier, W.P., 1995. Stratigraphic architecture of the Tonganoxie paleovalley fill (Lower Virgilian) in northeastern Kansas. *AAPG Bull.* 79, 1019–1043.
- Feldman, H.R., Franseen, E.K., Joeckel, R.M., Heckel, P.H., 2005. Impact of longer-term modest climate shifts on architecture of high-frequency sequences (cyclothem), Pennsylvanian of Midcontinent USA. *J. Sediment. Res.* 75, 350–368.
- Fielding, C.R., Frank, T.D., Isbell, J.L., 2008. The Late Paleozoic ice age—A review of current understanding and synthesis of global climate patterns. *Geological Society of America Special Paper* 441, 343–354.
- Gastaldo, R.A., Demko, T.M., 2011. The relationship between continental landscape evolution and the plant–fossil record: long term hydrologic controls on preservation. In: Allison, P.A., Bottjer, D.J. (Eds.), *Taphonomy: process and Bias Through Time*. Topics in Geobiology, 32, pp. 249–285.
- Greb, S.F., Eble, C.F., Hower, J.C., 1999. Depositional history of the Fire Clay coal bed (Late Duckmantian), eastern Kentucky, USA. *Int. J. Coal Geology* 40, 255–280.
- Greb, S.P., Andrews, W.M., Eble, C.F., DiMichele, W., Cecil, C.B., Hower, J.C., 2003. Desmoinesian coal beds of the Eastern Interior and surrounding basins: the largest tropical peat mires in Earth history. *Geological Society of America Special Paper* 370, 127–150.
- Hawkins, K., Davies, S.J., Mullins, G.L., Macquaker, J.H.S., 2013. Miospore distribution and sedimentological facies distribution as an insight to changing terrestrial palaeoequatorial floral communities during a Pennsylvanian glacio-eustatic sea level cycle. *Rev. Palaeobotan. Palynol.* 197, 166–178.
- Heckel, P.H., 1977. Origin of phosphatic black shale facies in Pennsylvanian cyclothem of mid-continent North America. *AAPG Bull.* 61, 1045–1068.
- Heckel, P.H., 1990. Evidence for global (glacial-eustatic) control over Upper Carboniferous (Pennsylvanian) cyclothem in midcontinent North America. *Geological Society, London. Special Publications* 55, 35–47.
- Heckel, P.H., 1991. Lost Branch Formation and revision of Upper Desmoinesian stratigraphy along midcontinent Pennsylvanian outcrop belt. *Kansas Geol. Surv. Geology Series* 4, 1–67.
- Heckel, P.H., 2008. Pennsylvanian cyclothem in Midcontinent North America as far-field effects of waxing and waning of Gondwana ice sheets. *Geological Society of America Special Paper* 441, 275–289.
- Heckel, P.H., Alekseev, A.S., Barrick, J.E., Boardman, D.R., Goreva, N.V., Nemyrovska, T.I., Ueno, K., Villa, E., Work, D.M., 2007. Cyclothem [“digital”] correlation and biostratigraphy across the global Moscovian–Kasimovian–Gzhelian stage boundary interval (Middle–Upper Pennsylvanian) in North America and eastern Europe. *Geology* 35, 607–610.
- Holterhoff, P., Cassady, K., 2012. Late Pennsylvanian and Early Permian black shale–limestone bed-set couplets of the Eastern Shelf, Midland Basin (Texas): climate-driven redox cycles of the inner platform realm. *American Association of Petroleum Geologists Annual Meeting, Search and Discovery Article #90142*.
- Horton, D.E., Poulsen, C.J., Montañez, I.P., DiMichele, W.A., 2012. Eccentricity-paced late Paleozoic climate change. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 331–332, 150–161.
- Hotton, C.L., Hueber, F.M., Griffing, D.H., Bridge, J.S., 2001. Early terrestrial plant environments: an example from the Emsian of Gaspé, Canada. In: Edwards, D., Gensel, P.G. (Eds.), *Plants Invade the Land*. pp. 179–203.
- Isbell, J.L., Lenaker, P.A., Askin, R.A., Miller, M.F., Babcock, L.E., 2003a. Reevaluation of the timing and extent of Late Paleozoic glaciation in Gondwana: role of the Transantarctic Mountains. *Geology* 31, 977–980.
- Isbell, J.L., Miller, M.F., Wolfe, K.L., Lenaker, P.A., 2003. Timing of Late Paleozoic glaciation in Gondwana: Was glaciation responsible for the development of Northern Hemisphere cyclothem? *Geological Society of America Special Paper* 370, 5–24.
- James, G.W., Baker, D.R., 1972. Geochemistry of a Pennsylvanian black shale (Excello) in the Midcontinent and the Illinois Basin. *Kansas Geol. Surv. Bull.* 204, 3–10.
- Joeckel, R.M., 1989. Geomorphology of a Pennsylvanian land surface; pedogenesis in the Rock Lake Shale Member, southeastern Nebraska. *J. Sediment. Res.* 59, 469–481.
- Joeckel, R.M., 1999. Paleosol in Galesburg Formation (Kansas City Group, Upper Pennsylvanian), northern Midcontinent, USA; evidence for climate change and mechanisms of marine transgression. *J. Sediment. Res.* 69, 720–737.
- Kerp, H., 1996. Post-Variscan Late Paleozoic Northern Hemisphere gymnosperms: the onset to the Mesozoic. *Rev. Palaeobotan. Palynol.* 90, 263–285.
- Kerp, H., Hamad, A.A., Vörding, B., Bandel, K., 2006. Typical Triassic Gondwanan floral elements in the Upper Permian of the paleotropics. *Geology* 34, 265–268.

- King, S.C., Cleal, C.J., Hilton, J., 2011. Common ground between two British Pennsylvanian wetland floras: using large, first-hand datasets to assess utility of historical museum collections. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 308, 405–417.
- Labandeira, C.C., Sepkoski, J.J., 1993. Insect diversity in the fossil record. *Science* 261, 310–315.
- Langenheim Jr., R.L., Nelson, W.J., 1992. Chapter 6: the cyclothemic concept in the Illinois Basin: a review. *Geol. Soc. Amer. Mem.* 180, 55–72.
- Liu, Y., Gastaldo, R.A., 1992. Characteristics of a Pennsylvanian ravine-ment surface. *Sediment. Geol.* 77, 197–213.
- Looy, C.V., 2007. Extending the range of derived Late Paleozoic conifers: *Lebowskia* gen. nov. (Majonicaceae). *Int. J. Plant Sci.* 168, 957–972.
- Lyons, P.C., Outerbridge, W.F., Triplehorn, D.M., Evans, H.T., Congdon, R.D., Caprio, M., Hess, J.C., Nash, W.P., 1992. An Appalachian isochron: a kaolinized Carboniferous air-fall volcanic-ash deposit (tonstein). *Geol. Soc. Amer. Bull.* 104, 1515–1527.
- McKee, E.D., Crosby, E.J., 1975. Paleotectonic investigations of the Pennsylvanian System in the United States. *US Geol. Surv. Professional Paper* 853.
- Montañez, I.P., Poulsen, C.J., 2013. The Late Paleozoic Ice Age: an evolving paradigm. *Ann. Rev. Earth Planetary Sci.* 41, 629–656.
- Naugolnykh, S.V., Kerp, H., 1996. Aspects of Permian palaeobotany and palynology. XV. On the oldest known peltasperms with radially symmetrical ovuliferous discs from the Kungurian (uppermost Lower Permian) of the Fore-Urals (Russia). *Rev. Palaeobotan. Palynol.* 91, 35–62.
- Nelson, W.J., Elrick, S.D., DiMichele, W.A., 2008. Evolution of a peat-contemporaneous channel: Galatia channel, Pennsylvanian of Illinois basin. *Geological Society of America, Abstracts with Programs* 40 (5), 81.
- Opluštil, S., 2005. Evolution of the Middle Westphalian river valley drainage system in central Bohemia (Czech Republic) and its palaeogeographic implication. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 222, 223–258.
- Opluštil, S., Pšenička, J., Libertín, M., Šimůnek, Z., 2007. Vegetation patterns of Westphalian and Lower Stephanian mire assemblages preserved in tuff beds of the continental basins of Czech Republic. *Rev. Palaeobotan. Palynol.* 143, 107–154.
- Opluštil, S., Edress, N.A., Sýkorová, I., 2013. Climatic vs. tectonic controls on peat accretion in non-marine setting; an example from the Zláčč Formation (Yeadonian-Bolsovian) in the Intra-Sudetic Basin (Czech Republic). *Int. J. Coal Geology* 116–117, 135–157.
- Peppers, R.A., 1996. Palynological correlation of major Pennsylvanian (Middle and Upper Carboniferous) chronostratigraphic boundaries in the Illinois and other coal basins. *Geol. Soc. Amer. Mem.* 188, 1–119.
- Peysner, C.E., Poulsen, C.J., 2008. Controls on Permo-Carboniferous precipitation over tropical Pangaea: A GCM sensitivity study. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 268, 181–192.
- Phillips, T.L., Peppers, R.A., Avcin, M.J., Laughnan, P.F., 1974. Fossil plants and coal: patterns of change in Pennsylvanian coal swamps of the Illinois Basin. *Science* 184, 1367–1369.
- Phillips, T.L., 1979. Reproduction of heterosporous arborescent lycopods in the Mississippian–Pennsylvanian of Euramerica. *Rev. Palaeobotan. Palynol.* 27, 239–289.
- Phillips, T.L., Peppers, R.A., DiMichele, W.A., 1985. Stratigraphic and interregional changes in Pennsylvanian coal-swamp vegetation: environmental inferences. *Int. J. Coal Geology* 5, 43–109.
- Poort, R.J., Kerp, J.H.F., 1990. Aspects of Permian palaeobotany and palynology. XI. On the recognition of true peltasperms in the Upper Permian of western and central Europe and a reclassification of species formerly included in *Peltaspermum* Harris. *Rev. Palaeobotan. Palynol.* 63, 197–225.
- Potter, P.E., 1963. Paleozoic sandstones of the Illinois Basin. *Illinois State Geological Survey. Report of Investigations* 217, 1–92.
- Poulsen, C.J., Pollard, D., Montañez, I.P., Rowley, D., 2007. Late Paleozoic tropical climate response to Gondwanan deglaciation. *Geology* 35, 771–774.
- Pšenička, J., Kerp, H., Opluštil, S., Elrick, S., DiMichele, W.A., Bek, J., Nelson, W.J., Ames, P.R., 2011. Preliminary report on the earliest callipterid assignable to the morphogenus *Rhachiphyllum* Kerp from the Late Moscovian (Asturian) Farmington Shale (Illinois Basin, USA). XVII International Congress on the Carboniferous and Permian, Perth, Programme and Abstracts, 105.
- Raymond, A., Costanza, S., 2007. Are wind-pollinated less diverse and longer ranging than insect-pollinated groups? *Geological Society of America Abstracts with Programs* 39 (6), 565.
- Rosenau, N.A., Tabor, N.J., Elrick, S.D., Nelson, W.J., 2013. Polygenetic history of paleosols In Middle–Upper Pennsylvanian cyclothems of the Illinois Basin, USA: Part II. Integrating geomorphology, climate, and glacioeustasy. *J. Sediment. Res.* 83, 637–668.
- Roscoe, S.J., Barrick, J.E., 2013. North American species of the conodont genus *Idiognathodus* from the Moscovian–Kasimovian boundary composite sequence and correlation of the Moscovian–Kasimovian stage boundary. *New Mexico Museum of Natural History and Science Bulletin* 60, 354–371.
- Rygel, M.C., Fielding, C.R., Frank, T.D., Birgenheier, L.P., 2008. The magnitude of Late Paleozoic glacioeustatic fluctuations: a synthesis. *J. Sediment. Res.* 78, 500–511.
- Schultz, R.B., Coveney, R.M., 1992. Time-dependent changes for Midcontinent Pennsylvanian black shales. *U.S.A. Chem. Geol.* 99, 83–100.
- Soil Survey Staff, 1998. *Keys to Soil Taxonomy*. United States Department of Agriculture. Natural Resources Conservation Service Monograph 19, 1–326.
- Souza, J.M., Iannuzzi, R., 2012. Dispersal Syndromes of fossil Seeds from the Lower Permian of Paraná Basin, Rio Grande do Sul, Brazil. *Anais da Academia Brasileira de Ciências* 84, 43–68.
- Stephenson, M.H., Millward, D., Leng, M.J., Vane, C.H., 2008. Palaeoecological and possible evolutionary effects of Early Namurian (Serpukhonian, Carboniferous) glacioeustatic cyclicity. *J. Geol. Soc.* 165, 993–1005.
- Tabor, N.J., Poulsen, C.J., 2008. Palaeoclimate across the Late Pennsylvanian–Early Permian tropical palaeolatitudes: a review of climate indicators, their distribution, and relation to palaeophysiological climate factors. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 268, 293–310.
- Taylor, T.N., Millay, M.A., 1979. Pollination biology and reproduction in early seed plants. *Rev. Palaeobotan. Palynol.* 27, 329–355.
- Tidwell, W.D., Jennings, J.R., Beus, S.S., 1992. A Carboniferous flora from the Surprise Canyon Formation in the Grand Canyon, Arizona. *J. Paleontology* 66, 1013–1021.
- Uhl, D., Lausberg, S., Noll, R., Stapf, K.R.G., 2004. Wildfires in the Late Paleozoic of central Europe – an overview of the Rotliegend (Upper Carboniferous–Lower Permian) of the Saar–Nahe Basin (SW-Germany). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 207, 23–35.
- Wang, J., Pfefferkorn, H.W., Zhang, Y., Feng, Z., 2012. Permian vegetational Pompeii from Inner Mongolia and its implications for landscape paleoecology and paleobiogeography of Cathaysia. *Proceedings of the National Academy of Sciences* 109, 4927–4932.
- Watney, W.L., Wong, J.C., French Jr., J.A., 1989. Computer simulation of Upper Pennsylvanian (Missourian) carbonate-dominated cycles in western Kansas. *Kansas Geol. Surv. Bull.* 233, 415–430.
- Webb, C.O., Ackerly, D.D., McPeck, M.A., Donoghue, M.J., 2002. Phylogenies and community ecology. *Ann. Rev. Ecol. Syst.* 33, 475–505.
- Weller, J.M., 1931. The conception of cyclical sedimentation during the Pennsylvanian Period. *Illinois State Geol. Surv. Bull.* 60, 163–177.
- Willard, D.A., Phillips, T.L., Lesnikowska, A.D., DiMichele, W.A., 2007. Paleoecology of the Late Pennsylvanian-age Calhoun coal bed and implications for long-term dynamics of wetland ecosystems. *Int. J. Coal Geology* 69, 21–54.
- Wilson, J.P., Knoll, A.H., Holbrook, N.M., Marshall, C.R., 2008. Modeling fluid flow in *Medullosa*, an anatomically unusual Carboniferous seed plant. *Paleobiology* 34, 472–493.
- Wing, S.L., DiMichele, W.A., 1995. Conflict between local and global changes in plant diversity through geological time. *Palaios* 10, 551–564.
- Zangerl, R., Richardson, E.S., 1963. The paleoecological history of two Pennsylvanian black shales. *Fieldiana Geology Memoirs* 4, 1–352.