



Hydrology, Environment (Surface Geochemistry)

Iron and zinc isotope fractionation during uptake and translocation in rice (*Oryza sativa*) grown in oxic and anoxic soils[☆]



Tim Arnold^a, Tamara Markovic^a, Guy J.D. Kirk^b, Maria Schönbacher^c,
Mark Rehkämper^a, Fangjie J. Zhao^d, Dominik J. Weiss^{a,e,*}

^a Imperial College London, Department of Earth Science and Engineering, London, United Kingdom

^b Cranfield University, School of Energy, Environment & Agrifood, Cranfield, United Kingdom

^c Eidgenössische Technische Hochschule, Earth Sciences, Zürich, Switzerland

^d Rothamsted Research, Sustainable Soil and Grassland Systems Department, Harpenden, United Kingdom

^e Stanford University, School of Earth Science, Stanford, United States

ARTICLE INFO

Article history:

Received 30 November 2014

Accepted after revision 11 May 2015

Available online 2 July 2015

Keywords:

Isotopes

Rice (*Oryza sativa*)

Anaerobic Soils

Iron

Zinc

ABSTRACT

Stable isotope fractionation is emerging quickly as a powerful novel technique to study metal uptake and translocation in plants. Fundamental to this development is a thorough understanding of the processes that lead to isotope fractionation under differing environmental conditions. In this study, we investigated Zn and Fe isotope fractionation in rice grown to maturity in anaerobic and aerobic soils under greenhouse conditions. The overall Zn isotope fractionation between the soil and above ground plant material was negligible in aerobic soil but significant in anaerobic soil with isotopically lighter Zn in the rice plant. The observed range of fractionation is in line with previously determined fractionations of Zn in rice grown in hydroponic solutions and submerged soils and emphasizes the effect of taking up different chemical forms of Zn, most likely free and organically complexed Zn. The Zn in the grain was isotopically lighter than in the rest of the above ground plant in rice grown in aerobic and anaerobic soils alike. This suggests that in the course of the grain loading and during the translocation within the plant important biochemical and/or biophysical processes occur. The isotope fractionation observed in the grains would be consistent with an unidirectional controlled transport from shoot to grain with a fractionation factor of $\alpha \approx 0.9994$. Iron isotopes showed an isotopic lighter signature in shoot and grain compared to the bulk soil or the leachate in aerobic and anaerobic soils alike. The negative direction of isotopic fractionation is consistent with possible changes in the redox state of Fe occurring during the uptake and translocation processes. The isotope fractionation pattern between shoots and grain material are different for Zn and Fe which finally suggests that different mechanisms operate during translocation and grain-loading in rice for these two key micronutrients.

© 2015 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

Soils deficient in zinc (Zn) and iron (Fe) constrain rice (*Oryza sativa*) production in large parts of the world (Dobermann and Fairhurst, 2000) and deficiencies of Zn and Fe in human populations with rice-based diets cause

[☆] Revised version submitted for publication to C. R. Geoscience.

* Corresponding author. Imperial College London, Department of Earth Science and Engineering, London, United Kingdom.

E-mail address: d.weiss@imperial.ac.uk (D.J. Weiss).

major health problems for millions of people (Graham, 2007). Identifying nutrient efficient rice lines, applying appropriate breeding techniques, and understanding the underlying mechanisms of Zn and Fe efficiency, are important for tackling this large societal challenge (Wissuwa et al., 2008).

Recent work demonstrated significant isotope fractionation during uptake and translocation of trace metals including Fe, Cu and Zn from soils to plants in a wide variety of plant species and this suggests that natural stable isotope compositions provide a powerful technique to study the chemical and biological processes that control micronutrient movement from soils into plants (Alvarez-Fernandez et al., 2014). Isotopic fractionations differ between plant species and between genotypes of the same species, as well as with soil and nutritional conditions (Arnold et al., 2010a; Aucour et al., 2011; Deng et al., 2014; Tang et al., 2012; Weiss et al., 2005; Weinstein et al., 2011).

Work on the isotopic fractionation during Zn uptake in hydroponic cultures showed that plant shoots are enriched with ^{64}Zn relative to ^{66}Zn in comparison to the source, consistent with an unidirectional process during transport of free Zn^{2+} across cell membranes (Aucour et al., 2011; Deng et al., 2014; Weiss et al., 2005). Plants grown in natural soils, however, show negligible or slight heavy isotope enrichment in the shoots (Arnold et al., 2010a; Tang et al., 2012; Viers et al., 2007). Recent work suggests that uptake of a Zn(II)-phytosiderophore (PS–Zn) complex could account for an observed heavy isotopic fractionation in Zn uptake by rice (Arnold et al., 2010a). Likewise Smolders and co-workers account heavy isotopic fractionation during Zn uptake by tomato plants in resin-buffered hydroponics by the uptake of a PS–Zn complex (Smolders et al., 2013). Investigations of Zn complexation by organic molecules using experiments in the laboratory (Jouvin et al., 2009) and ab initio calculations (Fujii and Albarède, 2012) support this model. Few studies determined the stable isotope fractionation of Zn during translocation in higher plants but enrichment with light isotopes in leaves with increasing distance from the root (Caldelas et al., 2011; Moynier et al., 2009; von Blankenburg et al., 2009) and during the translocation from root to shoot (Caldelas et al., 2011; Jouvin et al., 2012; Tang et al., 2012) have been reported.

To date, Fe isotope fractionation has not been studied in rice but a significant body of work exists for graminaceae (Guelke and von Blankenburg, 2007; Kiczka et al., 2010b; von Blankenburg et al., 2009; Guelke et al., 2010). The observed range of isotope fractionation is around 2.25‰ per atomic mass unit (pamu). Initial pot studies found an enrichment with light Fe during plant uptake in strategy-I plants as opposed to an enrichment with heavy isotopes in strategy-II ones (Guelke and von Blankenburg, 2007). These observations are in line with known plant uptake and isotope fractionation mechanism, whereby a reduction reaction and subsequent uptake of ferrous iron is responsible for the enrichment with light Fe isotopes in strategy-I plants, and the complexation of Fe^{3+} by organic ligands and the subsequent uptake of the PS–Fe complex is responsible for the enrichment with heavy isotopes in

strategy-II plants. Subsequent work found significant variation in isotopic fractionation among different graminaceous species, showing positive and negative isotope signatures suggesting that the controls are far more complex (Guelke-Stelling and von Blankenburg, 2012; Kiczka et al., 2010b). One proposed explanation is that strategy-II plants possess transporters for ferrous Fe and thus can take up Fe^{2+} directly (Cheng et al., 2007; Kim and Guerinot, 2007). Kiczka et al., 2010b suggested two consecutive stages during Fe uptake to explain light Fe enrichment in graminaceae, and in particular processes preceding active transport such as mineral dissolution which favours for trace metals the removal of light isotopes (Chapman et al., 2009; Kiczka et al., 2010a, 2011; Weiss et al., 2014; Wiederhold et al., 2006, 2007a, 2007b) and selective Fe uptake at the plasma membrane level. These more recent observations emphasize the possible effect of mixing of different chemical forms on the Fe and other trace metal isotope signature found in stems, grains and leaves during the transport in phloem and xylem (Guelke-Stelling and von Blankenburg, 2012; Moynier et al., 2013). Although Fe speciation during root uptake is a major contributor to the final isotopic signature, possible mechanisms that influence isotope distribution within plants include successive oxidation and reduction steps during translocation, ligand exchange reactions, remobilisation from older plant tissues and mixing effects of short- (phloem) and long-distance (xylem) transport (Alvarez-Fernandez et al., 2014; Yoneyama et al., 2010).

The aim of the present study was to investigate the controls of stable isotope fractionation of Zn and Fe in rice (*Oryza sativa*) grown under aerobic and anaerobic soil conditions. We conducted experiments in a nutrient-sufficient soil with a widely grown rice genotype. We analysed soil, leachable fractions, shoots and grains, and we discuss the observed fractionation patterns in light of the possible mechanisms outlined above.

2. Materials and methods

2.1. Plant growth experiments

Rice (*Oryza sativa* L. cv. Oochikara) was grown in pots in a greenhouse at Rothamsted Research under anaerobic (flooded) and aerobic conditions as described elsewhere (Xu et al., 2008). The soil was taken from the plough layer (0–20 cm depth) of an arable field at Rothamsted. It is a moderately well-drained Aquic Paleudalf (USDA classification) or Luvisol (FAO classification) with silty clay loam texture (26% clay, 53% silt, 21% sand), 2% organic C, 0.2% total N, 5.93 g kg⁻¹ amorphous Fe oxides (by ammonium oxalate extraction) and pH 6.4. The soil was air-dried, sieved to < 8 mm and homogenized. Two watering treatments (aerobic and flooded) with four pots per treatment were randomly arranged on a bench inside a greenhouse (day/night temperatures 28/25 °C, light period 16 h per day with natural sunlight supplemented with sodium vapour lamps to maintain light intensity of > 350 μmol m⁻²s⁻¹). Five pre-germinated rice seeds were planted in each pot.

To grow rice under aerobic conditions, plastic pots with holes at the bottom were placed on saucers and filled with 1 kg of soil, while for anaerobic conditions pots without holes were used. Fertilisers (120 mg N·kg⁻¹ soil as NH₄NO₃/(NH₄)₂SO₄, 30 mg P·kg⁻¹ and 75.5 mg K·kg⁻¹ soil as K₂HPO₄) were added to the soil and mixed thoroughly. Deionised water was added to full saturation with 2 cm of standing water for anaerobic rice, and to 70% of the soil's water holding capacity for aerobic rice. The two watering regimes were maintained throughout the experiment by daily additions of deionised water. Two further doses of 50 mg of N (NH₄NO₃) were added to each pot at 33 and 70 days after planting. The plants were harvested at grain maturity (117 days after planting). Stems were cut at 2 cm above soil surface, rinsed with deionised water, and dried at 60 °C for 48 h. The samples were separated into straw and grain (including husk), ground, pooled in one sample per pot and then aliquots were processed as described below. Root material was not collected for analysis because a large fraction of the roots had died and degenerated when the plants reached maturity. Only the above ground shoot materials were used.

2.2. Reagents, sample preparation, ion exchange chromatography and mass spectrometry

Sample preparation was carried out in Class-10 laminar flow benches within class 1000 clean room facilities. Mineral acids except for HF were purified from Analar grade by sub boiling distillation in quartz stills. Distilled HF (~28 M Ultrapure grade) and H₂O₂ (~30% v/v Aristar grade) were purchased from VWR. All acid dilutions used > 18.2 MΩ cm⁻¹ grade H₂O from a Millipore purification system.

Oven-dried plant samples were ground using a porcelain pestle and mortar with liquid nitrogen and passed through a 0.5 mm² sieve. Plant samples were digested using a CEM Microwave Accelerated Reaction System (MARS) with 100-mL XP-1500 Teflon vessels. 5 mL of 15 M HNO₃, 3 mL 30% H₂O₂, and 0.6 mL 28 M HF were added to approximately 0.35 g of material. The acid-solid mixture was ramped to 210 °C and 250 psi and held for 90 min. Soil samples (50 mg) were digested using the same protocol. Soil leaches were used to estimate the composition of the Zn pool available to the plants. 30 mL 0.1 M HCl was added to 0.3 g of air-dried soil, and the resulting suspension was stirred continuously for 48 h at 25 °C. The suspension was centrifuged, and the supernatant was collected.

After digestion, samples were evaporated to dryness and the residue was dissolved in ~1 mL 7 M HCl with 0.05 mL 30% H₂O₂ added to prevent reduction of ferric Fe. The solutions were dried again and re-dissolved in 2.5 mL 7 M HCl containing 0.001% H₂O₂ by volume. These solutions were split into three aliquots for Zn and Fe concentration measurements, for isotopic analysis, and for archive. The 0.5-mL aliquots for concentration measurement were diluted to 3.5 mL and made up to 1 M HCl and the measured using a Varian quadrupole ICPMS.

The separation of Zn and Fe from the matrix for subsequent isotope ratio measurements utilized Bio-Rad

Poly-Prep columns filled with 2 mL of AG MP-1 100–200 mesh anion exchange resin. The resin was conditioned with 7 M HCl (10 mL) and successively washed with 0.1 M HNO₃ (10 mL), H₂O (10 mL), 0.1 M HNO₃ (10 mL) and H₂O (10 mL). The resin was re-conditioned with 7 M HCl-0.001% H₂O₂ (7 mL) and the samples loaded (1 mL solution in 7 M HCl-0.001% H₂O₂). The majority of matrix elements were eluted using 30 mL 7 M HCl-0.001% H₂O₂. This was followed by elution of Fe using 17 mL 2 M HCl and Zn using 12 mL 0.1 M HCl. The resin was then washed with 10 mL 0.5 M HNO₃. The Zn and Fe fractions were dried down, dissolved in 0.2 mL 15 M HNO₃, and dried to drive off residual HCl. Finally, 1 mL 0.1 M HNO₃ was added and the samples refluxed for > 2 hours to ensure complete dissolution. The solutions were then ready to be analysed with the Nu Plasma MC-ICPMS.

The total procedural blank contributions, accounting for the dissolution and ion exchange procedures, were below 1% of the total Zn and Fe for the samples with the lowest concentrations. The effect of these blank contributions on the final isotopic results was therefore negligible compared to the measurement uncertainty and no blank correction was applied.

Zinc isotopes were analysed using the double spike method (Arnold et al., 2010b). An isotope spike enriched with ⁶⁴Zn and ⁶⁷Zn was added to the sample aliquot to achieve a total of 1000 ng of Zn and a spike: sample mass ratio of 1. Values are reported as δ⁶⁶Zn relative to the widely used standard JMC-Lyon, i.e. δ⁶⁶Zn_{JMC-Lyon} = [(⁶⁶Zn/⁶⁴Zn)_{sample} / (⁶⁶Zn/⁶⁴Zn)_{JMC-Lyon} - 1] × 10³. The Fe isotope analyses were performed by standard-sample bracketing using the reference material IRMM-014 as the standard, as detailed elsewhere (Chapman et al., 2009). Values are reported as δ⁵⁶Fe relative to the IRMM-14 standard, δ⁵⁶Fe_{IRMM-014} = [(⁵⁶Fe/⁵⁴Fe)_{sample} / (⁵⁶Fe/⁵⁴Fe)_{IRMM-014} - 1] × 10³.

Certified reference materials (rye grass BCR-281 and blend ore BCR-027) and single element industrial solutions (Johnson Matthey and Puratronic Alpha Aesar) were analysed to determine accuracy and precision of the isotope measurements (see Table 1 for details).

3. Results and discussion

3.1. Zinc content and isotopic ratios of plants and soil

The average Zn concentration of the shoot tissue grown in aerobic soil is 75.5 ± 12.3 μg·g⁻¹ (mean ± 1SD, n = 4) and approx. 2.3 higher than in rice grown in anaerobic soil (32.6 ± 3.5 μg·g⁻¹, ± 1SD, n = 4, Table 1). All measured concentrations are well above the threshold for deficiency, which is below 10 μg·g⁻¹ (Dobermann and Fairhurst, 2000; Smolders et al., 2013), suggesting the plants were Zn sufficient in both soils. This conclusion is supported by the shoot and grain biomasses, which were not significantly different for the two water regimes, despite the difference in Zn concentration of the shoot (Table 1). The lower Zn uptake under anaerobic conditions is expected because of the reduced Zn solubility in submerged soils (Kirk, 2004). Under aerobic soil conditions, the shoots, i.e. stems and leaves, contain a greater proportion of the total plant Zn (75% versus 63% under anaerobic conditions). The total Zn concentration

Table 1

Dry mass, element concentrations and δ -values for Zn and Fe in shoots, grain and total material above ground of rice plants grown in aerobic and anaerobic soil. Plants were harvested 117 days after sowing. Also shown are the isotopic compositions of soil measured in the 0.1 M HCl extract and in bulk soil and in standard reference materials measured during this and previous studies (Caldelas et al., 2011; Chapman et al., 2006, 2009).

Sample	Treatment	Comment	n	Zn/ $\mu\text{g g}^{-1}$	$\delta^{66}\text{Zn}_{\text{JMC-Lyon}}/\text{‰}$	Fe/ $\mu\text{g g}^{-1}$	$\delta^{56}\text{Fe}_{\text{IRMM-014}}/\text{‰}$	Dry mass/g	
Rice	Anaerobic soil	Shoot	Measured	4	32.6 ± 3.5	0.61 ± 0.09	95.2 ± 5.2	-0.32 ± 0.16	11.9 ± 1.8
		Grain	Measured	4	28.5 ± 0.5	-0.18 ± 0.10	13.8 ± 0.6	-0.39 ± 0.19	8.0 ± 1.6
		Total plant above ground	Calculated by mass balance	4		0.32 ± 0.09		-0.33 ± 0.16	
Rice	Aerobic soil	Shoot	Measured	4	75.5 ± 12.3	0.73 ± 0.12	96.1 ± 11.2	-0.40 ± 0.07	12.5 ± 2.6
		Grain	Measured	4	33.9 ± 3.6	-0.00 ± 0.07	14.8 ± 1.3	-0.23 ± 0.16	9.3 ± 2.4
		Total plant above ground	Calculated by mass balance	4		0.51 ± 0.04		-0.35 ± 0.07	
Soil	Bulk	0.1 M HCl extract	Measured	2		0.36 ± 0.04		0.05 ± 0.01	
			Measured	2		0.59 ± 0.02		-0.27 ± 0.02	
Reference material	Rye grass	This study	Caldelas et al. (2011)	3		0.40 ± 0.09		0.32 ± 0.07	
						0.50 ± 0.10			
	BCR 027 Blend Ore	This study	Chapman et al. (2006)	4		0.25 ± 0.06			
						0.33 ± 0.07			
	Johnson Matthey	This study	Chapman et al. (2009)	3				0.44 ± 0.04	
								0.49 ± 0.09	
Puratronic Alpha Aesar	This study	Chapman et al. (2009)	7				0.12 ± 0.13		
							0.10 ± 0.09		

Isotope data are given as mean \pm 2 SD and all the other data as mean \pm 1 SD. The plants were harvested after 117 days.

of the soil is $105 \mu\text{g.g}^{-1}$ whilst the 0.1 M HCl soil leach extracted $26 \mu\text{g.g}^{-1}$.

The Zn isotope compositions of the rice shoots, grains and the total above ground plant material are reported in Table 1. The total above ground plant value was calculated from the mass of the shoot and grain material and their corresponding Zn contents and isotope compositions. We find that the shoots of the rice plants grown under both soil regimes are isotopically significantly heavier than the bulk soil or leach (Fig. 1). The grains, in contrast, are isotopically light. The calculated Zn isotope composition of the total above ground plant material is closer to that of the shoot and soil leach and 0.19‰ heavier in the rice grown under aerobic conditions (Table 1). This difference is small but significant within the 95% confidence interval.

We note a small difference in the isotope ratios in rice grown in aerobic or anaerobic soils. The shoots of rice grown in the aerobic soil yield a heavier isotope composition than the Zn in the labile soil pool. This is balanced by a bias towards light isotopes of Zn between shoot and grain, resulting in a small net negative bias between Zn in the total plant and the Zn in the soil leach. Under anaerobic soil conditions, the enrichment with heavy isotopes between soil leach and shoot transfer is smaller, and the net light

bias between total above ground plant and soil solution is larger (Fig. 1).

Bulk and soil leach fractions show isotopic compositions ($\delta^{66}\text{Zn}_{\text{JMC-Lyon}}$) of 0.36 ± 0.04 and $0.59 \pm 0.02\text{‰}$ (mean \pm 1 SD, $n = 2$), respectively (Table 1). By mass balance, the Zn fraction remaining in the solid phase has a $\delta^{66}\text{Zn}_{\text{JMC-Lyon}}$ of ca. 0.28‰ . This lies well within the range of igneous rocks and/or the inferred range of geological materials (Cloquet et al., 2008). It is possible that the 0.1 M HCl leachable fraction represents Zn from polluted atmospheric deposition, which can account for the heavier isotopic signature compared to the bulk soil (Weiss et al., 2014; Gioia et al., 2008).

3.2. Isotope fractionation processes of Zn during plant uptake and translocation

The isotopic fractionation between the different reservoirs is assessed employing the $\Delta^{66}\text{Zn}$ notation calculated using the following equation (Weiss et al., 2013):

$$\Delta^{66}\text{Zn}_{\text{A-B}} = \delta^{66}\text{Zn}_{\text{A}} - \delta^{66}\text{Zn}_{\text{B}}$$

where $\Delta^{66}\text{Zn}_{\text{A-B}}$ is the isotopic fractionation between reservoirs A and B (i.e. soil leach, bulk soil, grain, shoot or




Fig. 1. (Colour online.) Zinc (a) and Fe (b) isotope variations, expressed as $\delta^{66}\text{Zn}_{\text{JMC-Lyon}}$ and $\delta^{56}\text{Fe}_{\text{IRMM-014}}$, respectively, between bulk soil, soil leachate, shoot, grain and total above ground material.

above ground plant) and $\delta^{66}\text{Zn}_{\text{A,B}}$ are the isotopic signatures of the reservoir A and B. The results are given in Table 2.

The observed fractionations between the total above ground plant and the soil leach (-0.27‰ in anoxic soils and -0.08‰ in oxic soils) are well in line with that found in rice grown in hydroponic solution (Smolders et al., 2013; Weiss et al., 2005) and in submerged soils (Arnold et al., 2010a). The signatures in these previous studies were explained by the kinetically controlled uptake of free Zn^{2+} in the hydroponic studies and by the uptake of PS–Zn complexes in the submerged soils. Complexation to a phytosiderophore

ligand in the soil solution prefers isotopically heavy Zn as shown by experimental and computational work (Fujii and Albarède, 2012; Fujii et al., 2014; Jouvin et al., 2009). Zinc has low solubility in aerobic and anaerobic soils and graminaceous plant roots, including rice, therefore secrete phytosiderophores to absorb the resulting Zn-phytosiderophore complexes (Reid et al., 1996; von Wiren et al., 1996; Widodo et al., 2010). We expect that phytosiderophore secretion is more pronounced in aerobic soil to solubilise Fe (III) and this indeed accounts for the higher Zn capture and the incorporation of heavier isotopes under aerobic soil conditions observed during this study (Fig. 1). We note that under Zn depletion in the soil solution, the isotopic composition of the total plant above the surface would trend towards that of the initial soil reservoir because the amount taken up approaches the total amount available in solution. The increased uptake of Zn observed under aerobic conditions and the isotopic composition of the total plant above the surface being closer to that of the soil leachate would be in line with this interpretation. However, mass balance calculations suggest that the rice extracted only a small fraction of the Zn that was available for extraction from the soil. The soil per pot contained approximately 26 mg Zn compared to 1.26 mg and 0.62 mg that were present in the plants under aerobic and anaerobic soil conditions, respectively (Table 1).

The grains show a significant enrichment with light Zn isotopes relative to the shoot material in rice grown under aerobic and anaerobic conditions alike (Table 2). The Zn budget of the shoot is accumulated over most of the growth season, whereas the grains largely accumulate their Zn during the grain-filling period of approximately 30 days. Previous work suggests that the rapid accumulation of Zn during the relatively short grain-filling period is controlled by the remobilisation of Zn from various parts of the plant and not by increased uptake from the soil medium (Nishiyama et al., 2012; Yoneyama et al., 2010). Our data indeed agrees with a stable isotope fractionation model for a unidirectional process such as translocation from shoot to grain (Fig. 2), the Rayleigh model is valid assuming that (i) the Zn is remobilized from the shoot during grain-loading and the soil as additional source is excluded, (ii) the Zn pool of the shoot is not significantly replenished from the soil during this time period, and (iii) the Zn transport is not quantitative. The unidirectional model equation is given by:

$$F^{(\alpha_{B-A}-1)} = \frac{(1000 + \delta^{66}\text{Zn}_A)}{(1000 + \delta^{66}\text{Zn}_{A,0})}$$

where α is the isotopic fractionation factor between shoot and grain, F the fraction of Zn in the shoot, $\delta^{66}\text{Zn}_{\text{A},0}$ the initial isotopic composition of the shoot prior to the grain-filling period, and $\delta^{66}\text{Zn}_A$ the isotopic composition of the shoot corresponding to the fraction, F , which remains in the shoot. Our experimental results suggest an initial offset between grain and shoot of -0.6‰ ($\alpha \approx 0.9994$) followed by a slight decrease to the measured values. Within the framework of this model, a greater extent of remobilisation

Table 2Isotopic fractionations, expressed as $\Delta^{66}\text{Zn}$ and $\Delta^{56}\text{Fe}$, between shoot, grain, total shoot and soil extract in 0.1 M HCl.

Treatment	$\Delta^{66}\text{Zn}_{\text{shoot-soil leach}}/\text{‰}$	$\Delta^{66}\text{Zn}_{\text{grain-shoot}}/\text{‰}$	$\Delta^{66}\text{Zn}_{\text{plant above ground-soil leach}}/\text{‰}$	$\Delta^{56}\text{Fe}_{\text{shoot-soil}}/\text{‰}$	$\Delta^{56}\text{Fe}_{\text{grain-shoot}}/\text{‰}$	$\Delta^{56}\text{Fe}_{\text{plant above ground-soil leach}}/\text{‰}$
Anaerobic	0.02 ± 0.09	-0.79 ± 0.16	-0.27 ± 0.09	-0.37 ± 0.16	-0.06 ± 0.10	-0.38 ± 0.16
Aerobic	0.14 ± 0.12	-0.67 ± 0.17	-0.08 ± 0.04	-0.45 ± 0.07	0.14 ± 0.09	-0.40 ± 0.07

($F < 0.6$) would result in even larger isotopic effects. We note the good fit of the model in our and previous studies testing the fractionation of stable Zn isotopes in Zn hyper accumulators and Zn tolerant plants (Aucour et al., 2011; Deng et al., 2014; Tang et al., 2012) but we acknowledge too that the mechanisms for Zn transport into the root and subsequently into the grain via xylem and/or phloem are highly complex (Alvarez-Fernandez et al., 2014; Nishiyama et al., 2012; Yamaji and Ma, 2014; Yoneyama et al., 2010) and a one-source one-sink model well be far too simplistic. The xylem is the only vascular tissue that reaches the developing grain and the grain tissue (Patrick and Offler, 2001). During the transport via the phloem, Zn must enter/re-enter the symplasm at least three times to reach the grain, i.e., at the root prior to the endodermis, during the subsequent transition into the phloem and during the crossover from the phloem into the grain tissue (Broadley et al., 2007).

3.3. Iron content and isotopic ratios of the plants and soil

Plant Fe concentrations and the total amount of Fe do not differ significantly between the rice plants grown in

the two contrasting soil environments (Table 1). In both treatments, the shoot Fe concentrations are well above the threshold range for deficiency, i.e., 50 to 100 $\mu\text{g}\cdot\text{g}^{-1}$ (Dobermann and Fairhurst, 2000). The Fe isotopic composition of the 0.1 M HCl leachate is about 0.3‰ lighter in $\delta^{56}\text{Fe}_{\text{IRMM-014}}$ than the bulk soil (Table 1). This is likely due to partial leach of selected mineral phases with an isotopically light signature and/or a kinetic mechanism whereby the lighter isotopes are readily removed (Chapman et al., 2009; Kiczka et al., 2010a; Wiederhold et al., 2006). The $\delta^{56}\text{Fe}_{\text{IRMM-014}}$ value of the soil leach is -0.27‰ (Table 1).

3.4. Isotope fractionation of Fe during uptake, translocation and grain-filling

The fractionations between plant tissues and the source are calculated for the Fe relative to the bulk soil as this enables us to compare our results with previous work (Guelke and von Blanckenburg, 2007; Guelke-Stelling and von Blanckenburg, 2012). We find an enrichment with isotopically light Fe between bulk soil and shoot of up to 0.5‰ in aerobic and anaerobic soils alike (Table 2). The




Fig. 2. Isotope fractionation model for unidirectional translocation of Zn from shoot to grain (expressed in fraction of above ground Zn in grain; see text for details regarding the model). The two data points represent the average $\Delta^{66}\text{Zn}$ values between grains and shoot for rice grown under anaerobic and aerobic conditions. Full arrows: data from this study, open arrows: data from literature including transfer from (1) seawater into algae (John et al., 2007), (2) seed to leaf in lentils and tuber to leaf in bamboo (Moynier et al., 2009), and (3) solution into tomato plant (Weiss et al., 2005). Only data from controlled systems are included, where sources and sinks can be unequivocally identified.

largest difference in $\Delta^{56}\text{Fe}$ value is between shoot and bulk soil of plants grown on aerobic soils (0.45‰). The Fe taken up into the shoots has a significant lighter isotopic composition compared to the bulk soil but only insignificantly lighter relative to the soil leach (Table 1 and Fig. 1).

A notable finding is the similarity in the Fe isotopic composition between rice grown in aerobic and anaerobic soils, and the small fractionation between reservoirs. This suggests that the different physical-chemical conditions in the soils, i.e. redox potential, oxygen fugacity, solubility, do not affect the isotope signature in the plant, despite of the different Fe redox states and the variable amounts of Fe(III)–PS complexes likely present. The extent of enrichment with light isotopes in the shoot material observed is significantly smaller than that found in previous work (Guelke and von Blanckenburg, 2007; Kiczka et al., 2010b) and this observation suggests significant uptake of Fe in its reduced form. Ferrous Fe is taken up without the need for preceding phyto siderophore-facilitated solubilisation in the rice rhizosphere (Ishimaru et al., 2006).

The rice does not show significant enrichment with light isotopes in the grain in disagreement with previous work on oat (Guelke and von Blanckenburg, 2010; Guelke-Stelling and von Blanckenburg, 2012). At present, this difference is not easily explainable and further work is needed. Iron, similar to Zn, is transported into the symplasm via the apoplasm before grain-loading. The lack of isotopic fractionation could suggest a quantitative transport of complexed Fe (III) from the soil solution to the grain.

4. Conclusions

We find large negative fractionation of Zn during the translocation of from shoots into grain. The direction and magnitude of the shoot to grain fractionation is consistent with other studies and with the concept of enrichment with light isotopes into higher levels of the food chain. The different extend of isotope fractionation between Zn in the soil leach and the total above ground plant suggests that a mixture of different Zn species are taken up depending on the soil conditions, likely including free Zn^{2+} and PS–Zn complexes. Within experimental error, Fe isotopes showed the same isotopic pattern between the different plant reservoirs and in rice grown in the two different soil environments. Finally, the different isotope patterns observed for Fe and Zn suggest that the remobilisation and subsequent transport does not occur via the same mechanism.

Acknowledgements

This paper is dedicated to Jean, Caroline, Olivier and Jules. The work was partly funded by a grant from the UK's Biotechnology and Biological Sciences Research Council (BBSRC, Grant Ref. BB/J011584/1) under the Sustainable Crop Production Research for International Development program, a joint multinational initiative of BBSRC, the UK Government's Department for International Development and (through a grant awarded to BBSRC) the Bill & Melinda

Gates Foundation. We acknowledge financial support from Stanford University (to DJW through the Cox Visiting Fellowship) and from Imperial College London (to TA through a PhD scholarship). Rothamsted Research is an institute of the Biotechnology and Biological Sciences Research Council of the UK.

References

- Alvarez-Fernandez, A., Diaz-Benito, P., Abadia, A., Lopez-Millan, A., Abadia, J., 2014. Metal species involved in long distance metal transport in plants. *Front. Plant Sci.* 5, 1–20.
- Arnold, T., Kirk, G.J.D., Wissuwa, M., Frei, M., Zhao, F.J., Weiss, D.J., 2010a. Evidence for the mechanisms of zinc efficiency in rice using isotope discrimination. *Plant Cell Environ.* 33, 370–381.
- Arnold, T., Schonbachler, M., Rehkammer, M., Dong, S., Zhao, F.J., Kirk, G.J., Coles, B.J., Weiss, D.J., 2010b. Measurement of zinc stable isotope ratios in biogeochemical matrices by double-spike MC-ICPMS and determination of the isotope ratio pool available for plants from soil. *Anal. Bioanal. Chem.* 398, 3115–3125.
- Aucour, A.M., Pichat, S., Macnair, M.R., Oger, P., 2011. Fractionation of stable zinc isotopes in the zinc hyperaccumulator *Arabidopsis halleri* and nonaccumulator *Arabidopsis petraea*. *Environ. Sci. Technol.* 45, 9212–9217.
- Broadley, M.R., White, P.J., Hammond, J.P., Zelko, I., Lux, A., 2007. Zinc in plants. *New Phytol.* 173, 677–702.
- Caldelas, C., Dong, S., Araus, J.L., Weiss, D.J., 2011. Zinc isotopic fractionation in *Phragmites australis* in response to toxic levels of zinc. *J. Exp. Bot.* 62, 2169–2178.
- Chapman, J., Mason, T.F.D., Weiss, D.J., Coles, B.J., Wilkinson, J.J., 2006. Chemical separation and isotopic variations of Cu and Zn from five geological reference material. *Geostand. Geoanal. Res.* 30, 5–16.
- Chapman, J.B., Weiss, D.J., Shan, Y., Lemberger, M., 2009. Iron isotope fractionation during leaching of granite and basalt by hydrochloric and oxalic acids. *Geochim. Cosmochim. Acta* 73, 1312–1324.
- Cheng, L., Wang, F., Shou, H., Huang, F., Zheng, L., He, F., Li, J., Zhao, F.J., Ueno, D., Ma, J.F., Wu, P., 2007. Mutation in nicotianamine aminotransferase stimulated the Fe(II) acquisition system and led to iron accumulation in rice. *Plant Physiol.* 145, 1647–1657.
- Cloquet, C., Carignan, J., Lehmann, M.F., Vanhaecke, F., 2008. Variations in the isotopic composition of zinc in the natural environment and the use of zinc isotopes in geosciences: a review. *Anal. Bioanal. Chem.* 390, 451–463.
- Deng, T., Cloquet, C., Tang, Y.-T., Sterckeman, T., Echevarria, G., Estrade, N., Morel, J.-L., Qiu, R.-L., 2014. Nickel and zinc isotope fractionation in hyperaccumulating and nonaccumulating plants. *Environ. Sci. Technol.* 48, 11926–11933.
- Dobermann, A., Fairhurst, T., 2000. Rice nutrient disorders and nutrient management. Potash and Phosphate Institute, International Rice Research Institute, Manila, Singapore.
- Fujii, T., Albarède, F., 2012. Ab initio calculation of the Zn isotope effect in phosphates, citrates, and malates and applications to plants and soil. *PLoS ONE* 7, 0030726. <http://dx.doi.org/10.1371/journal.pone.0030726>.
- Fujii, T., Moynier, F., Blichert-Toft, J., Albarède, F., 2014. Density functional theory estimation of isotope fractionation of Fe, Ni, Cu and Zn among species relevant to geochemical and biological environments. *Geochim. Cosmochim. Acta* 140, 553–576.
- Gioia, S., Weiss, D.J., Coles, B.J., Arnold, T., Babinski, M., 2008. Accurate and precise measurements of Zn isotopes in aerosols. *Anal. Chem.* 80, 9776–9780.
- Graham, R.D., 2007. Nutritious subsistence food systems. *Adv. Agron.* 92, 1–74.
- Guelke, M., von Blanckenburg, F., 2007. Fractionation of stable iron isotopes in higher plants. *Environ. Sci. Technol.* 41, 1896–1901.
- Guelke, M., von Blanckenburg, F., Schoenberg, R., Staubwasser, M., Stuerzel, H., 2010. Determining the stable Fe isotope signature of plant available iron in soils. *Chem. Geol.* 277, 269–280.
- Guelke-Stelling, M., von Blanckenburg, F., 2012. Fe isotope fractionation caused by translocation of iron during growth of bean and oat as models of strategy I and II plants. *Plant Soil* 352, 217–231.
- Ishimaru, Y., Suzuki, M., Tsukamoto, T., Suzuki, K., Nakazono, M., Kobayashi, T., Wada, Y., Watanabe, S., Matsuhashi, S., Takahashi, M., Nakanishi, H., Mori, S., Nishizawa, K., 2006. Rice plants take up iron as a Fe^{3+} phyto siderophore and as Fe^{2+} . *Plant J.* 45, 335–346.
- John, S.G., Geis, R.W., Saito, M.A., Boyle, E.A., 2007. Zinc isotope fractionation during high-affinity and low-affinity zinc transport by

- the marine diatom *Thalassiosira oceanica*. *Limnol. Oceanogr.* 53, 2710–2714.
- Jouvin, D., Louvat, P., Juillot, F., Marechal, C., Benedetti, M., 2009. Zinc isotopic fractionation: why organic matters. *Environ. Sci. Technol.* 43, 5747–5754.
- Jouvin, D., Weiss, D.J., Mason, T.F., Bravin, M.N., Louvat, P., Zhao, F., Ferec, F., Hinsinger, P., Benedetti, M.F., 2012. Stable isotopes of Cu and Zn in higher plants: evidence for Cu reduction at the root surface and two conceptual models for isotopic fractionation processes. *Environ. Sci. Technol.* 46, 2652–2660.
- Kiczka, M., Wiederhold, J.G., Frommer, J., Kraemer, S.M., Bourdon, B., Kretschmar, R., 2010a. Iron isotope fractionation during proton- and ligand-promoted dissolution of primary phyllosilicates. *Geochim. Cosmochim. Acta* 74, 3112–3128.
- Kiczka, M., Wiederhold, J.G., Frommer, J., Voegelin, A., Kraemer, S.M., Bourdon, B., Kretschmar, R., 2011. Iron speciation and isotope fractionation during silicate weathering and soil formation in an alpine glacier forefield chronosequence. *Geochim. Cosmochim. Acta* 75, 5559–5573.
- Kiczka, M., Wiederhold, J.G., Kraemer, S.M., Bourdon, B., Kretschmar, R., 2010b. Iron isotope fractionation during Fe uptake and translocation in alpine plants. *Environ. Sci. Technol.* 44, 6144–6150.
- Kim, S.A., Guerinet, M.L., 2007. Mining iron: iron uptake and transport in plants. *FEBS Letters* 581, 2271–2280.
- Kirk, G.J.D., 2004. *The biogeochemistry of submerged soils*. Cambridge University Press, Cambridge.
- Moynier, F., Fujii, T., Wanga, K., Foriel, J., 2013. Ab initio calculations of the Fe(II) and Fe(III) isotopic effects in citrates, nicotianamine, and phytosiderophore, and new Fe isotopic measurements in higher plants. *C. R. Geoscience* 345, 230–240.
- Moynier, F., Pichat, S., Pons, M., Fike, D., Balter, V., Albarede, F., 2009. Isotopic fractionation and transport mechanisms of Zn in plants. *Chem. Geol.* 267, 125–130.
- Nishiyama, R., Kato, M., Nagata, S., Yanagisawa, S., Yoneyama, T., 2012. Identification of Zn-nicotianamine and Fe-2'-Deoxymugineic acid in the phloem sap from rice plants (*Oryza sativa* L.). *Plant Cell Physiol.* 53, 381–390.
- Patrick, J.W., Offler, C.E., 2001. Compartmentation of transport and transfer events in developing seeds. *J. Exp. Bot.* 52, 551–564.
- Reid, R.J., Brookes, J.D., Tester, M.A., Smith, F.A., 1996. The mechanism of zinc uptake in plants – Characterisation of the low-affinity system. *Planta* 198, 39–45.
- Smolders, E., Versieren, L., Dong, S., Mattioli, N., Weiss, D.J., Petrov, I., Degryse, F., 2013. Isotopic fractionation of Zn in tomato plants reveals the role of root exudates on Zn uptake. *Plant Soil* 370, 605–613.
- Tang, Y.-T., Cloquet, C., Sterckeman, T., Echevarria, G., Carignan, J., Qiu, R.-L., Morel, J.-L., 2012. Fractionation of stable zinc isotopes in the field-grown zinc hyperaccumulator *Noccaea caerulea* and the zinc-tolerant plant *Silene vulgaris*. *Environ. Sci. Technol.* 46, 9972–9979.
- Viers, J., Oliva, P., Nonell, A., Gélabert, A., Sonke, J.E., Freydier, R., Gainville, R., Dupré, B., 2007. Evidence of Zn isotopic fractionation in a soil – plant system of a pristine tropical watershed (Nsimi, Cameroon). *Chem. Geol.* 239, 124–137.
- von Blankenburg, F., Von Wiren, N., Guelke, M., Weiss, D.J., Bullen, T.D., 2009. Metal stable isotope fractionation by higher plants. *Elements* 5, 375–380.
- von Wiren, N., Marschner, H., Romheld, V., 1996. Roots of iron-efficient maize absorb phytosiderophore chelated Zinc. *Plant Physiol.* 111, 1119–1125.
- Weinstein, C., Moynier, F., Wang, K., Paniello, R., Foriel, J., Catalano, J., Pichat, S., 2011. Isotopic fractionation of Cu in plants. *Chem. Geol.*
- Weiss, D.J., Boye, K., Caldelas, C., Fendorf, S., 2014. Fractionation of zinc isotopes during early proton and ligand controlled dissolution of biotite granite and implications for plant studies. *Soil Sci. Soc. Am. J.* 78, 171–179.
- Weiss, D.J., Harris, C., Maher, K., Bullen, T., 2013. A teaching exercise to introduce stable isotope fractionation of metals into geochemistry courses. *J. Chem. Edu.* 90, 1014–1017.
- Weiss, D.J., Mason, T.F.D., Zhao, F.J., Kirk, G.J.D., Coles, B.J., Horstwood, M.S.A., 2005. Isotopic discrimination of Zn in higher plants. *New Phytol.* 165, 703–710.
- Widodo, Broadley, M.R., Rose, T., Frei, M., Parisca-Tanaka, J., Yoshihashi, T., Thompson, M., Hammond, J.P., Aprile, A., Close, T., Ismail, A.M., Wissuwa, M., 2010. Response to zinc deficiency of two rice lines with contrasting tolerance is determined by root growth maintenance and organic acid exudation rates, and not by zinc-transporter activity. *New Phytol.*
- Wiederhold, J.G., Kraemer, S.M., Teutsch, N., Borer, P.M., Halliday, A.N., Kretschmar, R., 2006. Iron isotope fractionation during proton-promoted, ligand-controlled, and reductive dissolution of goethite. *Environ. Sci. Technol.* 40, 3787–3793.
- Wiederhold, J.G., Teutsch, N., Kraemer, S.M., Halliday, A.N., Kretschmar, R., 2007a. Iron isotope fractionation during pedogenesis on redox-imorphic soils. *Soil Sci. Soc. Am. J.* 1840–1850.
- Wiederhold, J.G., Teutsch, N., Kraemer, S.M., Halliday, A.N., Kretschmar, R., 2007b. Iron isotope fractionation in oxic soils by mineral weathering and podzolization. *Geochim. Cosmochim. Acta* 71, 5821–5833.
- Wissuwa, M., Ismail, A.M., Graham, R.D., 2008. Rice grain zinc concentrations as affected by genotype, native soil-zinc availability and zinc fertilisation. *Plant Soil* 306, 37–48.
- Xu, X.Y., McGrath, S.P., Meharg, A.A., Zhao, F.J., 2008. Growing rice aerobically markedly decreases arsenic accumulation. *Environ. Sci. Technol.* 42, 5574–5579.
- Yamaji, N., Ma, J.F., 2014. The node, a hub for mineral nutrient distribution in graminaceous plants. *Trends Plant Sci.* 19, 556–563.
- Yoneyama, T., Goshō, T., Kato, M., Goto, S., Hayashi, H., 2010. Xylem and phloem transport of Cd, Zn and Fe into the grains of rice plants (*Oryza sativa* L.) grown in continuously flooded Cd-contaminated soil. *Soil Sci. Plant. Nutr.* 56, 445–453.