



INSTITUT DE FRANCE
Académie des sciences

Comptes Rendus

Géoscience

Sciences de la Planète

Jean-François Pastre, Évelyne Debard, Catherine Ferrier, Michel Fialin, Bernard Gély, Bertrand Kervazo, Frédéric Maksud, Fatima Mokadem, Sébastien Nomade, Nicolas Rividi and Ségolène Saulnier-Copard

A cryptotephra from the Upper Pleistocene volcanism of the Bas-Vivarais in the sedimentary infilling of the Chauvet-Pont d'Arc cave (Ardèche, France)

Volume 353, issue 1 (2021), p. 67-79

Published online: 4 May 2021

<https://doi.org/10.5802/crgeos.52>



This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*Les Comptes Rendus. Géoscience — Sciences de la Planète sont membres du
Centre Mersenne pour l'édition scientifique ouverte*

www.centre-mersenne.org

e-ISSN : 1778-7025



Original Article — Stratigraphy, Sedimentology

A cryptotephra from the Upper Pleistocene volcanism of the Bas-Vivarais in the sedimentary infilling of the Chauvet-Pont d'Arc cave (Ardèche, France)

Jean-François Pastre^{*,a}, Évelyne Debard^b, Catherine Ferrier^c, Michel Fialin^d, Bernard Gély^e, Bertrand Kervazo^f, Frédéric Maksud^g, Fatima Mokadem^a, Sébastien Nomade^h, Nicolas Rividi^d and Ségolène Saulnier-Copard^a

^a Laboratoire de Géographie Physique, UMR 8591 CNRS, Universités de Paris I et XII, 1 place Aristide Briand, 92195 Meudon Cedex, France

^b 25 rue Paul Chevreton, 69370 Saint-Didier-au-Mont-d'Or, France

^c Université de Bordeaux, PACEA, UMR 5199 CNRS-UB-MC, Allée Geoffroy-Saint-Hilaire, CS 5003, 33615 Pessac Cedex, France

^d Camparis, UPMC, 4 place Jussieu, Tour 46-3^{ème} étage, France

^e DRAC Auvergne-Rhône-Alpes, Pôle Patrimoine, Service Régional de l'Archéologie, Le Grenier d'Abondance, 6 quai Saint-Vincent, 69001 Lyon, France

^f 6 rue du Serment, 24000 Périgueux, France

^g UMR 5608, Laboratoire TRACES, SMP3C, Ministère de la culture et de la communication, Direction Régionale des Affaires Culturelles d'Occitanie, Service Régional de l'Archéologie, Pôle de Toulouse, Hôtel Saint-Jean, 32 rue de la Dalbade, BP 811, 31080 Toulouse Cedex 6, France

^h Laboratoire des Sciences du Climat et de l'Environnement (IPSL-CEA-CNRS-UVSQ) et Université Paris-Saclay, Orme des Merisiers, Bâtiment 714, 91190 Gif-sur-Yvette, France

E-mails: jean-francois.pastre@lgp.cnrs.fr (J.-F. Pastre), evelyne.debard@free.fr (É. Debard), catherine.ferrier@u-bordeaux.fr (C. Ferrier), michel.fialin@upmc.fr (M. Fialin), bernard.gely@culture.gouv.fr (B. Gély), bertrand.kervazo@orange.fr (B. Kervazo), frederic.maksud@culture.gouv.fr (F. Maksud), fatima.mokadem@lgp.cnrs.fr (F. Mokadem), Sebastien.Nomade@lsce.ipsl.fr (S. Nomade), nicolas.rividi@upmc.fr (N. Rividi), segolene.saulnier-copard@lgp.cnrs.fr (S. Saulnier-Copard)

* Corresponding author.

Abstract. A cryptotephra belonging to the Upper Pleistocene volcanism of the Bas-Vivarais was identified for the first time in the sedimentary infilling of the entrance area of the Chauvet-Pont d'Arc cave. This slightly reworked tephra fall is characterized by its heavy minerals assemblage, among which magnesian olivine, enstatite, and chromian diopside issued from peridotites and olivine and diopside of basaltic origin. This composition likely refers to a phreatomagmatic eruption of a maar crater. The ^{14}C datings of the cryptotephra beds are contemporaneous with the Aurignacian settlement and paintings, that is, an age slightly older or close to 36 ka. Among the maar craters from the Bas-Vivarais, the Ray-Pic maar have the closest $^{40}\text{Ar}/^{39}\text{Ar}$ age ($36.2 \pm 11.3/32.2 \pm 11.1$ ka). By contrast, the Vestide du Pal, the biggest maar crater of this country, seems to have a younger age, but the $^{40}\text{Ar}/^{39}\text{Ar}$ dating was measured on a basalt which could have occurred later and been the source of the tephra fall, since its tephra contain the same minerals of unknown origin as those found in Chauvet cave. This discovery strengthens the previous hypothesis according to which the humans living in the Chauvet-Pont d'Arc area during the Upper Paleolithic have witnessed eruptions from the Bas-Vivarais volcanic field and other phenomena.

Keywords. Chauvet-Pont d'Arc cave, Cryptotephra, Phreatomagmatism, Bas-Vivarais volcanism, Upper Pleistocene.

Manuscript received 25th February 2021, revised and accepted 22nd March 2021.

1. Introduction

The Chauvet-Pont d'Arc cave is situated in France in the Ardèche valley near Vallon-Pont-d'Arc ($44^{\circ} 23' 14''$ N, $4^{\circ} 25' 07''$ E, Figure 1A, B, D). Discovered in 1994, it is universally known for the richness and the age of its Paleolithic paintings [Clottes, 2001, Clottes and Geneste, 2007, Delannoy and Geneste, 2020]. The ^{14}C datings of charcoal and paintings indicate two periods of occupation attributed to the Aurignacian and to the Gravettian (from 37.5 to 33.5 ka cal BP and from 31 to 28 ka cal BP; Clottes *et al.*, 1995, Cuzange *et al.*, 2007, Quiles *et al.*, 2016, Valladas *et al.*, 2005). These dates coincide with one of the activity phases of the Bas-Vivarais volcanism. Several recent $^{40}\text{Ar}/^{39}\text{Ar}$ datings have actually dated these eruptions between 29 ± 10 ka and 35 ± 8 ka (95% probability uncertainty Nomade *et al.*, 2016, Sasco, 2015, Sasco *et al.*, 2017). These dates fully cover the ^{14}C AMS datings [Quiles *et al.*, 2016] and the thermoluminescence (TL) datings [Guibert *et al.*, 2015], which correspond to the human settlements responsible for numerous parietal representations in the cave. Among them, spray-shape signs (Figure 1C) were hypothetically ascribed to depiction of strombolian volcanic eruptions [Nomade *et al.*, 2016].

In the context of the pluridisciplinary study of the cave financed by the French Culture Ministry the analysis of heavy minerals of the sediments was undertaken. This study led to the discovery of volcanic minerals attributed to a cryptotephra subcontemporaneous with the sedimentary infilling of the entrance area of the cave. These results, presented

here, corroborate the contemporaneity of the recent volcanism of the Bas-Vivarais with the history of the cave during a period close to its occupation by Aurignacians.

2. Stratigraphical and sedimentological context

The volcanic minerals were identified in the sediments of a limited archeological pit located at about twenty meters from the old entrances which are now filled by deposits related to several processes (Figure 2). Four phases were distinguished in the clogging of the entrance [Debard *et al.*, 2016]:

- The first one constitutes the origin of a gravity scree (= lower scree) mainly due to the gelification of the porch and entry area, to the mechanical expansion of the vaults as well as to the reworking of old superficial deposits. The spreading of the deposits on about forty meters up to the entrance of the Chamber of the Bear Hollows ("Salle des Bauges") is attributed to mobilization by creep and solifluction.
- The second one also corresponds to a gravity scree (intermediate scree) less expanded than the lower one. Their very similar components suggest a same origin.
- The third phase corresponds to blocks generated by the collapse of the cliff. This collapse mainly occurred in two main phases determined by ^{36}Cl cosmogenic dating




Figure 1. (A) Location of the Chauvet cave in France, (B) location of the Chauvet cave in Ardèche area, (C) example of spray-shape sign, (D) perspective view of the Bas-Vivarais volcanism seen from Chauvet cave.

($23,500 \pm 1200$ and $21,500 \pm 1000$ years—Sadier *et al.*, 2012) and gave rise to the definitive cave closure. This set was later covered by fine brown sediments originating from reworking of superficial deposits.

- After the closing of the porch, the ultimate registered events correspond to the desquamation of the equilibrium vaults resulting in small sedimentary clasts, to water runoff, to recent concretions growth, to the settling down of the scree.

The archeological pit (named GE1) carried out between 2006 and 2010 [Gély and Maksud, 2009, Gély *et al.*, 2012] is located in a small gallery connecting the old porch to the Chamber of the Bear Hollows (“Salle des Bauges”) and corresponding to one of the natural paths used by large vertebrates and prehistoric humans (Figure 2). Of small dimensions (50×50 cm), 80 cm deep, the pit intersected the lower

scree and, near the surface, a gravel of clasts issued from the desquamation of the vault.

Six beds have been defined (Figure 3a, b):

- Bed 1: near the surface, this layer is mainly composed of small plates of desquamation (c.1a). Underneath is found dry gravel of elements righted in counterslope (c.1b). Among the clasts of the first level was a fistulous concretion which was dated by U/Th to $13,315 \pm 79$ years [Genty, 2008]. This dating as well as the one obtained on the stalagmite Chau-stm 8, which seals the scree near the archeological pit and whose basis is dated at $11,500 \pm 170$ years [Genty *et al.*, 2005] allow to date this bed between the Bølling-Allerød warming and the beginning of the Holocene (Lateglacial). Level c.1b supplied a flint attributed to the Upper Paleolithic, a trace of the last human incursion before




Figure 2. Location of the archeological pit GE1 located about twenty meters from the Paleolithic entrances. Équipe Grotte Chauvet (Ministère de la Culture, CNRS, Universités).

the porch was finally clogged up and attributed to the Gravettian (31,000–28,000 cal BP; Quiles et al., 2016).

- Bed 2: this bed is a small gravel secondarily packed in a sandy-silty reddish matrix. Some clasts show a soft surface identified as bears' polish. Bed 2b corresponds to the reworking of the uppermost part of bed 3.
- Bed 3: this bed is about 10 cm thick, it is less gravelly, with a sandy and micaceous matrix very rich in charcoals responsible of its dark color. This layer contrasts with the reddish

color of the remaining part of the sequence. It corresponds to the reworking of hearths probably at the origin of the significant thermal impacts observed close to the archeological pit on the ceiling and the walls [Ferrier et al., 2014]. Rubefied clasts or with blackish marks have been found in this bed. Dating of several charcoals (i.e. ranging from 36,000–37,000 cal BP; Quiles et al., 2016) attributes these hearths to the Aurignacian occupancy.

- Bed 4: this bed is divided in two, the upper part corresponding to a small dense gravel




Figure 3. (a) Archeological pit GE1: view of the southern and western sections and dating elements. Picture by C. Ferrier (Equipe Grotte Chauvet—Ministère de la Culture, CNRS, Universités). (b) Log of the archeological pit GE1.

with angular granules due to runoff, while its lower part is coarser. The matrix is sandy-silty reddish brown becoming more yellow toward the bottom.

- Bed 5: this bed is made of an angular gravel packed in a red sandy sediment, more silty toward the base. At its base, a charcoal dated from between 36,361 and 34,797 cal BP [Quiles *et al.*, 2016] is contemporaneous to the first Aurignacian occupation of the cave.
- Bed 6: also, coarse, contains wall elements (one block and clasts) polished by bears. The matrix is more sandy, brownish.
- In beds 2 to 5, the presence of sorted lamellar microstructures and occasionally layered caps at the surface of the clasts shows the role of cold weather (mainly frost periods) in this part of the scree and its spreading by solifluction toward the Chamber of the Bear Hollows.

The volcanic minerals characterizing the Chauvet cryptotephra were identified in the beds 6 to 1b,

indicating a reworking of the initial tephra fall from the outside.

3. Material and methods

The mineralogical study was carried out on the 63 μm –2 mm sand fraction separated by wet sieving in an aqueous solution of sodium hexametaphosphate (dispersant) and removing of carbonates (post-deposition calcite coatings) by HCl 1N/2 hours treatment. Results are based on the study of ten samples distributed in the various beds of the archeological pit. The percentages between the silty-clayey fraction and the clayey fraction characterize clayey-silty sands and clayey-sandy silts. The examination of the light sandy fraction reveals quartz-rich sands (angular or slightly blunt quartz) poor in feldspars, issued from the regional granitic and metamorphic basement. Heavy minerals were separated with bromoform (samples CHAU 1–4 = beds 1b, 2, 3, 4) or with sodium polytungstate (samples CHAU 8–9 = beds 5a and 6, samples ARD 25–28 = beds 3, 4,

5, 6). Thus, they represent the heavy fraction ($d > 2.8$), including some minerals of the basement and all the volcanic minerals potentially present, except possible plagioclases of basaltic origin, not analyzed. These volcanic minerals either correspond to minerals typical of basaltic magma or to mantellic minerals issued from ultramafic xenoliths (peridotites/lherzolites). The percentages of heavy minerals vary between 0.85 and 3% of the sand fraction. The minerals have been identified with a polarization microscope and a binocular lens. In order to establish their frequency, which is slightly variable due to the grain size of the samples, several samples were counted on a squared grid. To specify their respective characterizations and origins, the various minerals identified by optical study were then analyzed with an electron microprobe. The handpicked minerals were included in epoxy resin and polished with abrasive papers and diamond powders. They were then analyzed with a WDS Cameca SX 100 microprobe at the Camparis center (Sorbonne University, Paris VI/CNRS) with column conditions 15 keV, 10 nA. These results were then compared to those of the analysis performed on minerals issued from tephra or peridotitic nodules from the Upper Pleistocene volcanism of Bas-Vivarais (Ardèche).

4. Mineralogy and characterization of the cryptotephra

The results of the heavy minerals counting of six samples are reported in Table 1 and Figure 4. They make it possible to appraise the frequencies of the various species and possibly to link them to the modal composition of the original host rocks. All the samples are largely dominated by olivine (73.8 to 89.7%). This one is mainly composed of xenomorph grains which can result from the mechanical fragmentation of peridotites nodules or from basaltic magma. Some automorphic specimens clearly originate from the last. The orthopyroxenes (xenomorph enstatite 3.9–7.6%) and the green clinopyroxenes (xenomorph chromiferous diopside, 2.5–6.6%) then represent minerals characteristic of peridotites xenoliths [e.g. Berger, 1981]. The brown clinopyroxene (diopside) only plays a minor role (0.7–2%). It probably originates from a basaltic magma together with other pyroxenes such as those occurring in the alkali basalts from the Massif Central. The brown amphiboles, also




Figure 4. Heavy minerals composition of the beds containing the cryptotephra.

little abundant (0.2–2%) are probably peridotites amphiboles as those described by Berger [1981].

These minerals are attributable to a basaltic volcanism rich in peridotites nodules (lherzolites) as well as lower crustal material which can only correspond to the Upper Pleistocene volcanism of the Bas-Vivarais located about 35 km away [e.g. Berger, 1973, 1981, Sasco et al., 2017]. They come from a tephra brought by the wind at the probable origin of most of the other drifts (basement silts and sands) and scattered in the cave environment. The homogeneity of the minerals assemblage and the freshness of the minerals (e.g. automorphic olivine) lead one to think that this slightly reworked tephra is subcontemporaneous with the sediments including the latter. Thus, it can be defined as a cryptotephra. This fall probably results from a phreatomagmatic eruption associated with the formation of a maar. The explosive products (base surges and falls) are actually usually rich in peridotites minerals (see discussion). The origin of the fall from a strombolian eruption is much less probable. In fact, this type of eruption effectively produces basaltic ashes and scoriae which are not present in the Chauvet samples and are also generally poor in minerals.

The other heavy silicates are basement minerals, which play only a minor role (Table 1). These include garnets (0.7 to 5.9%), andalusite, sillimanite-fibrolite, tourmaline, and rutile. They could be alluvial minerals from the Ardèche river, coming from the Massif Central and reworked from old fluvial terraces by runoff or gravity process. But they were more probably brought by the wind. Some could also come from the cryptotephra, the explosive products

Table 1. Heavy minerals spectra of samples containing Chauvet cryptotephra

Sample Bed	CHAU 1 1b	CHAU 2 2	CHAU 3 3	CHAU 4 4	CHAU 8 5a	CHAU 9 6
Brown amphibole	2.01	0.25	0.39	0.28	0.99	0.51
Enstatite	7.38	4.28	3.9	7.26	7.58	3.78
Chromiferous diopside	5.37	2.52	6.64	6.14	5.58	2.41
Brown diopside	0.67	1.26	1.17	1.95	0.39	0.34
Olivine	77.18	86.64	80.07	73.74	80.43	89.5
Andalusite	2.01	0.75	0.78	0.83	0.59	0.34
Sillimanite-fibrolite	0.67		0.39		0.39	0.17
Garnet	0.67	1.26	3.51	5.86	1.79	1.37
Tourmaline	0.67	0.5		0.28		
Rutile	0.67					
Opaques	2.68	2.52	3.12	3.63	2.19	1.37

of the maars being mainly rich in basement minerals. The opaque minerals, little represented (2.5 to 3.6%), correspond either to Fe–Ti oxides issued from the regional basaltic volcanism (magnetite, titanomagnetite), or to silicates and oxides of unknown origin (work in progress), or to rare chromium spinels (about 23% of Cr_2O_3) probably issued from peridotites.

5. Microprobe geochemical study of minerals

In order to precisely characterize the mineralogy and chemical composition of the minerals, a mineralogical study using an electron microprobe was undertaken and compared with the available analyses [Berger, 1981] and new analyses of proximal volcanic falls (base surges) as well as peridotites from the Ardèche volcanism. Various characteristic analyses are reproduced on Table 2.

Olivines are mainly characterized by the variation of their MgO contents. The analysis of automorphic grains reveals compositions comprised between 43.67 and 46.30% of MgO ($\text{Fo} = 83.1\text{--}86.6\%$). Such compositions, poor in Mg, are typical of basaltic phenocrystals as confirmed by their automorphic habitus. After all the cross-checking we have done, it seems that the limit of composition between basaltic and peridotite olivines is at about 47% MgO. The lowering of magnesium in these basaltic olivines (e.g. Table 2, OL1–2) results in their iron enrichment ($\text{FeOt} = 12.65\text{--}15.39\%$ for the most ones) as shown

by the diagram of Figure 5. Except for visibly broken specimens from basaltic origin with MgO contents <47%, the other xenomorphic olivines are more magnesian ($\text{MgO} = 47.40\text{--}48.96\%$) and characterize olivines from peridotites (e.g. Table 2, OL3–4, Figure 3). However, these olivines are slightly less magnesian than the olivine from peridotites of the Vestide du Pal and Chambon maars analyzed by Berger [1981], which mostly exceed 49% of MgO (48.71–49.82%); but based on six analyses performed on the phreatomagmatic tuff of Vestide. We have noticed that they were less magnesian and compatible with the compositions of the Chauvet cryptotephra (44.57–44.98–45.37–47.08–47.44–48.02% MgO). For comparison, the olivines from the Ray-Pic maar peridotites have compositions varying between 47.89 and 49.50% MgO. The olivines of microperidotite enclaves found within the strombolian tuff of Gravenne de Montpezat, were also analyzed. They are slightly more magnesian and mainly exceed 49% MgO. Although small, these variations seem to indicate significant slight differences of composition for the olivines of peridotites xenoliths from the Ardèche volcanism.

The orthopyroxenes necessarily come from peridotites of which they are one of the characteristic minerals with MgO contents near 32–33%. These are enstatites $\text{En}_{87.8\text{--}89.8}$, $\text{Wo}_{0.93\text{--}1.27}$ (Figure 6). Their analytical results (e.g. Table 2, EN1–2) are close to those of Berger [1981] concerning the orthopyroxenes of peridotites from the Vestide du Pal and Chambon

Table 2. Selected microprobe analyses of minerals issued from Chauvet cryptotephra (sample CHAU 4, bed 4)

	OL1	OL2	OL3	OL4	EN1	EN2	CD1	CD2	BD1	BD2
SiO ₂	39.99	40.15	40.96	40.62	56.04	55.48	52.49	52.88	46.68	45.86
Al ₂ O ₃	0.03	0.01	0.10	0.01	3.30	3.40	4.30	4.88	8.80	8.88
TiO ₂	0.00	0.03	0.00	0.04	0.04	0.08	0.33	0.24	2.18	2.53
Cr ₂ O ₃	0.00	0.03	0.02	0.02	0.19	0.29	0.81	0.65	0.05	0.07
FeOt	12.72	13.08	9.95	9.38	7.37	6.21	2.71	2.34	6.14	7.86
MnO	0.18	0.13	0.18	0.14	0.16	0.09	0.02	0.02	0.13	0.20
MgO	46.30	45.74	48.50	48.96	32.69	32.22	16.47	15.77	13.28	12.83
CaO	0.18	0.19	0.03	0.07	0.55	0.63	21.75	21.65	21.62	20.23
Na ₂ O	0.05	0.00	0.02	0.00	0.08	0.06	1.04	1.37	0.75	0.98
NiO	0.33	0.24	0.29	0.34	0.02	0.07	0.02	0.02	0.01	0.05
Total	99.84	99.65	100.12	99.61	100.48	99.58	99.98	99.85	99.71	99.53
Si	0.997	1.003	1.003	0.998	1.933	1.923	1.905	1.916	1.736	1.721
Al	0.001	0.000	0.003	0.000	0.134	0.139	0.181	0.208	0.386	0.392
Ti	0.000	0.000	0.000	0.000	0.001	0.002	0.009	0.006	0.061	0.071
Cr	0.000	0.000	0.000	0.000	0.005	0.008	0.023	0.018	0.001	0.002
Fe	0.265	0.273	0.203	0.192	0.212	0.180	0.082	0.071	0.191	0.246
Mn	0.003	0.002	0.003	0.002	0.004	0.002	0.000	0.000	0.004	0.006
Mg	1.720	1.703	1.77	1.794	1.680	1.716	0.891	0.852	0.736	0.718
Ca	0.004	0.005	0.000	0.002	0.020	0.023	0.846	0.840	0.862	0.813
Na	0.002	0.000	0.001	0.000	0.005	0.004	0.073	0.096	0.054	0.071
Ni	0.006	0.005	0.005	0.006	0.000	0.002	0.000	0.000	0.000	0.001
Total	3.003	2.995	2.995	2.999	3.999	4.002	4.018	4.011	4.035	4.045
Fo	86.6	86.2	89.7	90.3						
Wo					1.06	1.22	46.49	47.65	48.17	45.74
En					87.81	89.4	48.98	48.31	41.14	40.38
Fs					11.11	9.37	4.52	4.02	10.68	13.87

OL1–2: basaltic olivine—OL3–4: peridotite olivine—EN1–2: enstatite—CD1–2: chromiferous diopside—BD1–2: brown diopside. Olivines: numbers of ions on the basis of 4 O, pyroxenes: numbers of ions on the basis of 6 O.

maars. Like them, they are slightly chromiferous ($\text{Cr}_2\text{O}_3 = 0.22\text{--}0.34\%$). The other orthopyroxenes we analyzed (Ray-Pic maar, Borée maar, Gravenne de Montpezat scoria cone, Moula rockshelter tephra) showed relatively similar compositions.

The green clinopyroxene corresponds to the common clinopyroxene of peridotites [Berger, 1981]. This is chromiferous diopside (calcic clinopyroxene $\text{Wo}_{45\text{--}48}$, Figure 5). The results of the ten analyses performed (e.g. Table 2, CD1–2) show slightly less

chromium-rich pyroxenes than those analyzed by Berger [1981] for peridotites of the Vestide du Pal and Chambon maars: $0.54\text{--}0.81\%$ compared to $0.75\text{--}1.36\%$. This difference is not explained. Only one sample differs with $1.34/1.48\%$ Cr_2O_3 .

The brown clinopyroxenes correspond to diopsides $\text{Wo}_{44.97\text{--}48.30}\text{--En}_{40.01\text{--}42.34}\text{--Fs}_{10.68\text{--}13.87}$ (e.g. Table 2 BD1–2, Figure 6). These are basaltic pyroxenes common in the alkali basalts from the Massif Central and attributed to the basaltic magma pulverized




Figure 5. MgO/total FeO diagram of the olivines of Chauvet cryptotephra.




Figure 6. Wo–En–Fs diagram of the pyroxenes of Chauvet cryptotephra.

by the phreatomagmatic eruption. They show typical amounts of Al_2O_3 (8.51–9.46%), TiO_2 (2.15–2.53%), and FeOt (6.14–7.86%).

Analysis of some opaque minerals revealed the existence of several minerals of unspecified origin. The first one presents a chemical composition with 27–34.5% SiO_2 , 5–7% K_2O , 15.4–20.3% FeOt, 13.9–17.7% Al_2O_3 , 5.8–8.8% MgO, 3.8–4.5% TiO_2 . The second one is mainly composed of FeO (67–75% FeOt), but contains low percentages of SiO_2 , Al_2O_3 , CaO, MnO, and P_2O_5 and cannot be ascribed to a true oxide.

6. Discussion, origin, and possible age of the cryptotephra

As previously seen, the mantellic minerals (peridotites minerals = magnesian olivine + orthopyroxene + chromiferous diopside) and the basaltic

minerals (olivine + brown diopside) characterizing the cryptotephra of the Chauvet-Pont d'Arc cave can be ascribed without any doubt to a phreatomagmatic eruption related to a maar of the Upper Pleistocene volcanism of the Bas-Vivarais. They were pulverized and mixed by the phreatomagmatic explosions. The absence of basaltic ash (microscoriae) attributable to a strombolian eruption, the mixing of mantellic and basaltic minerals and the lower explosivity of this type of eruption led us to exclude a strombolian eruption at the origin of the cryptotephra.

The volcanism of the Bas-Vivarais is known since the work of Faujas de Saint-Fond [1778]. Its volcanic products are mainly alkaline basalts derived from a unique enriched mantle source [Chauvel and Jahn, 1984, Downes, 1987]. It benefited in particular from the works of Berger [e.g. Berger, 1973, 1981, 2007] and is distinguished by the frequency of its ultramafic xenoliths [Berger, 1981, Berger and Forette, 1975]. It comprises 17 eruptive centers. Its activity is characterized by phreatomagmatic eruptions (maars) preceded or generally followed by a strombolian activity associated with the emission of basaltic flows filling pre-existing valleys.

The karst of the Bas-Vivarais, situated to the southeast, is characterized by the importance of its Upper Pleistocene infillings [Debard, 1988, 1997]. The signature of volcanic eruptions from the Bas-Vivarais was evidenced in three of these karstic infillings: the Moula rockshelter in Soyons [Debard and Pastre, 2008, Pastre *et al.*, 1994], the Marzal II aven (or Flahault aven), and the Devès du Reynaud aven in Saint-Remèze [Debard and Pastre, 2008]. Among them, the in situ tephra found in Soyons is the more representative example of a phreatomagmatic fall comparable to that detected in Chauvet. Among the recent maars of the Bas-Vivarais, four present an explosive activity and can be at the origin of the cryptotephra we described above: the Ray-Pic, the Vestide du Pal, the Chambon, and the Sapède [Berger, 1981]. This last explosion crater can a priori be excluded, due to its essentially basaltic type [Berger, 1981] as confirmed by our own observations and analyses. Derived tephra are therefore mainly composed of basalt clasts and their olivines are mainly basaltic in composition with 44.01–46.55% MgO ($\text{Fo} = 86.3\text{--}87\%$). The Chambon maar emitted a base surge and lahars visible in the Fau valley above the hamlet. The mineralogical and microchemical study

shows that these products both contain peridotite and basaltic minerals similar to Chauvet' ones. The peridotites olivines (48.43–48.97% MgO) are however more abundant than the basaltic ones in the sample we analyzed. This maar cut in the basement could have been large enough to disperse widely its products, but, due to little extended dispersal in the proximal area, it was probably less extended than those of the following maars. Among the following eruptions, the Vestide du Pal maar constitutes one of the largest explosive craters in the Massif Central (diameter = 1700 m, Berger, 1973, 1981). It emitted important surges visible in its eastern and southern parts (Suc du Pal). The mineralogical study of these products rich in basement minerals pulverized by explosions, show that they contain both peridotite and basaltic minerals. These ones however comprise peridotite olivines (47.05–50% MgO) and peridotite pyroxenes. The basaltic olivines (44.57–45.37% MgO) are rare. However, this composition makes them compatible with the composition of Chauvet cryptotephra ones. The Ray-Pic maar also emitted large surges visible along the road D 215 to the east of the Ray-Pic cascade. The mineralogical study of these tephtras also rich in basement minerals shows that they contain basaltic and peridotite minerals. The olivine shows a composition varying between 44.63 and 49.78% MgO (Fo = 85.2–90.9%), which clearly argues for these two origins and compositions close to those of Chauvet.

The cryptotephra-bearing beds are well dated due to the presence of microcharcoals ascribed to frequenting of the site by Aurignacians. ^{14}C AMS datings gave ages comprised between 36,361–34,797 (bed c5) and 36,100–34,880 (bed c2a) years cal BP [Quiles *et al.*, 2016]. Thus, the cryptotephra must have an age greater than or close to 36 ka. The Bas-Vivarais volcanism benefited from ^{14}C datings [Berger, 1973, Berger *et al.*, 1975, Berger, 2007], from TL datings [Guérin, 1983, Guérin and Gillot, 2007, Sanzelle *et al.*, 2000], from paleomagnetism measurements [Rochette *et al.*, 1993, Sasco, 2015, Sasco *et al.*, 2017], and from K–Ar and $^{40}\text{Ar}/^{39}\text{Ar}$ datings [Sasco, 2015, Sasco *et al.*, 2017]. Among them, the $^{40}\text{Ar}/^{39}\text{Ar}$ datings (*ibid.*) seem to be the most accurate. The $^{40}\text{Ar}/^{39}\text{Ar}$ age of the Vestide du Pal maar would thus be comprised between 25.8 ± 7.7 and 27.2 ± 8.2 ka (*ibid.*). It would thus belong to the recent phase of the Ardèche volcanism. This age seems too young to fit

with the potential age of the Chauvet cryptotephra. Nevertheless, it was performed on a basalt following the phreatomagmatic eruption and not on the tephtras coming from the maar explosions. It is also the same for the age of the Chambon maar dated at 18.8 ± 9.6 ka (*ibid.*). On the other hand, the age of the Ray-Pic maar comprised between 36.2 ± 11.3 and 32.2 ± 11.1 ka (*ibid.*) is well compatible with the ^{14}C datings of the beds containing the minerals of the Chauvet cave. This maar belonging to the middle phase of the Bas-Vivarais volcanism could be at the origin of the cryptotephra we discovered. Nevertheless, its products do not contain the peculiar minerals which are currently studied to define their origin (terrestrial or extraterrestrial).

The correlation with the other tephtras identified in the karstic infillings of the region can be examined. The tephtra of the Devès du Reynaud aven [Debard and Pastre, 2008] is characterized by the presence of basaltic glass and its richness in biotite to which a peculiar provenance can be attributed. This composition is clearly different from the Chauvet cryptotephra. In the Marzal II (or Flahault), the basaltic and mantellic minerals reworked in set 5 [Debard and Pastre, 2008] can also be attributed to a phreatomagmatic emission of the Upper Pleistocene volcanism of Ardèche. The data obtained by biochronology suggest that this fall could belong to the recent or middle phase of Ardèche volcanism. The minerals of this infilling could originate from the same fall as the minerals of the Chauvet cave's cryptotephra or originate from another eruption like those of Vestide du Pal or Chambon. The tephtra identified in the Moula rockshelter corresponds to a beautiful in situ phreatomagmatic fall [Debard and Pastre, 2008, Pastre *et al.*, 1994]. Rich in basement minerals, it contains both peridotite and basaltic minerals as illustrated by the composition of its olivines varying between 43.97 and 48.87% MgO. Its initial dating around 40 ka [Pastre *et al.*, 1994] was contested according to a TL dating of 72 ± 12 ka [Sanzelle *et al.*, 2000]. It is indeed worth noticing that this older age is hardly reconcilable with the sedimentological and paleoclimatic data suggesting a periglacial context rather than a temperate one such as the Saint-Germain 2 episode [Debard and Pastre, 2008]. In the hypothesis of an age close to 40 ka, its correlation with the phreatomagmatic eruption of the Ray-Pic maar is possible. This hypothesis is in any

case compatible with the composition of the products identified in these two entities.

7. Conclusion

The cryptotephra from the Chauvet-Pont d'Arc cave provides a new example of tephra fall from the Upper Pleistocene volcanism of the Bas-Vivarais, trapped in the karst of this area. Its possible attribution to the phreatomagmatic eruption of the Ray-Pic maar could be supported by the mineralogical and geochemical compositions of its products containing mantellic and basaltic minerals. It is also in agreement with the recent $^{40}\text{Ar}/^{39}\text{Ar}$ datings of the Ardèche volcanism [Sasco, 2015, Sasco *et al.*, 2017], giving an age subcontemporaneous with the beds containing the charcoals well dated by ^{14}C [Quiles *et al.*, 2016] and contemporaneous with the Aurignacian occupation and art, that is slightly older than or close to 36 ka. In the other hand, the cryptotephra could also originate from the Vestide du Pal maar, as it also contains peculiar minerals whose origin is still unknown and which were also found in sedimentary infilling of the Chauvet cave. The discovery of this slightly reworked tephra in this remarkable Paleolithic cave provides an additional evidence of the contemporaneity of the Ardèche volcanism with the Aurignacian occupation of this region. Our findings are however not the bulletproof that will legitimate the hypothesis according to which the spray-shape signs of the cave can be ascribed to the first depiction of a volcanic eruption [Nomade *et al.*, 2016], but they surely reinforced the hypothesis that Aurignacians have been the witnesses of volcanic activity in the Ardèche region and that volcanic derived material did fall in the vicinity of Chauvet-Pont d'Arc cave about 36 ka ago.

Acknowledgments

We thank the French Culture Ministry and the ARPA (Association de Recherche Paléoécologique en Archéologie) for their financial support. We are especially grateful to Jean-Michel Geneste, former director of the scientific team of the Chauvet cave who has agreed to fund some of the analyses and to Carole Fritz (CNRS), actual research manager of the Chauvet team. Warm thanks are addressed to Jane Reilly for her help in the first English translation of this paper. We are grateful to Hélène Paquet for her review of this paper.

References

- Berger, E. (1973). *Le volcanisme récent de l'Ardèche*. PhD thesis, Thèse, Univ. Orsay. 402 p.
- Berger, E. (1981). Enclaves ultramafiques, mégacrystaux et leurs basaltes-hôtes en contexte océanique (Pacifique sud) et continental (Massif central français). Thèse d'Etat, Univ. Paris-sud, Orsay, 470 p.
- Berger, E. (2007). *Les jeunes volcans d'Ardèche*. Sud-Ouest éditions. 2879018293, 187 p.
- Berger, E., Brousse, R., and Maury, R. C. (1975). Age récent ($11\,770 \pm 270$ ans BP) des dernières éruptions du Vivarais. *C. R. Acad. Sci., Paris, Ser. D*, 280, 419–422.
- Berger, E. and Forette, M. C. (1975). Réactions des basaltes alcalins des Causses, du Vivarais et du Velay (France) avec les xénocrystaux et les minéraux des enclaves homéogènes. *Bull. Soc. Fr. Mineral. Cristal.*, 98, 366–373.
- Chauvel, C. and Jahn, B. M. (1984). Nd–St isotope and REE geochemistry of alkali basalts from the Massif Central, France. *Geochim., Cosmochim. Acta*, 48, 93–110.
- Clottes, J. (2001). *La Grotte Chauvet. L'art des origines*. Seuil Edition, Paris. 224 pp.
- Clottes, J., Chauvet, J.-M., Brunel-Deschamps, É., Hillaire, Ch., Daugas, J.-P., Arnold, M., Cachier, H., Évin, J., Fortin, Ph., Oberlin, C., Tisnerat, N., and Valladas, H. (1995). Les peintures préhistoriques de la Grotte Chauvet-Pont d'Arc (Ardèche, France) : datations directes et indirectes par la méthode du radiocarbone. *C. R. Acad. Sci., Paris, Ser. IIa*, 320, 1133–1140.
- Clottes, J. and Geneste, J.-M. (2007). La Grotte Chauvet, dix ans après. *Dossiers d'archéologie*, 324, 10–19. special issue “Grottes ornées en France”.
- Cuzange, M. T., Delqué-Kolic, E., Goslar, R., Meiort Grootes, P., Higham, T., Kaltnecker, E., Nadau, M.-J., Oberlin, C., Paterne, M., van der Plicht, J., Bronk Ramsey, Ch., Valladas, H., Clottes, J., and Geneste, J.-M. (2007). Radiocarbon inter-comparaison program for Chauvet Cave. *Radiocarbon*, 49(2), 339–347.
- Debard, É. (1988). Le Quaternaire du Bas-Vivarais d'après l'étude des remplissages d'avens, de porches de grottes et d'abris sous-roche. Dynamique sédimentaire, paléoclimatologie et chronologie. Thèse d'Etat. *Doc. Labo. géol. Lyon*, 103. 317 p.

- Debard, É. (1997). Les remplissages karstiques du Bas-Vivarais : karstogénèse, sédimentogénèse et archéologie. *Quaternaire*, 8, 305–317.
- Debard, É., Ferrier, C., and Kervazo, B. (2016). Grotte Chauvet-Pont d'Arc (Ardèche) : évolution morphosédimentaire de l'entrée. Implication sur les occupations et sur la conservation des vestiges. *Quaternaire*, 27(1), 3–14.
- Debard, É. and Pastre, J. F. (2008). Nouvelles données sur les téphras pléistocènes piégés dans les remplissages karstiques ardéchois (S-E France). *Quaternaire*, 19(2), 107–116.
- Delannoy, J.-J. and Geneste, J.-M., editors (2020). *Monographie de la Grotte Chauvet-Pont d'Arc. Vol. 1, Atlas de la Grotte Chauvet-Pont d'Arc*, volume 111-1 of *Documents d'archéologie française*. Éditions de la Maison des Sciences de l'homme.
- Downes, H. (1987). Relationship between geochemistry and textural type in spinel peridotite xenoliths, Massif Central, France. *Earth Planet. Sci. Lett.*, 82, 121–135.
- Faujas de Saint-Fond, B. (1778). *Recherches sur les volcans éteints du Vivarais et du Velay*. Joseph Cuchet Edition, Grenoble.
- Ferrier, C., Debard, É., Kervazo, B., Brodard, A., Guibert, P., Baffier, D., Feruglio, V., Gély, B., Geneste, J.-M., and Maksud, F. (2014). Les parois chauffées de la grotte Chauvet-Pont d'Arc (Ardèche, France) : caractérisation et chronologie. *Paleo*, 25, 59–78.
- Gély, B., Boche, E., and Maksud, F. (2012). Rapport triennal 2010–2012. In *Études pluridisciplinaires à la grotte Chauvet-Pont d'Arc (Ardèche). Rapports d'activités 2010–2012*, pages 215–242. (coordinateur J.-M. Geneste) (unpublished report, DRAC Auvergne-Rhône-Alpes).
- Gély, B. and Maksud, F. (2009). Sondage GE-1 dans la galerie d'entrée - Travaux 2009 et synthèse des résultats 2006–2009. In *Études pluridisciplinaires à la grotte Chauvet-Pont d'Arc (Ardèche). Rapports d'activités triennaux 2007–2009 et Perspectives 2010–2012*, pages 211–233. (coordinateur J.-M. Geneste) (unpublished report, DRAC Auvergne-Rhône-Alpes).
- Genty, D. (2008). Datations et analyses isotopiques des stalagmites de la grotte Chauvet du Secteur Rouge. In Geneste, J. M., editor, *La grotte Chauvet à Vallon Pont d'Arc (Ardèche), Rapports d'activités 2008*, pages 155–160. (unpublished report, DRAC Auvergne-Rhône-Alpes).
- Genty, D., Blamart, D., and Ghaleb, B. (2005). Apport des stalagmites pour l'étude de la grotte Chauvet: datations absolues U/Th (TIMS) et reconstitution paléoclimatique par les isotopes stables de la calcite. In : Geneste, J.M. (dir.), *La grotte Chauvet à Vallon-Pont d'Arc : un bilan des recherches pluridisciplinaires. Actes de la séance de la Société préhistorique française, Lyon, 2003*. *Bull. Soc. Prehist. Fr.*, 102(1), 45–62.
- Guérin, G. (1983). La thermoluminescence des plagioclases. Méthode de datation du volcanisme, applications au domaine volcanique français : Chaîne des Puys, Mont-Dore et Cézallier, Bas Vivarais. Thèse d'Etat. Université Pierre et Marie Curie, 258 p.
- Guérin, G. and Gillot, P. Y. (2007). Nouveaux éléments de chronologie du volcanisme pléistocène du Bas-Vivarais, (Ardèche, France) par thermoluminescence. *C. R. Geosci.*, 339, 40–49.
- Guibert, P., Brodard, A., Quiles, A., Geneste, J.-M., Baffier, D., Debard, É., and Ferrier, C. (2015). When were the walls of the Chauvet Cave heated? A chronological approach by thermoluminescence. *Quat. Geochronol.*, 29, 36–47.
- Nomade, S., Genty, D., Sascio, R., Scao, V., Feruglio, V., Baffier, D., Guillou, H., Bourdier, C., Valladas, H., Reigner, E., Debard, É., Pastre, J. F., and Geneste, J. M. (2016). A 36,000-year-old volcanic eruption depicted in the Chauvet-Pont d'Arc Cave (Ardèche, France)? *PLoS One*, 11(1), article no. e0146621.
- Pastre, J. F., Debard, É., and Chennaoui, K. (1994). Un téphra-repère du volcanisme phréatomagmatique du Vivarais dans la séquence Pléistocène supérieur de l'abri Moula (Soyons, Ardèche, France). *C. R. Acad. Sci. Paris, Ser. II*, 319, 937–943.
- Quiles, A., Valladas, H., Bocherens, H., Delqué-Kolic, E., Kaltnecker, E., van der Plicht, J., Delannoy, J. J., Feruglio, V., Fritz, C., Monney, J., Philippe, M., Tosello, G., Clottes, J., and Geneste, J. M. (2016). A high-precision chronological model for the decorated Upper Paleolithic cave of Chauvet-Pont d'Arc, Ardèche, France. *Proc. Natl Acad. Sci. USA*, 113(17), 4670–4675.
- Rochette, P., Bertrand, H., Braun, C., and Berger, E. (1993). La province volcanique Pléistocène Supérieur du Bas-Vivarais (Ardèche, France) : propagation de fentes crustales en échelons? *C. R. Acad. Sci. Paris, Ser. II*, 316, 913–920.

- Sadier, B., Delannoy, J. J., Benedetti, L., Bourlès, D. L., Jaillet, S., Geneste, J. M., Lebatard, A. E., and Arnold, M. (2012). Further constraints on the Chauvet cave artwork elaboration. *Proc. Natl Acad. Sci. USA*, 109(21), 8002–8006.
- Sanzelle, S., Pilleyre, T., Montert, M., Fain, J., Miallier, D., Camus, G., de Goër, H., and Defleur, A. (2000). Datation par thermoluminescence : étude d'une corrélation possible entre le maar de la Vestide-du-Pal et un niveau de téphra de la Baume-Moula-Guercy (Ardèche, France). *C. R. Acad. Sci. Paris, Ser. II*, 330, 541–546.
- Sasco, R. (2015). Développement d'un outil chronostratigraphique pour les archives climatiques : datations absolues (K/Ar, $^{40}\text{Ar}/^{39}\text{Ar}$) et paléomagnétisme appliqués aux laves. Thèse de doctorat, Université Paris-sud.
- Sasco, R., Guillou, H., Nomade, S., Scao, V., Maury, R. C., Kissel, C., and Wandres, C. (2017). $^{40}\text{Ar}/^{39}\text{Ar}$ and unspiked ^{40}K – ^{40}Ar dating of upper Pleistocene volcanic activity in the Bas-Vivarais (Ardèche, France). *J. Volcanol. Geotherm. Res.*, 341, 301–314.
- Valladas, H., Tisnerat-Laborde, N., Cachier, H., Kaltnecker, E., Arnold, M., Oberlin, C., and Evin, J. (2005). Bilan des datations carbone 14 effectuées sur des charbons de bois de la grotte Chauvet. *Bull. Soc. Préhist. Fr.*, 102(1), 109–113.