



ELSEVIER

Contents lists available at SciVerse ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Probability Theory

The shape of the $(2 + 1)$ D SOS surface above a wall [☆]*La forme de l'interface SOS $(2 + 1)$ -dimensionnelle au-dessus d'un mur*Pietro Caputo ^a, Eyal Lubetzky ^b, Fabio Martinelli ^a, Allan Sly ^c, Fabio Lucio Toninelli ^d^a Università Roma Tre, Largo S. Murialdo 1, 00146 Roma, Italy^b Microsoft Research, One Microsoft Way, Redmond, WA 98052, USA^c UC Berkeley, Berkeley, CA 94720, USA^d CNRS and ENS Lyon, laboratoire de physique, 46 allée d'Italie, 69364 Lyon, France

ARTICLE INFO

Article history:

Received 15 July 2012

Accepted 18 July 2012

Available online 1 August 2012

Presented by the Editorial Board

ABSTRACT

We give a full description for the shape of the classical $(2 + 1)$ D Solid-On-Solid model above a wall, introduced by Temperley (1952) [14]. On an $L \times L$ box at a large inverse-temperature β the height of most sites concentrates on a single level $h = \lfloor \frac{1}{4\beta} \log L \rfloor$ for most values of L . For a sequence of diverging boxes the ensemble of level lines of heights $(h, h - 1, \dots)$ has a scaling limit in Hausdorff distance iff the fractional parts of $\frac{1}{4\beta} \log L$ converge to a noncritical value. The scaling limit is explicitly given by nested distinct loops formed via translates of Wulff shapes. Finally, the h -level lines feature $L^{1/3+o(1)}$ fluctuations from the side boundaries.

© 2012 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Nous donnons une description complète de la forme typique de l'interface SOS $(2 + 1)$ -dimensionnelle au-dessus d'un mur, introduite par Temperley (1952) [14]. Dans une boîte $L \times L$ à basse température $T = 1/\beta$, la hauteur de la plupart des sommets se concentre au niveau $h = \lfloor \frac{1}{4\beta} \log L \rfloor$, pour la plupart des valeurs de L . Pour une suite croissante de boîtes, l'ensemble de lignes de niveau de hauteur $(h, h - 1, \dots)$ admet une limite au sens de la distance de Hausdorff ssi la partie fractionnaire de $\frac{1}{4\beta} \log L$ converge à une valeur non critique. La limite d'échelle est donnée explicitement par des boucles imbriquées, formées par des translations de formes de Wulff. Enfin, la distance entre le bord de la boîte et la ligne de niveau h a des fluctuations $L^{1/3+o(1)}$.

© 2012 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

The $(2 + 1)$ -dimensional *Solid-On-Solid* model is a crystal surface model whose definition goes back to Temperley [14] in 1952 (also known as the Onsager–Temperley sheet). At low temperatures, the model approximates the interface between the plus and minus phases in the 3D Ising model. See, e.g., [7–9] for notable previous works on this model.

[☆] This work was supported by the European Research Council through the “Advanced Grant” PIRELSS 228032.

E-mail addresses: caputo@mat.uniroma3.it (P. Caputo), eyal@microsoft.com (E. Lubetzky), martin@mat.uniroma3.it (F. Martinelli), sly@stat.berkeley.edu (A. Sly), fabio-lucio.toninelli@ens-lyon.fr (F.L. Toninelli).

The configuration space of the model on a finite box $\Lambda \subset \mathbb{Z}^2$ with a floor (wall) at 0 and zero boundary conditions is the set Ω_Λ of all height functions η on \mathbb{Z}^2 such that $\Lambda \ni x \mapsto \eta_x \in \mathbb{Z}_+$ whereas $\eta_x = 0$ for all $x \notin \Lambda$. The probability of $\eta \in \Omega_\Lambda$ is given by the Gibbs distribution

$$\pi_\Lambda(\eta) = \frac{1}{Z_\Lambda} \exp\left(-\beta \sum_{x \sim y} |\eta_x - \eta_y|\right), \quad (1)$$

where $\beta > 0$ is the inverse-temperature, $x \sim y$ denotes a nearest-neighbor bond in the lattice \mathbb{Z}^2 and the normalizing constant Z_Λ is the partition-function.

Despite progress on understanding the typical height of the surface (notably the work of Bricmont, El Mellouki and Fröhlich [2] in 1986 and the recent companion paper of the authors [3]), little was known on its actual 3D shape. The fundamental problem is the following:

Question 1.1. Consider the ensemble of all level lines of the low temperature $(2 + 1)_D$ SOS on an $L \times L$ box, rescaled to the unit square:

- (i) Do these jointly converge to a scaling limit as $L \rightarrow \infty$, e.g., in Hausdorff distance?
- (ii) If so, can the limit be explicitly described?
- (iii) For finite large L , what are the fluctuations of the level lines around their limit?

In this work we fully resolve parts (i) and (ii) and partially answer part (iii). En route, we also establish that for most values of L the surface height concentrates on the single level $\lfloor \frac{1}{4\beta} \log L \rfloor$.

1.1. Main results

Our first result addresses the distribution of the surface height.

Theorem 1 (Height concentration). Fix $\beta > 0$ to be sufficiently large and define

$$H(L) = (1/4\beta) \log L. \quad (2)$$

Let $E_h = \{\eta: \#\{x: \eta_x = h\} \geq \frac{9}{10} L^2\}$ be the event that at least $\frac{9}{10}$ of the sites are at height h . Then the SOS measure on the $L \times L$ box $\Lambda \subset \mathbb{Z}^2$ with inverse-temperature β satisfies

$$\lim_{L \rightarrow \infty} \pi_\Lambda(E_{\lfloor H \rfloor - 1} \cup E_{\lfloor H \rfloor}) = 1. \quad (3)$$

Furthermore, the typical height of the configuration is governed by $\alpha(L) = H - \lfloor H \rfloor$ as follows: Let Λ_n be a diverging sequence of boxes with side-lengths L_n . There exists $0 < \alpha_c(\beta) < 1$ so that

- (i) If $\liminf_{n \rightarrow \infty} \alpha(L_n) > \alpha_c$ then $\lim_{n \rightarrow \infty} \pi_{\Lambda_n}(E_{\lfloor H \rfloor}) = 1$.
- (ii) If $\limsup_{n \rightarrow \infty} \alpha(L_n) < \alpha_c$ then $\lim_{n \rightarrow \infty} \pi_{\Lambda_n}(E_{\lfloor H \rfloor - 1}) = 1$.

Remark. The constant $\frac{9}{10}$ in the definition of E_h can be replaced by $1 - \varepsilon$ for any arbitrarily small $\varepsilon > 0$ provided that β is chosen large enough. Moreover, the critical fractional value $\alpha_c(\beta)$ satisfies $\alpha_c = (1 + \varepsilon\beta) \frac{\log(4\beta)}{4\beta}$ where $\varepsilon\beta \rightarrow 0$ as $\beta \rightarrow \infty$. Therefore, for large enough (fixed) β most values of $\alpha(L)$ will yield $\pi_\Lambda(E_{\lfloor H \rfloor}) \rightarrow 1$.

We now address the scaling limit of the ensemble of level lines, formally defined next.

Definition 1.2. Let η be an SOS configuration on a box $\Lambda \in \mathbb{Z}^2$. The h -level lines ($h \geq 1$) are a collection of disjoint self-avoiding loops in the dual \mathbb{Z}^{2*} formed as follows. Let \mathcal{E}_h be the set of all bonds $e' \in \mathbb{Z}^{2*}$ whose dual edge $e = (x, y)$ is such that $\eta_x \geq h$ whereas $\eta_y < h$, or vice versa. Due to the zero boundary conditions on η , the bonds of \mathcal{E}_h can then be uniquely partitioned into a finite number of edge-disjoint simple loops as follows. Whenever four bonds in \mathcal{E}_h meet at a vertex we separate them along the NW-oriented diagonal going through the intersection. The final collection of loops comprises the h -level lines.

The next theorem gives a necessary and sufficient condition for the existence of a scaling limit to the full ensemble of level lines, and furthermore provides an explicit description for the limit.

Theorem 2 (Shape Theorem). Fix $\beta > 0$ to be sufficiently large and let L_n be a diverging sequence of side-lengths. Set $H_n = (1/4\beta) \log L_n$ and $\alpha_n = H_n - \lfloor H_n \rfloor$. For an SOS surface on a box Λ_n of side-length L_n , let $(\mathcal{L}_0^{(n)}, \mathcal{L}_1^{(n)}, \dots)$ be the collections of loops whose area is at least $\log^2 L$ belonging to the level lines of heights $(\lfloor H_n \rfloor, \lfloor H_n \rfloor - 1, \dots)$, respectively. Then:

- (a) With high probability, the level lines of every height $h > \lfloor H \rfloor$ consist of microscopic loops, each of which spans an area of at most $\log^2 L$. In addition, for each $i \geq 1$ w.h.p. $\mathcal{L}_i^{(n)}$ consists of exactly one loop whereas $\mathcal{L}_0^{(n)}$ is w.h.p. either empty or contains a single loop.
- (b) The rescaled loop ensemble $\frac{1}{L_n}(\mathcal{L}_0^{(n)}, \mathcal{L}_1^{(n)}, \dots)$ converges almost surely to a limit $(\mathcal{W}_0, \mathcal{W}_1, \dots)$ in Hausdorff distance iff $\lim_{n \rightarrow \infty} \alpha_n$ exists and differs from α_c , the critical fractional value given in Theorem 1. More precisely, in this case almost surely $\limsup_{n \rightarrow \infty} \sup_i d_{\mathcal{H}}(\frac{1}{L_n} \mathcal{L}_i^{(n)}, \mathcal{W}_i) = 0$.
- (c) The scaling limit $(\mathcal{W}_0, \mathcal{W}_1, \dots)$ is explicitly given as a nested sequence of distinct loops invariant under $\pi/4$ rotations, each of which is constructed as follows. Let $\alpha_* = \lim_{n \rightarrow \infty} \alpha_n$.
 - If $\alpha_* > \alpha_c$ then for any $i \geq 0$ the loop \mathcal{W}_i is the boundary of the union of all possible translates within the unit square of $\mathcal{W}_*(r_i)$ with an explicit $0 < r_i(\alpha_*) < 1$, where $\mathcal{W}_*(r)$ is the r -dilation of a convex smooth shape (the area 1 SOS Wulff shape).
 - If $\alpha_* < \alpha_c$ then \mathcal{W}_0 is empty and the remaining loops $\{\mathcal{W}_i\}_{i \geq 1}$ are constructed as above.

We remark that for large enough β each \mathcal{W}_i has an overlap of length at least $1/2$ with each side-boundary and positive distance from the corners. The fluctuations of the loop $\mathcal{L}_0^{(n)}$ (the plateau at level $\lfloor H \rfloor$) from its limit $L_n \times \mathcal{W}_0$ along the side boundaries are now addressed.

Theorem 3 (Cube-root fluctuations). In the setting of Theorem 2 suppose $\liminf_{n \rightarrow \infty} \alpha_n > \alpha_c$. Consider $\mathcal{L}_0^{(n)}$, the collection of macroscopic loops belonging to the level line at height $\lfloor H_n \rfloor$ of the SOS surface. Then the maximal vertical fluctuation of $\mathcal{L}_0^{(n)}$ from the boundary interval $[(\frac{1}{4}L_n, 0), (\frac{3}{4}L_n, 0)]$ is w.h.p. of order $L_n^{1/3+o(1)}$. More precisely, let $\rho(x) = \min\{y : (x, y) \in \mathcal{L}_0^{(n)}\}$ be the vertical fluctuation of $\mathcal{L}_0^{(n)}$ from the bottom boundary at coordinate x . Then for any $\varepsilon > 0$ and large enough n ,

$$\pi_{\Lambda_n} \left(L_n^{1/3-\varepsilon} < \sup_{\frac{1}{4}L_n \leq x \leq \frac{3}{4}L_n} \rho(x) < L_n^{1/3+\varepsilon} \right) > 1 - e^{-L_n^\varepsilon}.$$

As a direct corollary one can deduce an upper bound on the fluctuations of all level lines.

Corollary 4 (Cascade of fluctuation exponents). In the same setting of Theorem 3, let $\rho(i, x)$ be the vertical fluctuation of $\mathcal{L}_i^{(n)}$ from the bottom boundary at coordinate x . Let $0 < \xi < 1$ and let $i = \lfloor \xi H_n \rfloor$. Then for any $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} \pi_{\Lambda_n} \left(\sup_{\frac{1}{4}L_n \leq x \leq \frac{3}{4}L_n} \rho(i, x) > L_n^{\frac{1-\xi}{3}+\varepsilon} \right) = 0.$$

Proof. Choose $\ell = (1 + o(1))L^{1-\xi}$ such that $\lfloor H(\ell) \rfloor = \lfloor H_n \rfloor - i$ while $\alpha(\ell) > 2\alpha_c$ (recalling that α_c is arbitrarily small for large β). Fix a coordinate $\frac{1}{4}L_n \leq x \leq \frac{3}{4}L_n$ on the bottom boundary and consider the box $B = [x - \frac{\ell}{2}, x + \frac{\ell}{2}] \times [0, \ell]$. By imposing zero boundary conditions on the box B we obtain a surface that is clearly stochastically dominated by the original one. By the choice of $\alpha(\ell)$, Theorem 1 ensures w.h.p. the existence of a macroscopic loop \mathcal{L}'_0 at height $\lfloor H(\ell) \rfloor$ for the new surface in B . Theorem 3 guarantees that the vertical fluctuation of \mathcal{L}'_0 at coordinate x is at most $\ell^{1/3+\varepsilon}$ except with probability $e^{-\ell^\varepsilon}$. A union bound extends this bound to the entire interval $x \in [\frac{1}{4}L_n, \frac{3}{4}L_n]$. Hence, w.h.p. the macroscopic loop $\mathcal{L}_i^{(n)}$ of the original surface must be within $\ell^{1/3+\varepsilon} = (1 + o(1))L^{(1-\xi)/3+\varepsilon}$ from the bottom interval. \square

Detailed proofs of Theorems 1–3 will be given in a forthcoming paper.

2. Related work and open problems

2.1. Limiting shape

In the two papers [12,13] the authors studied the limiting shape of the low temperature 2D Ising with minus boundary under a prescribed small positive external field, proportional to the inverse of the side-length L . The behavior of the droplet of plus spins in this model is qualitatively similar to the behavior of the top loop \mathcal{L}_0 in our case. Here, instead of an external field, it is the subtle entropic repulsion phenomenon which induces the surface to rise to level $\lfloor H \rfloor$ and creates a macroscopic loop \mathcal{L}_0 . In line with this connection, the description of the limiting shape of the plus droplet in the aforementioned works exactly coincides with our limit \mathcal{W}_0 (with a different Wulff shape).

An important difference between the two models is of course the fact that in our case there exist $\lfloor H \rfloor \geq c \log L$ levels (rather than just one), which are interacting in two nontrivial ways. First, they are dependent as they are nested by definition. Second, they can weakly either attract or repel one another depending on the local geometry and height. Moreover, the box boundary itself can attract or repel the level lines. A prerequisite to proving Theorem 2 is to overcome these “pinning” issues. As a side note we remark that at times such pinning issues have been overlooked in the relevant literature.

As for the fluctuations of the plus droplet from its limiting shape, it was argued in [12] that these should be normal (i.e., \sqrt{L}). However, due to the analogy mentioned above between the models, it follows from our proof of Theorem 3 that these fluctuations are in fact $L^{1/3+o(1)}$.

2.2. Cube root fluctuations

It turns out that governing the behavior of the top level loop \mathcal{L}_0 is a version of the usual measure associated with Ising-type contours (see, e.g., [5]) crucially tilted by a factor of the form $\exp((\lambda/L)\text{area}(\mathcal{L}_0))$ for some fixed $\lambda > 0$. Modulo this fact, whose proof is quite delicate, the $L^{1/3}$ fluctuations can be recovered by analyzing the behavior of the loop along mesoscopic boxes of dimensions $L^{2/3+o(1)} \times L^{1/3+o(1)}$, in the spirit of the approach of [5].

There is a rich literature of contour models featuring similar distributions and cube root fluctuations. In some of these works (e.g., [1,6,10,15]) the area term appears due to an externally imposed constraint (by conditioning and/or adding an external field). In others, modeling ordered random walks (e.g., [4,11] to name a few), the area effect on a contour is due to the contours preceding it, and the exact solvability of the model (e.g., via determinantal representations) plays an essential role in the analysis.

We stress that the area term driving the $L^{1/3}$ fluctuations in our setting arises naturally due to entropic repulsion. By the lack of exact solvability for the $(2+1)$ D SOS we must resort to cluster expansion techniques and contour analysis as in the framework of [5]. However, in the mesoscopic scale (required to establish the cube root fluctuations) this framework is particularly prone to the aforementioned pinning issues of level lines (to those above and below them, as well as the boundary), forming a major technical obstacle for the analysis.

2.3. Open problems

We conclude by mentioning two problems that remain unaddressed by our results. First, establish the exponents for the fluctuations of all intermediate level lines from the side boundaries. We believe the upper bound in Corollary 4 features the correct cascade of exponents. Second, find the correct fluctuation exponent of the level lines $\{\mathcal{L}_i^{(n)}\}$ around the curved part of their limiting shapes $\{L \times \mathcal{W}_i\}$.

References

- [1] K.S. Alexander, Cube-root boundary fluctuations for droplets in random cluster models, *Comm. Math. Phys.* 224 (3) (2001) 733–781.
- [2] J. Bricmont, A. El Mellouki, J. Fröhlich, Random surfaces in statistical mechanics: roughening, rounding, wetting, ..., *J. Stat. Phys.* 42 (5–6) (1986) 743–798.
- [3] P. Caputo, E. Lubetzky, F. Martinelli, A. Sly, F.L. Toninelli, Dynamics of $2+1$ dimensional SOS surfaces above a wall: slow mixing induced by entropic repulsion, preprint. Available at arXiv:1205.6884, 2012.
- [4] I. Corwin, A. Hammond, Brownian Gibbs property for Airy line ensembles, preprint. Available at arXiv:1108.2291, 2011.
- [5] R. Dobrushin, R. Kotecký, S. Shlosman, Wulff Construction. A Global Shape from Local Interaction, *Transl. Math. Monogr.*, vol. 104, American Mathematical Society, Providence, RI, 1992.
- [6] P.L. Ferrari, H. Spohn, Constrained Brownian motion: fluctuations away from circular and parabolic barriers, *Ann. Probab.* 33 (4) (2005) 1302–1325.
- [7] J. Fröhlich, T. Spencer, Kosterlitz–Thouless transition in the two-dimensional plane rotator and Coulomb gas, *Phys. Rev. Lett.* 46 (15) (1981) 1006–1009.
- [8] J. Fröhlich, T. Spencer, The Kosterlitz–Thouless transition in two-dimensional abelian spin systems and the Coulomb gas, *Comm. Math. Phys.* 81 (4) (1981) 527–602.
- [9] J. Fröhlich, T. Spencer, The Berezinskiĭ–Kosterlitz–Thouless transition (energy–entropy arguments and renormalization in defect gases), scaling and self-similarity in physics, *Prog. Phys.* 7 (1983) 29–138.
- [10] O. Hryniv, Y. Velenik, Universality of critical behaviour in a class of recurrent random walks, *Probab. Theory Related Fields* 130 (2) (2004) 222–258.
- [11] K. Johansson, Discrete polynuclear growth and determinantal processes, *Comm. Math. Phys.* 242 (1–2) (2003) 277–329.
- [12] R.H. Schonmann, S.B. Shlosman, Complete analyticity for 2D Ising completed, *Comm. Math. Phys.* 170 (2) (1995) 453–482.
- [13] R.H. Schonmann, S.B. Shlosman, Constrained variational problem with applications to the Ising model, *J. Stat. Phys.* 83 (5–6) (1996) 867–905.
- [14] H.N.V. Temperley, Statistical mechanics and the partition of numbers. II. The form of crystal surfaces, *Proc. Cambridge Philos. Soc.* 48 (1952) 683–697.
- [15] Y. Velenik, Entropic repulsion of an interface in an external field, *Probab. Theory Related Fields* 129 (1) (2004) 83–112.