



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Group theory

Projective modules in the intersection cohomology of Deligne–Lusztig varieties



Modules projectifs dans la cohomologie d'intersection des variétés de Deligne–Lusztig

Olivier Dudas^a, Gunter Malle^{b,1}^a Université Paris-Diderot, UFR de mathématiques, bâtiment Sophie-Germain, 5, rue Thomas-Mann, 75205 Paris cedex 13, France^b FB Mathematik, TU Kaiserslautern, Postfach 3049, 67653 Kaiserslautern, Germany

ARTICLE INFO

Article history:

Received 13 February 2014

Accepted 7 March 2014

Available online 13 April 2014

Presented by the Editorial Board

ABSTRACT

We formulate a strong positivity conjecture on characters afforded by the Alvis–Curtis dual of the intersection cohomology of Deligne–Lusztig varieties. This conjecture provides a powerful tool to determine decomposition numbers of unipotent ℓ -blocks of finite reductive groups.

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Nous formulons une conjecture de positivité sur le dual d'Alvis–Curtis du caractère obtenu à partir de la cohomologie d'intersection d'une variété de Deligne–Lusztig. Cette conjecture se révèle être un outil puissant pour déterminer les nombres de décompositions des ℓ -blocs unipotents des groupes réductifs finis.

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Deligne–Lusztig varieties

Let \mathbf{G} be a connected reductive linear algebraic group over an algebraically closed field of positive characteristic p , and $F : \mathbf{G} \rightarrow \mathbf{G}$ be a Steinberg endomorphism. There exists a positive integer δ such that F^δ defines a split \mathbb{F}_{q^δ} -structure on \mathbf{G} (with $q \in \mathbb{R}_+$), and we will choose δ minimal for this property. We set $G := \mathbf{G}^F$, the finite group of fixed points.

We fix a pair (\mathbf{T}, \mathbf{B}) consisting of a maximal torus contained in a Borel subgroup of \mathbf{G} , both of which are assumed to be F -stable. We denote by W the Weyl group of \mathbf{G} , and by S the set of simple reflections in W associated with \mathbf{B} . Then F^δ acts trivially on W .

Following Deligne and Lusztig, there exists a variety $X(w)$, for any $w \in W$, endowed with an action of $G \times \langle F^\delta \rangle$, whose ℓ -adic cohomology gives rise to the so-called *unipotent* representations of G .

E-mail addresses: dudas@math.jussieu.fr (O. Dudas), malle@mathematik.uni-kl.de (G. Malle).

¹ The author gratefully acknowledges financial support by ERC Advanced Grant 291512.

1.1. Basic sets for finite reductive groups

Let ℓ be a prime number and (K, \mathcal{O}, k) be an ℓ -modular system. We assume that it is large enough for all the finite groups encountered. Furthermore, since we will be working with ℓ -adic cohomology, we will assume throughout this note that K is a finite extension of \mathbb{Q}_ℓ .

Let H be a finite group. Representations of H will always be assumed to be finite-dimensional. Recall that every projective kH -module lifts to a representation of H over K . The character afforded by such a representation will be referred to as a *projective character*. Integral linear combinations of projective characters will be called *virtual projective characters*.

Throughout this paper, we shall make the following assumptions on ℓ :

- $\ell \neq p$ (non-defining characteristic),
- ℓ is good for \mathbf{G} and $\ell \nmid |(Z(\mathbf{G})/Z(\mathbf{G})^\circ)^F|$.

In this situation, the unipotent characters lying in a given unipotent ℓ -block of G form a *basic set* of this block [6,5]. Consequently, the restriction of the decomposition matrix of the block to the unipotent characters is invertible. In particular, every (virtual) unipotent character is a virtual projective character, up to adding and removing some non-unipotent characters.

1.2. A positivity conjecture

Let D_G denote the Alvis–Curtis duality, with the convention that if ρ is a cuspidal unipotent character, then $D_G(\rho) = (-1)^{\text{rk}_F(\mathbf{G})} \rho$ where $\text{rk}_F(\mathbf{G}) = |S/F|$ is the F -semisimple rank of \mathbf{G} . For $w \in W$, we denote by $IH^i(\bar{X}(w))$ the i th intersection cohomology group of the Zariski closure $\bar{X}(w)$ of $X(w)$ in \mathbf{G}/\mathbf{B} , with coefficients in the trivial local system. We define Q_w to be the restriction to G of the virtual character

$$\sum_{i \in \mathbb{Z}} (-1)^i [D_G(IH^i(\bar{X}(w)))],$$

and by $Q_w[\lambda]$ the generalized λ -eigenspace of F^δ for $\lambda \in k^\times$. By [11, Thm. 3.8, 5.11.4 and 6.8.6] this character can be computed using almost characters together with values of characters of the Iwahori–Hecke algebra associated with W on the Kazhdan–Lusztig basis.

Lusztig proved [11, Prop. 6.9 and 6.10] that up to a global sign, Q_w (and even $Q_w[\lambda]$) is always a nonnegative linear combination of unipotent characters (note that the assumption on q can be removed by [2, Cor. 3.3.22]). The sign is given by the a -value $a(w)$ of the two-sided cell in which w lies.

Proposition 1.1 (Lusztig). *For all $w \in W$ and all $\lambda \in k^\times$, $(-1)^{a(w)} Q_w[\lambda]$ is a sum of unipotent characters.*

Unipotent characters are only the unipotent part of virtual projective characters in general. We conjecture that the modular analogue of Proposition 1.1 should hold in general, that is that Q_w is actually a proper projective character whenever ℓ is not too small.

Conjecture 1.2. *Under the assumption on ℓ in Section 1.1, for all $w \in W$ and all $\lambda \in k^\times$, $(-1)^{a(w)} Q_w[\lambda]$ is the unipotent part of a projective character.*

This conjecture has been checked on many decomposition matrices, including the unipotent blocks with cyclic defect groups for exceptional groups, and the matrices that are determined in [3,4]. We will give some examples in Section 3.

Example 1. The closure of the Deligne–Lusztig variety $X(w_0)$ associated with the longest element w_0 of W is smooth and equal to \mathbf{G}/\mathbf{B} . Therefore, its intersection cohomology $IH^\bullet(\bar{X}(w_0))$ consists of copies of the trivial representation in degrees of a given parity. Since $a(w_0) = \ell(w_0)$, then $(-1)^{a(w_0)} Q_{w_0}$ is a nonnegative multiple of the Steinberg character. On the other hand, the Steinberg character is the unipotent part of a unique projective indecomposable module (given by a summand of a Gelfand–Graev module) and therefore the conjecture holds for w_0 .

2. Applications

In this section, we explain how to deduce properties of decomposition matrices using Conjecture 1.2.

2.1. Families of simple unipotent modules

Following [11, § 5], we denote by \leq_{LR} the partial order on two-sided cells of W . Recall that to each cell Γ corresponds a two-sided ideal $I_{\leq \Gamma} := \text{span}_{\mathbb{Q}}\{C_w | w \leq_{LR} \Gamma\}$ of the group algebra $\mathbb{Q}W$, where $(C_w)_{w \in W}$ is the Kazhdan–Lusztig basis of the corresponding Hecke-algebra with parameter ν . Moreover, given $\chi \in \text{Irr } W$, there is a unique two-sided cell Γ_χ such that χ occurs in $I_{\leq \Gamma_\chi} / I_{< \Gamma_\chi}$. To each two-sided cell Γ one can attach a so-called family \mathcal{F}_Γ of unipotent characters. They are defined as the constituents of R_χ for various χ such that $\Gamma_\chi = \Gamma$. By [11, Thm. 6.17], they form a partition of the set of unipotent characters of G .

We can use the characters Q_w to define families of simple unipotent kG -modules. For each Q_w we choose a virtual projective kG -module \tilde{Q}_w whose character coincides with Q_w up to adding/removing non-unipotent characters. We denote by $\text{Irr}_{\leq \Gamma} kG$ (resp. $\text{Irr}_{< \Gamma} kG$) the set of simple unipotent kG -modules N such that a projective cover P_N occurs in the virtual module \tilde{Q}_w for some $w \leq_{LR} \Gamma$ (resp. $w <_{LR} \Gamma$). We define the family of simple unipotent modules associated with Γ as $\text{Irr}_\Gamma kG = \text{Irr}_{\leq \Gamma} kG \setminus \text{Irr}_{< \Gamma} kG$. Since the regular representation is uniform (see for example [1, Cor. 12.14]), then every indecomposable projective module lying in a unipotent block occurs in some \tilde{Q}_w ; therefore every simple unipotent module belongs to at least one family. However, it is unclear whether this family is unique in general.

It is conjectured that, for ℓ not too small, the ℓ -modular decomposition matrix of G depends only on the order of q modulo ℓ . The following proposition gives some evidence toward this conjecture as well as to Geck’s conjecture on the unitriangular shape of the decomposition matrix [7, Conj. 3.4].

Proposition 2.1. *Let b be a unipotent ℓ -block of G . Assume that*

- (i) *Conjecture 1.2 holds, and*
- (ii) *$|\text{Irr}_\Gamma bkG| = |\mathcal{F}_\Gamma \cap \text{Irr } b|$ for any two-sided cell Γ of W .*

Then the unipotent part of the decomposition matrix of b has the following shape:

$$D_{\text{uni}} = \begin{pmatrix} D_{\mathcal{F}_1} & 0 & \cdots & 0 \\ * & D_{\mathcal{F}_2} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ * & \cdots & * & D_{\mathcal{F}_r} \end{pmatrix}$$

where \mathcal{F}_i runs over the families and where each $D_{\mathcal{F}_i}$ is a square matrix of size $|\mathcal{F}_i \cap \text{Irr } b|$. Furthermore, the entries of D_{uni} are bounded above independently of q and ℓ .

Proof. Given any simple kG -module N in the block, there exists by (ii) a unique two-sided cell Γ such that $N \in \text{Irr}_\Gamma bkG$ and P_N occurs in some Q_w for $w \in \Gamma$ (note that $w \not\leq_{LR} \Gamma$, otherwise N would belong to a smaller family). Since the matrix of the Q_w ’s is block triangular with respect to families, the proposition follows from (i). \square

Remark 1. Assumption (ii) ensures that families define a partition of the set of simple unipotent kG -modules. This assumption will hold whenever each unipotent character in the block is a linear combination of the unipotent almost characters cut by the block. This condition can easily be checked on Fourier matrices; for example, it holds whenever $\ell | (q \pm 1)$ and G is exceptional.

2.2. Determining decomposition numbers

The bounds on the entries of the decomposition matrix given by the Q_w ’s are often small enough to determine some of the decomposition numbers. We illustrate this on the principal Φ_2 -block of the group $F_4(q)$.

Proposition 2.2. *Assume that $(q, 6) = 1$, $q \equiv -1 \pmod{\ell}$ and $\ell > 11$. If Conjecture 1.2 holds then*

$$\begin{array}{c|cccc} \phi_{9,10} & 1 & & & \\ \phi'_{2,16} & 1 & 1 & & \\ \phi''_{2,16} & 1 & \cdot & 1 & \\ B_{2,\varepsilon} & 2 & \cdot & \cdot & 1 \\ \phi_{1,24} & 3 & 2 & 2 & 4 & 1 \end{array}$$

is a submatrix of the ℓ -modular decomposition matrix of $F_4(q)$. (Here, “ \cdot ” denotes zero entries.)

Proof. By [10], the above matrix is a submatrix of the decomposition matrix of $F_4(q)$, but with undetermined entries f, g, h, i, j in place of the entries larger than 1, which satisfy $f, h, i \geq 2$, $g \geq 4f - 5$ and $j \geq 4$. Let b be the block idempotent associated with the principal ℓ -block of $F_4(q)$. With $w = s_2s_3s_2s_3s_4s_3s_2s_1s_3s_2s_4s_3s_2s_1$ we have:

$$bQ_w = 32\phi_{9,10} + 72\phi_{2,16}'' + 72\phi_{2,16}' + 72B_2, \varepsilon + 288\phi_{1,24}.$$

If we denote by P_1, \dots, P_5 the unipotent part of the characters of the PIMs corresponding to the last five columns of the decomposition matrix of $F_4(q)$, we can decompose Q_w as

$$bQ_w = 32P_1 + 40P_2 + 40P_3 + (72 - 32f)P_4 + (288 - 32g - 40h - 40i - (70 - 32f)j)P_5.$$

By [Conjecture 1.2](#), the multiplicity of each P_k should be nonnegative, that is $72 - 32f \geq 0$ and $288 - 32g - 40h - 40i - (70 - 32f)j \geq 0$. Since $f \geq 2$ the first relation forces $f = 2$ (and therefore $g \geq 3$). The second becomes $288 - 32g - 40h - 40i - 8j \geq 0$. Since $288 = 32 \times 3 + 40 \times 2 + 40 \times 2 + 8 \times 4$ is the minimal value that the expression $32g + 40h + 40i + 8j$ can take, we deduce that $h = i = 2, g = 3$ and $j = 4$. \square

3. Further examples

3.1. A cuspidal module in the unitary group

We give here a non-trivial example for $SU_n(q)$ where [Conjecture 1.2](#) holds. The key point is to find a formula for the corresponding Kazhdan–Lusztig basis element in terms of well-identified elements of W . This is done using the geometric description of Kazhdan–Lusztig polynomials. The approach given below could be adapted to other groups, even when the corresponding Schubert variety is no longer smooth, using Bott–Samelson varieties instead.

Recall that the set of unipotent characters of $SU_n(q)$ is parameterized by partitions of n . Given such a partition λ , we denote by ρ_λ (resp. χ_λ) the corresponding unipotent character (resp. character of \mathfrak{S}_n), with the convention that ρ_{1^n} is the Steinberg character.

Proposition 3.1. *Let $\ell > n$ be a prime dividing $q + 1$. Then [Conjecture 1.2](#) holds for $SU_n(q)$ and $w = s_1 w_0 \in \mathfrak{S}_n$. Furthermore,*

$$\frac{(-1)^{a(w)}}{(n-1)!} Q_w = \rho_{21^{n-2}} + (n-1)\rho_{1^n}$$

is the unipotent part of a projective indecomposable $kSU_n(q)$ -module.

Here w_0 is the longest element of $W \cong \mathfrak{S}_n$ and s_i is the transposition $(i, i + 1)$.

Proof. Let w_I denote the longest element of the parabolic subgroup of W generated by s_2, \dots, s_{n-2} . Using the fact that the Schubert varieties associated with w_0, w_1 and $s_1 w_0$ are smooth, one finds the following relation between the corresponding Kazhdan–Lusztig basis elements:

$$C_{s_1 w_0 | v=1} = -C_{w_0 | v=1} + s_1 s_2 \cdots s_{n-1} C_{w_I | v=1}.$$

Given a partition λ , the multiplicity of ρ_λ in Q_w is $(-1)^{A(\lambda)} \chi_\lambda(C_{w | v=1} w_0)$, where $A(\lambda)$ denotes the degree in q of the degree polynomial of the unipotent character ρ_λ . For $w = s_1 w_0$ the value of χ_λ can be computed easily from the above formula. \square

3.2. Groups of small rank

We finish by computing for several groups G of small rank the contribution to the principal ℓ -block b by various Q_w 's. In the table below, we give in the second column the minimal integer d such that $\ell \mid (q^d - 1)$, and in the last column the decomposition of Q_w in the basis of projective indecomposable modules which is obtained from the decomposition matrices in [\[8,3,4,9\]](#). Note however that this might change when ℓ is small as the decomposition matrix will change (see for example $G_2(4)$ with $\ell = 5$). In any case, Q_w remains the unipotent part of a proper projective character. Our notation for the unipotent characters is as in the cited sources.

G	d	w	λ	$bQ_w[\lambda]$
$Sp_6(q)$	2	$s_1 s_2 s_1 s_2 s_3$	1	$3(\rho_{1,1^2} + St) + (B_2, 1^2 + 2St) + (\rho_{1^3, -} + 3St)$
$G_2(q)$	2	$s_1 s_2$	-1	$G_2[-1] + 2St$
${}^3D_4(q)$	6	$s_1 s_2 s_3 s_2$	1	$2(\phi_{2,2} + \phi'_{1,3} + St) + 2({}^3D_4[-1] + 2St)$
$SU_6(q)$	2	$s_1 s_3 s_4 s_3$	1	$2(\rho_{321} + 2\rho_{2^3} + 2\rho_{31^3} + 2\rho_{2^2 1^2} + 2\rho_{21^4} + 6\rho_{1^6})$
$F_4(q)$	12	$s_1 s_2 s_1 s_3 s_2 s_1 s_3 s_2 s_4 s_3 s_2 s_3$	ζ_{12}^5	$2F_4[\theta] + 20B_2, St + 4St$
			ζ_{12}^4	$4F_4[\theta] + F_4[i] + 11B_2, St + 10St$

One can even construct larger examples for which the conjecture holds. We give below five examples of computations of Q_w in $SU_{10}(q)$ for $\ell|(q+1)$. When $\ell > 17$, we can use [4, Thm. 6.2] to decompose them on the basis of projective indecomposable modules. It turns out that at least four of them are (up to a scalar) the character of a projective indecomposable module.

w	Q_w
$s_1s_3s_4s_3s_5s_4s_3s_6s_5s_4s_3s_7s_6s_5s_4s_3s_9s_8s_7s_6s_5s_4s_3$	$1440\Psi_{321^5}$
$s_1s_2s_1s_4s_5s_4s_6s_5s_4s_7s_6s_5s_4s_8s_7s_6s_5s_4$	$96\Psi_{32^21^3}$
$s_5s_4s_3s_2s_1s_4s_3s_2s_4s_3s_4s_9s_8s_7s_9s_8s_9$	$576\Psi_{32^31}$
$s_1s_2s_1s_4s_5s_4s_6s_5s_4s_7s_6s_5s_4s_9$	$96\Psi_{3^221^2}$
$s_2s_1s_2s_3s_7s_6s_5s_7s_6s_7s_9$	$16(\Psi_{4321} + (1 - \alpha)\Psi_{32^31} + \beta\Psi_{2^412})$

Here $\alpha, \beta \in \{0, 1\}$ are as in [4, Thm. 6.2]

Acknowledgements

We would like to thank Jean Michel for helping us to compute the characters of the modules Q_w in Chevie [12], especially for $SU_{10}(q)$.

References

[1] F. Digne, J. Michel, Representations of Finite Groups of Lie Type, Lond. Math. Soc. Stud. Texts, vol. 21, Cambridge University Press, 1991.
 [2] F. Digne, J. Michel, R. Rouquier, Cohomologie des variétés de Deligne–Lusztig, Adv. Math. 209 (2007) 749–822.
 [3] O. Dudas, A note on decomposition numbers for groups of Lie type of small rank, J. Algebra 388 (2013) 364–373.
 [4] O. Dudas, G. Malle, Decomposition matrices for low rank unitary groups, 2013, submitted for publication.
 [5] M. Geck, Basic sets of Brauer characters of finite groups of Lie type II, J. Lond. Math. Soc. 47 (1993) 255–268.
 [6] M. Geck, G. Hiss, Basic sets of Brauer characters of finite groups of Lie type, J. Reine Angew. Math. 418 (1991) 173–188.
 [7] M. Geck, G. Hiss, Modular representations of finite groups of Lie type in non-defining characteristic, in: Finite Reductive Groups, Luminy, 1994, in: Prog. Math., vol. 141, Birkhäuser Boston, Boston, MA, 1997, pp. 195–249.
 [8] F. Himstedt, F. Noeske, Decomposition numbers of $SO_7(q)$ and $Sp_6(q)$, preprint, 2013.
 [9] G. Hiss, F. Lübeck, The Brauer trees of the exceptional Chevalley groups of types F_4 and 2E_6 , Arch. Math. (Basel) 70 (1998) 16–21.
 [10] C. Köhler, Unipotente Charaktere und Zerlegungszahlen der endlichen Chevalleygruppen vom Typ F_4 , Dissertation, RWTH Aachen, Germany, 2006.
 [11] G. Lusztig, Characters of Reductive Groups over a Finite Field, Ann. Math. Stud., vol. 107, Princeton University Press, Princeton, NJ, 1984.
 [12] J. Michel, The GAP-part of the Chevie system, GAP 3-package available for download from <http://www.math.jussieu.fr/~jmicHEL/chevie/chevie.html>.