



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Dynamical systems

On the density of singular hyperbolic three-dimensional vector fields: a conjecture of Palis



Sur la densité de l'hyperbolicité singulière pour les champs de vecteurs en dimension trois : une conjecture de Palis

Sylvain Crovisier^{a,1}, Dawei Yang^{b,2}

^a CNRS – Laboratoire de mathématiques d'Orsay, Université Paris-Sud 11, 91405 Orsay, France

^b School of Mathematical Sciences, Soochow University, Suzhou, 215006, PR China

ARTICLE INFO

Article history:

Received 15 October 2014

Accepted 23 October 2014

Available online 20 November 2014

Presented by Étienne Ghys

ABSTRACT

In this note we announce a result for vector fields on three-dimensional manifolds: those who are singular hyperbolic or exhibit a homoclinic tangency form a dense subset of the space of C^1 -vector fields. This answers a conjecture by Palis. The argument uses an extension for local fibred flows of Mañé and Pujals–Sambarino's theorems about the uniform contraction of one-dimensional dominated bundles.

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Dans cette note, nous annonçons un résultat portant sur les champs de vecteurs des variétés de dimension 3 : ceux qui vérifient l'hyperbolicité singulière ou qui possèdent une tangence homocline forment un sous-ensemble dense de l'espace des champs de vecteurs C^1 . Ceci répond à une conjecture de Palis. La démonstration utilise une généralisation pour les flots fibrés locaux des théorèmes de Mañé et Pujals–Sambarino traitant de la contraction uniforme de fibrés unidimensionnels dominés.

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

We are concerned with dynamics that are typical in the space of dynamical systems. Hyperbolic systems are natural candidates since they form an open set. However, there are obstructions for hyperbolicity, such as homoclinic tangencies that produce rich behaviors as Newhouse phenomena [13]. Palis [14,15] conjectured that on surfaces, every diffeomorphism can be accumulated either by hyperbolic ones or by diffeomorphisms with a homoclinic tangency. Pujals and Sambarino

E-mail addresses: Sylvain.Crovisier@math.u-psud.fr (S. Crovisier), yangdw@suda.edu.cn, yangdw1981@gmail.com (D. Yang).

¹ S. Crovisier was partially supported by International Balzan Foundation, Balzan Research Project of J. Palis.

² D. Yang was partially supported by NSFC 11271152 and ANR project DynNonHyp BLAN08-2313375.

[17] managed to prove the conjecture in the C^1 topology. For higher dimensions, other homoclinic bifurcations have to be included in Palis conjecture, such as heterodimensional cycles. For the case of vector fields, new kinds of bifurcations, associated with singularities, and called singular cycles, have to be introduced. A version of Palis conjecture for vector fields can be formulated as:

Conjecture 1. (See Palis [16].) *Every vector field can be accumulated either by hyperbolic vector fields or by ones with a homoclinic bifurcation or with a singular cycle.*

Vector fields admit robustly non-hyperbolic transitive attractors with singularities [1,6]. Morales, Pacifico and Pujals [12] defined the *singular hyperbolicity* to characterize robust attractors with singularities in dimension 3. Palis gave a stronger version of his conjecture in dimension 3 (see also [2,11]):

Conjecture 2. (See Palis [16].) *Every vector field on a three-dimensional manifold can be accumulated either by singular hyperbolic vector fields or by ones with a homoclinic tangency (associated with a non-singular periodic orbit).*

Arroyo and Rodriguez Hertz [2] proved the first conjecture for three-dimensional manifolds in the C^1 topology. The goal of this work is to get a positive answer of the second one in the C^1 topology. Generalizations of singular hyperbolicity in higher dimension have been proposed [10,18], but it is not clear for us what should be the generalization of **Conjecture 2**. Since Newhouse phenomenon requires C^2 -smoothness, an even stronger result could be imagined, specific to C^1 -topology: singular hyperbolicity may be C^1 -dense in the space of three-dimensional vector fields.

2. Precise statements

Let M be a three-dimensional compact Riemannian manifold without boundary. A smooth vector field X on M generates a flow φ_t . A point x is *regular* if $X(x) \neq 0$; otherwise, x is *singular*. The set of singularities plays a particular role and is denoted by $\text{Sing}(X)$. The derivative $D\varphi_t$ of φ_t w.r.t. the space variable is called the *tangent flow*.

The dynamics of the flow is usually split in the following way. For any $\varepsilon > 0$, an ε -pseudo orbit from x to y is a sequence $\{x_i\}_{i=0}^n$ such that $x_0 = x$, $x_n = y$, $n \geq 1$ and such that $d(\varphi_{t_i}(x_i), x_{i+1}) < \varepsilon$ for any $i \in \{0, \dots, n-1\}$ and for some $\{t_i\}_{i=0}^{n-1}$ in $[1, 2]$. Any two points $x, y \in M$ are said to be *chain related* if for any $\varepsilon > 0$, there are ε -pseudo orbits from x to y and from y to x . If x is chain related to itself, then x is called a *chain recurrent point*. The (compact invariant) set of chain recurrent points is called the *chain recurrent set* of X . To be chain related is an equivalence relation on the chain recurrent set; the equivalence classes are called *chain recurrence classes* of X . They are compact, invariant (and pairwise disjoint).

Let Λ be an invariant compact set. An invariant continuous splitting $T_\Lambda M = E \oplus F$ (where E, F are non-trivial vector bundles) is *dominated* if there exist constants $C, \lambda > 0$ such that for any $t > 0$, any $x \in \Lambda$ and for any unit vectors $u \in E(x)$ and $v \in F(x)$, we have $\|D\varphi_t(u)\| \leq Ce^{-\lambda t} \|D\varphi_t(v)\|$.

The set Λ is *hyperbolic* if there exist a continuous invariant splitting $T_\Lambda M = E^s \oplus \langle X \rangle \oplus E^u$ w.r.t. $D\varphi_t$ and constants $C, \lambda > 0$ such that for any $x \in \Lambda$ and $t > 0$, one has $\|D\varphi_t|_{E^s(x)}\| \leq Ce^{-\lambda t}$ and $\|D\varphi_{-t}|_{E^u(x)}\| \leq Ce^{-\lambda t}$.

An attractor Λ is *singular hyperbolic* if there are a dominated splitting $T_\Lambda M = E^{ss} \oplus E^{cu}$ w.r.t. $D\varphi_t$ and constants $C > 0$ and $\lambda > 0$ such that:

- contraction: for any $t > 0$ and any $x \in \Lambda$, $\|D\varphi_t|_{E^{ss}(x)}\| \leq Ce^{-\lambda t}$.
- area-expansion: for any $t > 0$ and any $x \in \Lambda$, $|\text{Det}\varphi_{-t}|_{E^{cu}(x)}| \leq Ce^{-\lambda t}$.

A transitive repeller is *singular hyperbolic* if it is a singular hyperbolic attractor for $-X$.

We say that X is *singular hyperbolic* if the chain-recurrent set of X is the union of finitely pairwise disjoint invariant compact sets $\{\Lambda_i\}$ such that each Λ_i is a hyperbolic set, a singular hyperbolic attractor, or a singular hyperbolic repeller. X is *robustly singular hyperbolic* if any vector field C^1 -close to X is singular hyperbolic.

We say that X has a *homoclinic tangency* if X has a (non-singular) hyperbolic periodic orbit γ such that the stable manifold of γ and the unstable manifold of γ have some non-transverse intersection.

We announce that an answer to the above Palis **Conjecture 2** is:

Main Theorem. *In the C^1 topology, every three-dimensional vector field can be accumulated by robustly singular hyperbolic vector fields, or by vector fields with homoclinic tangencies.*

To prove this theorem, we only need to prove the following (which was already obtained by Arroyo and Rodriguez Hertz [2] for non-singular chain-recurrence classes).

Main Theorem Restated. *For a C^1 generic vector field X on a three-dimensional manifold that is far from homoclinic tangencies, any chain-recurrence class is hyperbolic or is a singular hyperbolic attractor or repeller.*

3. Dominated splittings for tangent vs. linear Poincaré flows

The dynamics induces another linear flow above the set of regular points. At any point $x \in M \setminus \text{Sing}(X)$, we introduce the plane $\mathcal{N}_x = X(x)^\perp$ and define the normal bundle $\mathcal{N} = \coprod_{x \in M \setminus \text{Sing}(X)} \mathcal{N}_x$. By orthogonal projection of the tangent flow $D\varphi_t$, one gets the *linear Poincaré flow* ψ_t on \mathcal{N} .

Let C be any chain-recurrence class (which is not an isolated singularity) of a C^1 -generic vector fields far from homoclinic tangencies. Using technics developed in the different works on robustly transitive sets [12,7,11] and Liao's estimation [8], Gan and Yang [5] have proved the following properties (up to replace X by $-X$):

1. Any non-isolated singularity $\sigma \in C$ is *Lorenz-like*: it has three real eigenvalues satisfying $\lambda_1 < \lambda_2 < 0 < -\lambda_2 < \lambda_3$. Moreover the (one-dimensional) strong stable manifold satisfies $W^{ss}(\sigma) \cap C = \{\sigma\}$.
2. The linear Poincaré flow on $C \setminus \text{Sing}(X)$ has a dominated splitting.
3. [5, Theorem C]. If the tangent flow $D\varphi_t$ on Λ has a dominated splitting and if C contains a singularity, then C is a singular hyperbolic attractor.

Thus our main theorem essentially reduces to comparing the dominated splitting of the linear Poincaré flow and of the tangent flow. (Note that the following result goes beyond C^1 -generic vector fields.)

Theorem 1 (*Dominated splitting for the tangent flow*). Consider any C^3 three-dimensional vector field X and any compact invariant set Λ with the following properties:

- every periodic orbit in Λ is a hyperbolic saddle;
- every singularity $\sigma \in \Lambda$ is Lorenz-like and $W^{ss}(\sigma) \cap \Lambda = \{\sigma\}$;
- Λ does not contain a minimal repeller whose dynamics is the suspension of an irrational rotation of the circle.

Then the tangent flow $D\varphi_t$ on Λ has a dominated splitting if and only if the linear Poincaré flow on $\Lambda \setminus \text{Sing}(X)$ has a dominated splitting.

4. Uniform contraction for dominated fibred dynamics

The existence of singularities introduces several difficulties: the regular orbits may separate when passing the singularities, which causes a lack of compactness and of uniformity. These problems have been overcome in previous works in two ways:

- by blowing up the singular set and extending the flow (introduced by Li–Gan–Wen [7]),
- by rescaling the flow near the singular set (in Liao's work [8] and in [5]).

These ideas may be applied to several flows associated with the initial flow φ_t : the flow itself, the sectional Poincaré flow, their tangent flows...

In this work we get a compactification of the rescaled sectional Poincaré flow that we define now. For any regular points x and time $t_0 \in \mathbb{R}$, the flow φ_t induces a local holonomy map from a neighborhood of x in $\exp_x(\mathcal{N}_x)$ to a neighborhood of $\varphi_{t_0}(x)$ in $\exp_{\varphi_{t_0}(x)}(\mathcal{N}_{\varphi_{t_0}(x)})$. This induces a local map P_{t_0} from \mathcal{N}_x to $\mathcal{N}_{\varphi_{t_0}(x)}$ that preserves 0 and gives a local flow P_t on the bundle \mathcal{N} in a neighborhood of the 0-section. It is called the *sectional Poincaré flow*. Its linearization at the 0-section is the linear Poincaré flow ψ_t .

For these flows, we define the rescaled sectional Poincaré flow and the rescaled linear Poincaré flow as:

$$P_t^*(v_x) = \|X(\varphi_t(x))\|^{-1} \cdot P_t(\|X(x)\| \cdot v_x) \quad \text{and} \quad \psi_t^*(v_x) = \frac{\|X(x)\|}{\|X(\varphi_t(x))\|} \cdot \psi_t(v_x).$$

Still ψ_t^* can be seen as the linearization of P_t^* .

Theorem 2 (*Compactification of the sectional Poincaré flow*). Assume that the flow φ_t is C^r , $r \geq 2$, and let K be an invariant compact set whose singularities $\text{Sing}(X) \cap K$ are hyperbolic. Then, there exists a flow $\widehat{\varphi}_t$ on a compact metric space \widehat{K} and a local flow \widehat{P}_t^* on a linear bundle $\widehat{\mathcal{N}}$ over the dynamics of $\widehat{\varphi}_t$ on \widehat{K} , which preserves the 0-section, such that:

- there exists an injective continuous fiber-preserving map $p: \mathcal{N} \rightarrow \widehat{\mathcal{N}}$ which conjugates the local flows P_t^* and \widehat{P}_t^* ;
- \widehat{P}_t^* is C^{r-1} -along fibers: it induces local C^{r-1} -diffeomorphisms $(\widehat{\mathcal{N}}_x, 0) \rightarrow (\widehat{\mathcal{N}}_{\widehat{\varphi}_t(x)}, 0)$ depending continuously on x for the C^{r-1} -topology.

Theorem 1 can be translated for the flow \widehat{P}_t^* with the following remarks:

- if the linear Poincaré flow ψ_t has a dominated splitting, then the 0-section in $\widehat{\mathcal{N}}$ has a dominated splitting $T\widehat{\mathcal{N}} = \widehat{E} \oplus \widehat{F}$ for the compactified rescaled flow \widehat{P}_t^* ,
- if moreover \widehat{E} is uniformly contracted by the linearization of \widehat{P}_t^* , then K has a dominated splitting $T_K M = E \oplus F$ for the tangent flow $D\varphi_t$, with $\dim(E) = 1$ and $\mathbb{R} \cdot X \subset F$.

Theorem 1 is thus a consequence of the theorem below for dominated local fibred flow \widehat{P}_t^* on a bundle $\widehat{\mathcal{N}}$ with 2-dimensional fibers. It requires an important *compatibility assumption* which reflects the fact that \widehat{P}_t^* has been obtained from the sectional Poincaré flow on a manifold:

1. There exists an open set $U \subset \widehat{K}$ and for close points $x, y \in U$ an identification map $\pi_{x,y}: \widehat{\mathcal{N}}_x \rightarrow \widehat{\mathcal{N}}_y$ which is “compatible” with the local flow \widehat{P}_t^* .
2. Along pieces of orbit in $\widehat{K} \setminus U$, the bundle \widehat{E} is uniformly contracted.

Theorem 3 (Uniform contraction of one-dimensional bundle for dominated local fibred flow). Let \widehat{P}_t^* be a local fibred flow which is C^2 -along fibers, on a bundle $\widehat{\mathcal{N}}$ with 2-dimensional fibers satisfying:

- the 0-section is invariant and has a dominated splitting $T\widehat{\mathcal{N}} = \widehat{E} \oplus \widehat{F}$;
- the compatibility assumption holds;
- E is contracted along each periodic orbit of the base space \widehat{K} of $\widehat{\mathcal{N}}$;
- the base space \widehat{K} of $\widehat{\mathcal{N}}$ does not contain any invariant compact set which is a repeller and whose dynamics is conjugate to the suspension of an irrational circle rotation.

Then \widehat{E} is uniformly contracted by the linearization of the flow \widehat{P}_t^* .

This theorem is a continuation of a sequence of works initiated by Mañé [9] and Pujals–Sambarino [17]: the latter proves the hyperbolicity of dominated invariant compact set for surface diffeomorphisms. Compared with the usual Pujals–Sambarino arguments, we meet the following difficulties:

- work with continuous time dynamics. It is not enough to consider the time-one map since the flow shears along the orbits. Bounded time evolution produces small shear, but we need to consider long-term behaviors. This difficulty already appears in Arroyo and Rodriguez Hertz’ result [2];
- there is no extension of the Pujals–Sambarino result to general dominated fibred dynamics (as one can see considering a product of a minimal base dynamics with the identity along fibers). Our proof uses strongly the identification maps $\pi_{x,y}$;
- since the identifications $\pi_{x,y}$ are only defined on U , we have to handle with the induced dynamics in U . This makes us to consider “induced hyperbolic returns” (as in [3]);
- the flow \widehat{P}_t^* is locally defined and the global arguments of [17] have to be replaced by local ones;
- we need to construct some Markovian boxes in a non-symmetric setting: the bundles \widehat{E} , \widehat{F} play different roles. We borrow some ideas from a recent work of Crovisier, Pujals and Sambarino [4].

References

- [1] V. Afraimovič, V. Bykov, L. Silnikov, The origin and structure of the Lorenz attractor, Dokl. Akad. Nauk SSSR 234 (1977) 336–339.
- [2] A. Arroyo, F. Rodriguez Hertz, Homoclinic bifurcations and uniform hyperbolicity for three-dimensional flows, Ann. Inst. Henri Poincaré, Anal. Non Linéaire 20 (2003) 805–841.
- [3] S. Crovisier, E. Pujals, Essential hyperbolicity and homoclinic bifurcations: a dichotomy phenomenon/mechanism for diffeomorphisms, Invent. Math. (2014), <http://dx.doi.org/10.1007/s00222-014-0553-9>, Preprint, arXiv:1011.3836.
- [4] S. Crovisier, M. Sambarino, E. Pujals, Hyperbolicity of the extremal bundles, in preparation.
- [5] S. Gan, D. Yang, Morse–Smale systems and non-trivial horseshoes for three-dimensional singular flows, Preprint, arXiv:1302.0946v1.
- [6] J. Guckenheimer, A strange, strange attractor, in: The Hopf Bifurcation Theorems and Its Applications, in: Applied Mathematical Series, vol. 19, Springer-Verlag, 1976, pp. 368–381.
- [7] M. Li, S. Gan, L. Wen, Robustly transitive singular sets via approach of extended linear Poincaré flow, Discrete Contin. Dyn. Syst. 13 (2005) 239–269.
- [8] S. Liao, On (η, d) -contractible orbits of vector fields, Syst. Sci. Math. Sci. 2 (1989) 193–227.
- [9] R. Mañé, Hyperbolicity, sinks and measure in one-dimensional dynamics, Commun. Math. Phys. 100 (1985) 495–524.
- [10] R. Metzger, C. Morales, Sectional-hyperbolic systems, Ergod. Theory Dyn. Syst. 28 (2008) 1587–1597.
- [11] C. Morales, M. Pacifico, A dichotomy for three-dimensional vector fields, Ergod. Theory Dyn. Syst. 23 (2003) 1575–1600.
- [12] C. Morales, M. Pacifico, E. Pujals, Robust transitive singular sets for 3-flows are partially hyperbolic attractors or repellers, Ann. Math. 160 (2004) 375–432.
- [13] S. Newhouse, The abundance of wild hyperbolic sets and nonsmooth stable sets for diffeomorphisms, Publ. Math. IHÉS 50 (1979) 101–151.
- [14] J. Palis, Homoclinic bifurcations, sensitive-chaotic dynamics and strange attractors, in: Dynamical Systems and Related Topics, Nagoya, 1990, in: Adv. Ser. Dynam. Systems, 1991, pp. 466–472.
- [15] J. Palis, A global view of dynamics and a conjecture of the denseness of finitude of attractors, Astérisque 261 (2000) 335–347.
- [16] J. Palis, A global perspective for non-conservative dynamics, Ann. Inst. Henri Poincaré, Anal. Non Linéaire 22 (2005) 485–507.
- [17] E. Pujals, M. Sambarino, Homoclinic tangencies and hyperbolicity for surface diffeomorphisms, Ann. Math. (2) 151 (2000) 961–1023.
- [18] S. Zhu, S. Gan, L. Wen, Indices of singularities of robustly transitive sets, Discrete Contin. Dyn. Syst. 21 (2008) 945–957.