



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Homological algebra/Algebraic geometry

Algebraic K -theory with coefficients of cyclic quotient singularities

Gonçalo Tabuada ^{a,b,c,1}^a Department of Mathematics, MIT, Cambridge, MA 02139, USA^b Departamento de Matemática, FCT, UNL, Portugal^c Centro de Matemática e Aplicações (CMA), FCT, UNL, Portugal

ARTICLE INFO

Article history:

Received 8 December 2015

Accepted after revision 26 January 2016

Available online 7 March 2016

Presented by Michel Duflo

ABSTRACT

In this note, by combining the work of Amiot–Iyama–Reiten and Thanhoffer de Völcsy–Van den Bergh on Cohen–Macaulay modules with the previous work of the author on orbit categories, we compute the algebraic K -theory with coefficients of cyclic quotient singularities.

© 2016 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Dans cette note, en combinant les travaux de Amiot–Iyama–Reiten et Thanhoffer de Völcsy–Van den Bergh sur les modules Cohen–Macaulay avec le travail précédent de l'auteur sur les catégories d'orbites, nous calculons la K -théorie algébrique avec coefficients des singularités quotient cycliques.

© 2016 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction and statement of results

Let k be an algebraically closed field of characteristic zero. Given an integer $d \geq 2$, consider the associated polynomial ring $S := k[t_1, \dots, t_d]$. Let G be a cyclic subgroup of $SL(d, k)$ generated by $\text{diag}(\zeta^{a_1}, \dots, \zeta^{a_d})$, where ζ is a primitive n^{th} root of unit and a_1, \dots, a_d are integers satisfying the following conditions: we have $0 < a_j < n$ and $\text{gcd}(a_j, n) = 1$ for every $1 \leq j \leq d$; we have $a_1 + \dots + a_d = n$. The group G acts naturally on S and the invariant ring $R := S^G$ is a Gorenstein isolated singularity of Krull dimension d . For example, when $d = 2$, the ring R identifies with the Kleinian singularity $k[u, v, w]/(u^n + vw)$ of type A_{n-1} .

The affine k -scheme $X := \text{Spec}(R)$ is singular. Following Orlov [5,4], we can then consider the associated dg category of singularities $\mathcal{D}_{\text{dg}}^{\text{sing}}(X)$, also known as matrix factorizations or maximal Cohen–Macaulay modules. Roughly speaking, this dg category encodes all the crucial information concerning the isolated singularity of X .

E-mail address: tabuada@math.mit.edu.

¹ The author was partially supported by the National Science Foundation CAREER Award #1350472 and by the Fundação para a Ciência e a Tecnologia (Portuguese Foundation for Science and Technology) through the project grant UID/MAT/00297/2013 (Centro de Matemática e Aplicações).

<http://dx.doi.org/10.1016/j.crma.2016.01.017>

1631-073X/© 2016 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

Let us denote by (Q, ρ) the quiver with relations defined by the following steps:

- (s1) consider the quiver with vertices $\mathbb{Z}/n\mathbb{Z}$ and with arrows $x_j^i: i \rightarrow i + a_j$, where $i \in \mathbb{Z}/n\mathbb{Z}$ and $1 \leq j \leq d$. The relations ρ are given by $x_{j'}^{i+a_j} x_j^i = x_j^{i+a_{j'}} x_{j'}^i$ for every $i \in \mathbb{Z}/n\mathbb{Z}$ and $1 \leq j, j' \leq d$;
- (s2) remove from (s1) all arrows $x_j^i: i \rightarrow i'$ with $i > i'$;
- (s3) remove from (s2) the vertex 0.

Consider the matrix $(n - 1) \times (n - 1)$ matrix C such that C_{ij} equals the number of arrows in Q from j to i (counted modulo the relations). Let us write M for the matrix $(-1)^{d-1} C(C^{-1})^T - \text{Id}$ and $M: \oplus_{r=1}^{n-1} \mathbb{Z}/l^v \rightarrow \oplus_{r=1}^{n-1} \mathbb{Z}/l^v$ for the associated (matrix) homomorphism, where l^v is a (fixed) prime power.

Theorem 1.1. *We have the following computation:*

$$\mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X); \mathbb{Z}/l^v) \simeq \begin{cases} \text{cokernel of } M & \text{if } i \geq 0 \text{ even,} \\ \text{kernel of } M & \text{if } i \geq 0 \text{ odd,} \\ 0 & \text{if } i < 0. \end{cases}$$

Thanks to [Theorem 1.1](#), the computation of the (nonconnective) algebraic K -theory with coefficients of the cyclic quotient singularities reduces to the computation of (co)kernels of explicit matrix homomorphisms! To the best of the author’s knowledge, these computations are new in the literature. In the particular case of Kleinian singularities of type A_n they were originally established in [\[8, §3\]](#).

Corollary 1.2.

- (i) *If there exists a prime power l^v and an even (resp. odd) integer $j \geq 0$ such that $\mathbf{K}_j(\mathcal{D}_{\text{dg}}^{\text{sing}}(X); \mathbb{Z}/l^v) \neq 0$, then for every even (resp. odd) integer $i \geq 0$ at least one of the groups $\mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X))$ and $\mathbf{K}_{i-1}(\mathcal{D}_{\text{dg}}^{\text{sing}}(X))$ is non-zero.*
- (ii) *If there exists a prime power l^v such that $\mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X); \mathbb{Z}/l^v) = 0$ for every $i \geq 0$, then the groups $\mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X))$, $i \geq 0$, are uniquely l^v -divisible.*


Proof. Combine the following universal coefficients sequence (see [\[8, §5\]](#))

$$0 \longrightarrow \mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X)) \otimes_{\mathbb{Z}} \mathbb{Z}/l^v \longrightarrow \mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X); \mathbb{Z}/l^v) \longrightarrow \{l^v\text{-torsion in } \mathbf{K}_{i-1}(\mathcal{D}_{\text{dg}}^{\text{sing}}(X))\} \longrightarrow 0$$

with the computation of [Theorem 1.1](#). \square

2. A low dimensional example

When $d = 3, n = 5, a_1 = 1$, and $a_2 = a_3 = 2$, the three steps (s1)–(s3) lead to the following quiver




with relations $xy = yx, yz = zy$, and $zx = xz$. Consequently, we obtain the following matrix

$$M = \begin{pmatrix} 0 & -1 & -3 & -3 \\ 1 & -1 & -4 & -6 \\ 3 & -2 & -10 & -13 \\ 3 & 0 & -11 & -19 \end{pmatrix}.$$

Since $\det(M) = 26$, we have $\mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X); \mathbb{Z}/l^v) = 0$ whenever $l \neq 2, 13$. In the remaining two cases, a computation shows that $\mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X); \mathbb{Z}/l^v) \simeq \mathbb{Z}/l$ for every $i \geq 0$. Thanks to [Corollary 1.2](#), this implies that for every $i \geq 0$, at least one of the groups $\mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X))$ and $\mathbf{K}_{i-1}(\mathcal{D}_{\text{dg}}^{\text{sing}}(X))$ is non-zero. Moreover, the groups $\mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X))$, $i \geq 0$, are uniquely l -divisible for every prime $l \neq 2, 13$.

3. A family of examples

When $n = d \geq 3$ and $a_1 = \dots = a_d = 1$, the above three steps (s1)–(s3) lead to the following quiver



with relations $x_j x_i = x_i x_j$. In the case where d is odd, we obtain the matrix

$$M_{ij} = \begin{cases} -\sum_{r=0}^{i-1} \binom{d}{r} \binom{d}{(j-i)+r} & \text{if } i < j, \\ -\sum_{r=1}^{i-1} \binom{d}{r}^2 & \text{if } i = j, \\ -\sum_{r=1}^{j-1} \binom{d}{(i-j)+r} \binom{d}{r} + \binom{d}{i-j} & \text{if } i > j, \end{cases}$$

where $\binom{d}{r}$ stands for the multicomination symbol.² Similarly, in the case where d is even, we obtain

$$M_{ij} = \begin{cases} \sum_{r=0}^{i-1} \binom{d}{r} \binom{d}{(j-i)+r} & \text{if } i < j, \\ -2 + \sum_{r=1}^{i-1} \binom{d}{r}^2 & \text{if } i = j, \\ \sum_{r=1}^{j-1} \binom{d}{(i-j)+r} \binom{d}{r} - \binom{d}{i-j} & \text{if } i > j. \end{cases}$$

Whenever d is a prime number, all the multicombinations

$$\binom{d}{r} = \binom{d+r-1}{r} = \frac{(d+r-1) \cdots d(d-1)!}{r!(d-1)!}, \quad 0 \leq r \leq d-2$$

are multiples of d . This implies that the homomorphism $M: \bigoplus_{r=1}^{d-1} \mathbb{Z}/d \rightarrow \bigoplus_{r=1}^{d-1} \mathbb{Z}/d$ is zero, and consequently that $\mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X); \mathbb{Z}/d) \simeq \bigoplus_{r=1}^{d-1} \mathbb{Z}/d$ for every $i \geq 0$. Thanks to Corollary 1.2(i), we hence conclude that for every $i \geq 0$ at least one of the groups $\mathbf{K}_i(\mathcal{D}_{\text{dg}}^{\text{sing}}(X))$ and $\mathbf{K}_{i-1}(\mathcal{D}_{\text{dg}}^{\text{sing}}(X))$ is non-zero.

4. Proof of Theorem 1.1

Let A be a finite dimensional k -algebra of finite global dimension. We write $\mathcal{D}^b(A)$ for the bounded derived category of (right) A -modules and $\mathcal{D}_{\text{dg}}^b(A)$ for the canonical dg enhancement of $\mathcal{D}^b(A)$. Consider the dg functors $\tau^{-1} \Sigma^d: \mathcal{D}_{\text{dg}}^b(A) \rightarrow \mathcal{D}_{\text{dg}}^b(A)$, $d \geq 0$, where τ stands for the Auslander–Reiten translation. Following Keller [3, §7.2], we can consider the associated dg orbit category $\mathcal{C}_A^{(d)} := \mathcal{D}_{\text{dg}}^b(A)/(\tau^{-1} \Sigma^d)^{\mathbb{Z}}$. Similarly to [8, Thm. 2.5] (consult [7, §2]), we have a distinguished triangle of spectra

$$\bigoplus_{r=1}^{\nu} \mathbf{K}(k; \mathbb{Z}/l^{\nu}) \xrightarrow{(-1)^d \Phi_A - \text{Id}} \bigoplus_{r=1}^{\nu} \mathbf{K}(k; \mathbb{Z}/l^{\nu}) \longrightarrow \mathbf{K}(\mathcal{C}_A^{(d)}; \mathbb{Z}/l^{\nu}) \longrightarrow \bigoplus_{r=1}^{\nu} \Sigma \mathbf{K}(k; \mathbb{Z}/l^{\nu}),$$

where ν stands for the number of simple (right) A -modules and Φ_A for the inverse of the Coxeter matrix of A . Consider the following (matrix) homomorphism

$$(-1)^d \Phi_A - \text{Id}: \bigoplus_{r=1}^{\nu} \mathbb{Z}/l^{\nu} \longrightarrow \bigoplus_{r=1}^{\nu} \mathbb{Z}/l^{\nu}. \tag{1}$$

As proved by Suslin in [6, Cor. 3.13], we have $\mathbf{K}_i(k; \mathbb{Z}/l^{\nu}) \simeq \mathbb{Z}/l^{\nu}$ when $i \geq 0$ is even and $\mathbf{K}_i(k; \mathbb{Z}/l^{\nu}) = 0$ otherwise. Consequently, making use of the long exact sequence of algebraic K -theory groups with coefficients associated with the above distinguished triangle of spectra, we obtain the following computations:

$$\mathbf{K}_i(\mathcal{C}_A^{(d)}; \mathbb{Z}/l^{\nu}) \simeq \begin{cases} \text{cokernel of (1)} & \text{if } i \geq 0 \text{ even,} \\ \text{kernel of (1)} & \text{if } i \geq 0 \text{ odd,} \\ 0 & \text{if } i < 0. \end{cases}$$

² Also usually known as the *multisubset* symbol.

Consider also the dg functors $S^{-1}\Sigma^d: \mathcal{D}_{\text{dg}}^b(A) \longrightarrow \mathcal{D}_{\text{dg}}^b(A)$, $d \geq 0$, where S stands for the Serre dg functor. The associated dg orbit category $\mathcal{C}_d(A) := \mathcal{D}_{\text{dg}}^b(A)/(S^{-1}\Sigma^d)^{\mathbb{Z}}$ is usually called the *generalized d -cluster dg category of A* ; see [1, §1.3] and the references therein. Since $S^{-1}\Sigma = \tau^{-1}$, we have $\mathcal{C}_d(A) \simeq \mathcal{C}_A^{(d-1)}$.

Now, let us take for A the k -algebra $kQ/\langle \rho \rangle$ associated with the quiver with relations (Q, ρ) . As proved independently by Amiot–Iyama–Reiten [1, §5] and Thanhoffer de Völcsey–Van den Bergh [2], we have $\mathcal{D}_{\text{dg}}^{\text{sing}}(X) \simeq \mathcal{C}_{d-1}(A)$. Consequently, it remains then only to show that the homomorphism (1), with d replaced by $d - 2$, agrees with the homomorphism M associated with the matrix $M := (-1)^{d-1}C(C^{-1})^T - \text{Id}$. On the one hand, the number of simple (right) A -modules agrees with the number of vertices of the quiver Q . This implies that $v = n - 1$. On the other hand, the inverse of the Coxeter matrix of A can be expressed as $-C(C^{-1})^T$, where C_{ij} equals the number of arrows in Q from j to i (counted modulo the relations). This implies that $(-1)^{d-2}\Phi_A - \text{Id} = (-1)^{d-1}C(C^{-1})^T - \text{Id} = M$, and hence concludes the proof.

Acknowledgement

The author is grateful to Michel Van den Bergh for useful discussions.

References

- [1] C. Amiot, O. Iyama, I. Reiten, Stable categories of Cohen–Macaulay modules and cluster categories, *Amer. J. Math.* 137 (3) (2015) 813–857.
- [2] L. de Thanhoffer, M. Van den Bergh, Explicit models for some stable categories of maximal Cohen–Macaulay modules, arXiv:1006.2021, 2016, *Math. Res. Lett.*, in press.
- [3] B. Keller, On triangulated orbit categories, *Doc. Math.* 10 (2005) 551–581.
- [4] D. Orlov, Triangulated categories of singularities and D -branes in Landau–Ginzburg models, *Proc. Steklov Inst. Math.* 246 (2004) 227–248.
- [5] D. Orlov, Derived categories of coherent sheaves and triangulated categories of singularities, in: *Algebra, Arithmetic, and Geometry: In Honor of Yu.I. Manin*, vol. II, in: *Progr. Math.*, vol. 270, Birkhäuser Boston, Inc., Boston, MA, USA, 2009, pp. 503–531.
- [6] A. Suslin, On the K -theory of local fields, in: *Proceedings of the Luminy Conference on Algebraic K -Theory*, Luminy, 1983. *J. Pure Appl. Algebra* 34 (2–3) (1984) 301–318.
- [7] G. Tabuada, \mathbb{A}^1 -homotopy invariants of dg orbit categories, *J. Algebra* 434 (2015) 169–192.
- [8] G. Tabuada, \mathbb{A}^1 -homotopy invariance of algebraic K -theory with coefficients and Kleinian singularities, arXiv:1502.05364, 2016, *Annals of K -theory*, in press.