



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Algebraic geometry

On the Chow ring of certain rational cohomology tori



Sur l'anneau de Chow de certaines variétés dont la cohomologie rationnelle est celle d'un tore

Zhi Jiang^{a,b}, Qizheng Yin^c^a Département de mathématiques, CNRS UMR 8628, Université Paris-Sud, Bâtiment 425, 91405 Orsay cedex, France^b Shanghai Center for Mathematical Sciences, Fudan University, No. 220 Handan Road, Yangpu District, Shanghai 200433, China^c Beijing International Center for Mathematical Research, Peking University, No. 5 Yiheyuan Road, Haidian District Beijing 100871, China

ARTICLE INFO

Article history:

Received 22 March 2017

Accepted 18 April 2017

Available online 28 April 2017

Presented by Claire Voisin

ABSTRACT

Let $f : X \rightarrow A$ be an abelian cover from a complex algebraic variety with quotient singularities to an abelian variety. We show that f^* induces an isomorphism between the rational cohomology rings $H^*(A, \mathbb{Q})$ and $H^*(X, \mathbb{Q})$ if and only if f^* induces an isomorphism between the Chow rings with rational coefficients $\text{CH}^*(A)_{\mathbb{Q}}$ and $\text{CH}^*(X)_{\mathbb{Q}}$.

© 2017 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

R É S U M É

Soit $f : X \rightarrow A$ un revêtement abélien d'une variété algébrique complexe à singularités-quotient dans une variété abélienne. Nous montrons que f^* induit un isomorphisme entre les anneaux de cohomologie rationnelle $H^*(A, \mathbb{Q})$ et $H^*(X, \mathbb{Q})$ si et seulement si f^* induit un isomorphisme entre les anneaux de Chow à coefficients rationnels $\text{CH}^*(A)_{\mathbb{Q}}$ et $\text{CH}^*(X)_{\mathbb{Q}}$.

© 2017 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Rational cohomology tori and the Chow ring

A complex algebraic variety X is called a rational cohomology torus if X is normal and

$$H^*(X, \mathbb{Q}) \simeq \wedge^* H^1(X, \mathbb{Q}).$$

In [3], the authors studied properties of rational cohomology tori. They showed that if X is a rational cohomology torus, then there exists a finite cover $a_X : X \rightarrow A$ to an abelian variety such that $a_X^* : H^*(A, \mathbb{Q}) \rightarrow H^*(X, \mathbb{Q})$ is an isomorphism. The morphism a_X is the universal morphism to an abelian variety and is called the Albanese morphism of X .

It is a general principle that Hodge structures control Chow groups. Hence it makes sense to ask the following question.

E-mail addresses: zhi.jiang@math.u-psud.fr, zhijiang@fudan.edu.cn (Z. Jiang), yinqizheng@math.pku.edu.cn (Q. Yin).

<http://dx.doi.org/10.1016/j.crma.2017.04.006>

1631-073X/© 2017 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Question 1.1. Let $f : X \rightarrow A$ be a finite morphism from a variety with quotient singularities to an abelian variety. Assume that $f^* : H^\bullet(A, \mathbb{Q}) \rightarrow H^\bullet(X, \mathbb{Q})$ is an isomorphism. Does f^* induce an isomorphism of Chow rings with rational coefficients between A and X ?

A variety is said to have quotient singularities if it is étale-locally the quotient of a smooth variety by a finite group. For such varieties there is a well-defined intersection theory with \mathbb{Q} -coefficients (see [7]). Note that a smooth rational cohomology torus cannot be of general type, but in every dimension > 2 , there exist rational cohomology tori of general type with quotient singularities (see [3]). Hence it is more interesting to consider the Chow ring of rational cohomology tori with quotient singularities.

Despite several constraints on a_X found in [3], we do not have a thorough classification of rational cohomology tori. Hence it seems difficult to answer the above question in general. But all known rational cohomology tori are actually abelian covers of abelian varieties (see [3, Section 4]). The goal of this note is to answer affirmatively the above question in the case of abelian covers of abelian varieties.

Let $f : X \rightarrow A$ be an abelian cover from a normal variety to an abelian variety with Galois group G . Then we have (see for instance [6])

$$f_*\mathcal{O}_X = \bigoplus_{\chi \in G^*} L_\chi^{-1},$$

where L_χ is a line bundle on A and G acts on L_χ^{-1} via the character χ . Assume further that X has quotient singularities. Define for each character $\chi \in G^*$ the projector

$$[\Gamma_\chi^X] := \frac{1}{|G|} \sum_{g \in G} \chi^{-1}(g) [\Gamma_g^X] \in \text{CH}^{\dim X}(X \times X)_\mathbb{C},$$

where Γ_g^X is the graph of g . The image of $\text{CH}^\bullet(X)_\mathbb{C}$ under $[\Gamma_\chi^X]$ is

$$\text{CH}^\bullet(X)_\mathbb{C}^\chi := \{\alpha \in \text{CH}^\bullet(X)_\mathbb{C} : g_*(\alpha) = \chi(g)\alpha \text{ for all } g \in G\}.$$

We have a decomposition

$$\text{CH}^\bullet(X)_\mathbb{C} = \bigoplus_{\chi \in G^*} \text{CH}^\bullet(X)_\mathbb{C}^\chi.$$

Theorem 1.2. Let $f : X \rightarrow A$ be an abelian cover from a variety with quotient singularities to an abelian variety with Galois group G . Then the following statements are equivalent:

- (1) X is a rational cohomology torus;
- (2) $h^i(A, L_\chi) = 0$ for all $i \in \mathbb{Z}$ and non-trivial $\chi \in G^*$;
- (3) $[\Gamma_\chi^X] = 0 \in \text{CH}^{\dim X}(X \times X)_\mathbb{C}$ for all non-trivial $\chi \in G^*$; in particular,

$$f^* : \text{CH}^\bullet(A)_\mathbb{Q} \rightarrow \text{CH}^\bullet(X)_\mathbb{Q}$$

is an isomorphism.

The implications (1) \Rightarrow (2) and (3) \Rightarrow (1) are straightforward. Indeed, if X is a rational cohomology torus, then $f^* : H^\bullet(A, \mathbb{Q}) \rightarrow H^\bullet(X, \mathbb{Q})$ is an isomorphism of \mathbb{Q} -Hodge structures. We have for all $i \in \mathbb{Z}$,

$$h^i(A, \mathcal{O}_A) = h^{0,i}(A) = h^{0,i}(X) = h^i(X, \mathcal{O}_X) = \sum_{\chi \in G^*} h^i(A, L_\chi^{-1}).$$

Hence $h^i(A, L_\chi^{-1}) = 0$ for all $i \in \mathbb{Z}$ and non-trivial $\chi \in G^*$. We also have $h^i(A, L_\chi) = 0$ by Serre duality.

If $[\Gamma_\chi^X] = 0 \in \text{CH}^{\dim X}(X \times X)_\mathbb{C}$ for all non-trivial $\chi \in G^*$, then by applying $[\Gamma_\chi^X]$ to $H^\bullet(X, \mathbb{C})$, we find $H^\bullet(X, \mathbb{C}) = H^\bullet(X, \mathbb{C})^G$. Hence $f^* : H^\bullet(A, \mathbb{Q}) \rightarrow H^\bullet(X, \mathbb{Q})$ is an isomorphism and X is a rational cohomology torus.

The proof of (2) \Rightarrow (3) will occupy the next section.

Remark 1.3. Since L_χ is semi-ample (see Step 1 in the proof of Proposition 2.2), condition (2) in Theorem 1.2 is also equivalent to the condition that $h^0(A, L_\chi) = 0$ for all non-trivial $\chi \in G^*$. Hence, under the assumption of Theorem 1.2, we know that X is a rational cohomology torus if and only if $p_g(X) = 1$.

2. Proof of Theorem 1.2

Lemma 2.1. *Let A be an abelian variety and let $t : A \rightarrow A$ be the translation by a torsion element. Then $[\Gamma_t^A] - [\Delta_A] = 0 \in \text{CH}^{\dim A}(A \times A)_{\mathbb{Q}}$, where Γ_t^A is the graph of t and Δ_A is the diagonal.*

Proof. Let $m \in \mathbb{Z}_{>0}$ be the order of t . Recall that $m^* : \text{CH}^{\bullet}(A \times A)_{\mathbb{Q}} \rightarrow \text{CH}^{\bullet}(A \times A)_{\mathbb{Q}}$ is an isomorphism (see [1]). Since $m^*[\Gamma_t^A] = m^*[\Delta_A] \in \text{CH}^{\dim A}(A \times A)_{\mathbb{Q}}$, we conclude that $[\Gamma_t^A] - [\Delta_A] = 0 \in \text{CH}^{\dim A}(A \times A)_{\mathbb{Q}}$. \square

Proposition 2.2. *Let $\rho : Y \rightarrow A$ be a cyclic cover from a variety with quotient singularities to an abelian variety with Galois group H . We write*

$$\rho_* \mathcal{O}_Y = \bigoplus_{\chi \in H^*} L_{\chi}^{-1}.$$

If $h^i(A, L_{\chi}) = 0$ for all $i \in \mathbb{Z}$ and non-trivial $\chi \in H^*$, then for all non-trivial $\chi \in H^*$,

$$\sum_{h \in H} \chi^{-1}(h)[\Gamma_h^Y] = 0 \in \text{CH}^{\dim Y}(Y \times Y)_{\mathbb{C}}.$$

Proof. We proceed by induction on the dimension of Y . The statement in dimension 0 is trivial. We then assume that Proposition 2.2 holds in dimensions $< \dim Y$.

Step 1. Reduction to lower dimensions.

Let χ be a generator of H^* . Since $h^i(A, L_{\chi}) = 0$ for all $i \in \mathbb{Z}$, the line bundle L_{χ} is not ample. Let K denote the neutral component of the morphism $\varphi_{L_{\chi}} : A \rightarrow \text{Pic}^0(A)$ induced by L_{χ} , and let $p : A \rightarrow B$ be the quotient by K . Then we have

$$L_{\chi} = p^*H + Q,$$

where H is an ample line bundle on B and Q is a torsion line bundle such that $Q|_K \neq \mathcal{O}_K$ (see for instance [2, Section 3.3]). As ρ is a cyclic cover and χ is a generator of H^* , for each $k \in \mathbb{Z}$, there exists an effective divisor D_k whose components are irreducible components of the branched divisor D such that

$$kL_{\chi} = L_{\chi^k} + D_k.$$

Moreover, for $m = |H|$, we know that $mL_{\chi} = D_m$ and that $D_m - D$ is effective (see [6]). In particular, we have $D = p^*D_B$ for some ample divisor D_B on B .

Consider the Stein factorization of the natural morphism $\varphi = p \circ \rho : Y \rightarrow B$,

$$\varphi : Y \xrightarrow{\varphi'} Z \xrightarrow{\varphi''} B.$$

A general fiber of φ' is an étale cover of K , which is an abelian variety that we denote by \tilde{K} . Hence φ' is the litaka fibration of Y in the sense of [5, Theorem 13].

Let $Y' := Z \times_B A$. Then we have a factorization

$$\rho : Y \xrightarrow{\rho'} Y' \xrightarrow{\rho''} A.$$

Both ρ' and ρ'' are cyclic covers with respective Galois groups $H' \subset H$ and $H'' := H/H'$. Hence the morphism $\varphi'' : Z \rightarrow B$ is also an H'' -cover. Moreover, we write

$$\varphi''_* \mathcal{O}_Z = \bigoplus_{\iota \in H''^*} L_{\iota, Z}^{-1}.$$

Then we have $p^*L_{\iota, Z} = L_{\chi_{\iota}}$, where χ_{ι} is the induced character of H . Hence $h^i(B, L_{\iota, Z}) = 0$ for all $i \in \mathbb{Z}$ and non-trivial $\iota \in H''^*$.

Further, we show that Y' and Z have quotient singularities. Recall from [7, Proposition 2.8] that Y is the coarse moduli space of a canonical smooth Deligne–Mumford stack \mathcal{Y} . By the universal property of the canonical stack (see [4, Theorem 4.6]), the action of H' on Y induces an action on \mathcal{Y} . The quotient $Y' = Y/H'$ is the coarse moduli space of the smooth Deligne–Mumford stack $[\mathcal{Y}/H']$, and hence has quotient singularities.

We take a quotient $A \rightarrow K$ such that $A \rightarrow B \times K$ is an isogeny. Each connected component of a general fiber of the induced morphism $Y' \rightarrow K$ is an étale cover of Z . Hence Z also has quotient singularities.

Since $\dim Z < \dim Y$, the induction hypothesis implies that for all non-trivial $\iota \in H''^*$,

$$\sum_{h \in H''} \iota^{-1}(h)[\Gamma_h^Z] = 0 \in \text{CH}^{\dim Z}(Z \times Z)_{\mathbb{C}}.$$

Step 2. Proof that $\sum_{h \in H''} \iota^{-1}(h)[\Gamma_h^{Y'}] = 0 \in \text{CH}^{\dim Y'}(Y' \times Y')_{\mathbb{C}}$ for all non-trivial $\iota \in H''^*$.

Consider the quotient $A \rightarrow K$ such that $A \rightarrow B \times K$ is an isogeny with Galois group L . Then the induced morphism $\pi : Y' = Z \times_B A \rightarrow Z \times K$ is also an étale cover with Galois group L .

Note that for all $h \in H''$,

$$(\pi \times \pi)^*[\Gamma_h^Z \times \Delta_K] = \sum_{l \in L} [\Gamma_h^Z \times_{\Delta_B} \Gamma_l^A] \in \text{CH}^{\dim Y'}(Y' \times Y')_{\mathbb{Q}}.$$

Consider for each $l \in L$ the embeddings $\Gamma_h^Z \times_{\Delta_B} \Gamma_l^A \subset \Gamma_h^Z \times_{\Delta_B} (A \times_B A) \subset Y' \times Y'$. By Lemma 2.1, we have $[\Gamma_l^A] = [\Delta_A] \in \text{CH}_{\dim Y'}(A \times_B A)_{\mathbb{Q}}$. Hence

$$[\Gamma_h^Z \times_{\Delta_B} \Gamma_l^A] = [\Gamma_h^Z \times_{\Delta_B} \Delta_A] \in \text{CH}_{\dim Y'}(\Gamma_h^Z \times_{\Delta_B} (A \times_B A))_{\mathbb{Q}}.$$

In particular,

$$[\Gamma_h^Z \times_{\Delta_B} \Gamma_l^A] = [\Gamma_h^Z \times_{\Delta_B} \Delta_A] \in \text{CH}^{\dim Y'}(Y' \times Y')_{\mathbb{Q}}.$$

Then we find

$$\begin{aligned} (\pi \times \pi)^* \left(\sum_{h \in H''} \iota^{-1}(h)[\Gamma_h^Z \times \Delta_K] \right) &= \sum_{h \in H''} \iota^{-1}(h) \left(\sum_{l \in L} [\Gamma_h^Z \times_{\Delta_B} \Gamma_l^A] \right) \\ &= |L| \sum_{h \in H''} \iota^{-1}(h)[\Gamma_h^Z \times_{\Delta_B} \Delta_A] \\ &= |L| \sum_{h \in H''} \iota^{-1}(h)[\Gamma_h^{Y'}] \in \text{CH}^{\dim Y'}(Y' \times Y')_{\mathbb{C}}. \end{aligned}$$

As $\sum_{h \in H''} \iota^{-1}(h)[\Gamma_h^Z \times \Delta_K] = 0 \in \text{CH}^{\dim Y'}(Z \times Z \times K \times K)_{\mathbb{C}}$, we also have


$$\sum_{h \in H''} \iota^{-1}(h)[\Gamma_h^{Y'}] = 0 \in \text{CH}^{\dim Y'}(Y' \times Y')_{\mathbb{C}}.$$

Step 3. Proof that $\sum_{h \in H'} \chi^{-1}(h)[\Gamma_h^Y] = 0 \in \text{CH}^{\dim Y}(Y \times Y)_{\mathbb{C}}$ for all non-trivial $\chi \in H'^*$.

By Kawamata’s theorem [5, Theorem 13], there exists an étale cover $\mu : \tilde{Y} \rightarrow Y$ with Galois group J induced by an étale cover of A , such that the following holds: consider the Stein factorization of the natural morphism $\nu = \varphi' \circ \mu : \tilde{Y} \rightarrow Z$,

$$\nu : \tilde{Y} \xrightarrow{\nu'} \hat{Z} \xrightarrow{\nu''} Z.$$

Then we have $\tilde{Y} \simeq \hat{Z} \times \tilde{K}$ with ν' the projection, J acts on \hat{Z} faithfully and on \tilde{K} faithfully by translation, and $Y \simeq (\hat{Z} \times \tilde{K})/J$. We put everything into a commutative diagram:



Since μ is induced by an étale cover of A , we know that $\rho' \circ \mu : \tilde{Y} \rightarrow Y'$ is still an abelian cover. It is clear that the Galois group of $\rho \circ \mu$ is $J \times H$ and hence the Galois group of $\rho' \circ \mu$ is $J \times H'$. Let $q : J \times H' \rightarrow H'$ be the projection.

Consider the morphism $\mu \times \mu : \tilde{Y} \times \tilde{Y} \rightarrow Y \times Y$. Note that for all $h \in H'$,

$$(\mu \times \mu)^*[\Gamma_h^Y] = \sum_{g \in q^{-1}(h)} [\Gamma_g^{\tilde{Y}}] \in \text{CH}^{\dim \tilde{Y}}(\tilde{Y} \times \tilde{Y})_{\mathbb{Q}}.$$

Hence for all non-trivial $\chi \in H'^*$,

$$(\mu \times \mu)^* \left(\sum_{h \in H'} \chi^{-1}(h)[\Gamma_h^Y] \right) = \sum_{(g,h) \in J \times H'} \chi^{-1}(h)[\Gamma_{(g,h)}^{\tilde{Y}}] \in \text{CH}^{\dim Y}(\tilde{Y} \times \tilde{Y})_{\mathbb{C}}.$$

Under the isomorphism $\tilde{Y} \simeq \hat{Z} \times \tilde{K}$, we have

$$[\Gamma_{(g,h)}^{\tilde{Y}}] = [\Gamma_g^{\tilde{Z}} \times \Gamma_{(g,h)}^{\tilde{K}}] \in \text{CH}^{\dim Y}(\tilde{Z} \times \tilde{Z} \times \tilde{K} \times \tilde{K})_{\mathbb{Q}}.$$

Then we find

$$\begin{aligned} (\mu \times \mu)^* \left(\sum_{h \in H'} \chi^{-1}(h) [\Gamma_h^Y] \right) &= \sum_{(g,h) \in J \times H'} \chi^{-1}(h) [\Gamma_g^{\tilde{Z}} \times \Gamma_{(g,h)}^{\tilde{K}}] \\ &= \sum_{g \in J} ([\Gamma_g^{\tilde{Z}}] \times \left(\sum_{h \in H'} \chi^{-1}(h) [\Gamma_{(g,h)}^{\tilde{K}}] \right)) \in \text{CH}^{\dim Y}(\tilde{Z} \times \tilde{Z} \times \tilde{K} \times \tilde{K})_{\mathbb{C}}. \end{aligned}$$

By Lemma 2.1, we have $[\Gamma_{(g,h)}^{\tilde{K}}] = [\Delta_{\tilde{K}}] \in \text{CH}^{\dim \tilde{K}}(\tilde{K} \times \tilde{K})_{\mathbb{Q}}$. Hence

$$\sum_{h \in H'} \chi^{-1}(h) [\Gamma_{(g,h)}^{\tilde{K}}] = 0 \in \text{CH}^{\dim \tilde{K}}(\tilde{K} \times \tilde{K})_{\mathbb{C}},$$

and thus $(\mu \times \mu)^* \left(\sum_{h \in H'} \chi^{-1}(h) [\Gamma_h^Y] \right) = 0 \in \text{CH}^{\dim Y}(\tilde{Y} \times \tilde{Y})_{\mathbb{C}}$. By the projection formula,

$$\sum_{h \in H'} \chi^{-1}(h) [\Gamma_h^Y] = 0 \in \text{CH}^{\dim Y}(Y \times Y)_{\mathbb{C}}.$$

Step 4. Conclusion.

Let $\chi \in H^*$ be a non-trivial character. If the restriction of χ to H' is trivial, then χ is the pull-back of a non-trivial character ι of H'' . As before,

$$\sum_{h \in H} \chi^{-1}(h) [\Gamma_h^Y] = (\rho' \times \rho')^* \left(\sum_{h \in H''} \iota^{-1}(h) [\Gamma_h^{Y'}] \right) \in \text{CH}^{\dim Y}(Y \times Y)_{\mathbb{C}}.$$

By Step 2, we see that $\sum_{h \in H} \chi^{-1}(h) [\Gamma_h^Y] = 0 \in \text{CH}^{\dim Y}(Y \times Y)_{\mathbb{C}}$.

If the restriction of χ to H' is non-trivial, we take a subset V of H consisting of representatives of each coset of H' . Then we have

$$\sum_{h \in H} \chi^{-1}(h) [\Gamma_h^Y] = \left(\sum_{h \in H'} \chi^{-1}(h) [\Gamma_h^Y] \right) \circ \left(\sum_{g \in V} \chi^{-1}(g) [\Gamma_g^Y] \right) \in \text{CH}^{\dim Y}(Y \times Y)_{\mathbb{C}}.$$

By Step 3, we also know that $\sum_{h \in H} \chi^{-1}(h) [\Gamma_h^Y] = 0 \in \text{CH}^{\dim Y}(Y \times Y)_{\mathbb{C}}$. \square

We are now ready to prove Theorem 1.2. Let $\chi \in G^*$ be a non-trivial character. Consider

$$G' := \text{Ker}(\chi : G \rightarrow \mathbb{C}^*) \subset G.$$

We take $X' := X/G'$ which has quotient singularities (see Step 1 in the proof of Proposition 2.2), and we have a factorization

$$f : X \xrightarrow{f'} X' \xrightarrow{f''} A.$$

Then f'' is a cyclic cover with Galois group $G'' := G/G' \subset \mathbb{C}^*$. Moreover, the character χ is the pull-back of a non-trivial character ι of G'' . It is straightforward to verify that the assumption of Proposition 2.2 is satisfied by X' and G'' . Hence

$$\sum_{g \in G''} \iota^{-1}(g) [\Gamma_g^{X'}] = 0 \in \text{CH}^{\dim X'}(X' \times X')_{\mathbb{C}}.$$

As before,

$$\sum_{g \in G} \chi^{-1}(g) [\Gamma_g^X] = (f' \times f')^* \left(\sum_{g \in G''} \iota^{-1}(g) [\Gamma_g^{X'}] \right) = 0 \in \text{CH}^{\dim X}(X \times X)_{\mathbb{C}},$$

and thus

$$[\Gamma_X^X] = \frac{1}{|G|} \sum_{g \in G} \chi^{-1}(g) [\Gamma_g^X] = 0 \in \text{CH}^{\dim X}(X \times X)_{\mathbb{C}}. \quad \square$$

Acknowledgements

We thank Barbara Fantechi, Lie Fu, Johannes Schmitt, and Zhiyu Tian for helpful discussions.

References

- [1] A. Beauville, Sur l'anneau de Chow d'une variété abélienne, *Math. Ann.* 273 (4) (1986) 647–651.
- [2] Ch. Birkenhake, H. Lange, *Complex Abelian Varieties*, second edition, *Grundlehren der Mathematischen Wissenschaften*, vol. 302, Springer-Verlag, Berlin, 2004.
- [3] O. Debarre, Z. Jiang, M. Lahoz, W. Savin, Rational cohomology tori, *Geom. Topol.* 21 (2) (2017) 1095–1130.
- [4] B. Fantechi, E. Mann, F. Nironi, Smooth toric Deligne–Mumford stacks, *J. Reine Angew. Math.* 648 (2010) 201–244.
- [5] Y. Kawamata, Characterization of abelian varieties, *Compos. Math.* 43 (2) (1981) 253–276.
- [6] R. Pardini, Abelian covers of algebraic varieties, *J. Reine Angew. Math.* 417 (1991) 191–213.
- [7] A. Vistoli, Intersection theory on algebraic stacks and on their moduli spaces, *Invent. Math.* 97 (3) (1989) 613–670.