



Harmonic analysis

Time-frequency analysis on the adèles over the rationals

*Analyse des temps-fréquences sur l'anneau des adèles des rationnels*Ulrik B.R. Enstad^a, Mads S. Jakobsen^b, Franz Luef^b^a University of Oslo, Department of Mathematics, Oslo, Norway^b Norwegian University of Science and Technology, Department of Mathematical Sciences, Trondheim, Norway

ARTICLE INFO

Article history:

Received 27 July 2018

Accepted 7 January 2019

Available online 28 January 2019

Presented by the Editorial Board

ABSTRACT

We show that the construction of Gabor frames in $L^2(\mathbb{R})$ with generators in $\mathcal{S}_0(\mathbb{R})$ and with respect to time-frequency shifts from a rectangular lattice $\alpha\mathbb{Z} \times \beta\mathbb{Z}$ is equivalent to the construction of certain Gabor frames for L^2 over the adèles over the rationals and the group $\mathbb{R} \times \mathbb{Q}_p$. Furthermore, we detail the connection between the construction of Gabor frames on the adèles and on $\mathbb{R} \times \mathbb{Q}_p$ with the construction of certain Heisenberg modules.

© 2019 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Nous montrons que la construction de trames (ou repères) de Gabor de $L^2(\mathbb{R})$ avec générateurs dans $\mathcal{S}_0(\mathbb{R})$ et des décalages de temps-fréquence dans un réseau rectangulaire $\alpha\mathbb{Z} \times \beta\mathbb{Z}$ est équivalente à la construction de certaines trames de Gabor pour L^2 sur les adèles des rationnels avec le groupe $\mathbb{R} \times \mathbb{Q}_p$. Nous analysons également les relations entre la construction de trames de Gabor sur les adèles et sur $\mathbb{R} \times \mathbb{Q}_p$ et la construction de certains modules de Heisenberg.

© 2019 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

The theory of Gabor systems and of their frame properties is available on any locally compact abelian (LCA) group [23]. However, the construction of explicit examples of Gabor frames with time-frequency shifts from lattices (discrete and co-compact subgroups) is mostly restricted to the elementary LCA groups \mathbb{R} , \mathbb{Z} , \mathbb{T} and $\mathbb{Z}/d\mathbb{Z}$, $d \in \mathbb{N}$. While the group of the p -adic numbers \mathbb{Q}_p does not contain any lattice, the group $\mathbb{R} \times \mathbb{Q}_p$ and also the adèle group over the rationals, $\mathbb{A}_{\mathbb{Q}}$, do contain discrete and co-compact subgroups. This makes these groups eligible for Gabor analysis.

Other efforts to do time-frequency and time-scale (wavelet) analysis on local fields include [1–3,27,33,42,43] and [44].

We mention that [5] provides a method for constructing Gabor frames on any group with time-frequency shifts from lattices. In case of the p -adics or other groups that do not contain lattices, other methods of building somewhat structured families of functions with the use of quasi-lattices have been suggested in [19].

E-mail addresses: ubenstad@math.uio.no (U.B.R. Enstad), mads.jakobsen@ntnu.no (M.S. Jakobsen), franz.luef@ntnu.no (F. Luef).

In this paper, we combine the established theory on the p -adic numbers, the adèles, Gabor analysis, the theory of the Feichtinger algebra, and the modulation spaces. Our main result (Theorem 4.2) shows that the construction of (dual) Gabor frames for $L^2(\mathbb{R})$ with generators in the Feichtinger algebra $\mathbf{S}_0(\mathbb{R})$ is equivalent to both (I) the construction of certain (dual) Gabor frames for $L^2(\mathbb{A}_{\mathbb{Q}})$ with generators in $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})$ and (II) the construction of certain (dual) Gabor frames for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ with generators in $\mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$.

In Section 2, we describe the groups \mathbb{Q}_p and $\mathbb{A}_{\mathbb{Q}}$. The Feichtinger algebra on the p -adic groups \mathbb{Z}_p and \mathbb{Q}_p , the adèles $\mathbb{A}_{\mathbb{Q}}$, and $\mathbb{R} \times \mathbb{Q}_p$ is described in Section 3. Finally, in Section 4 we state and prove our main result. In Section 4.1, we connect the construction of dual Gabor frames in $L^2(\mathbb{A}_{\mathbb{Q}})$ and $L^2(\mathbb{R} \times \mathbb{Q}_p)$ with the construction of idempotent elements in twisted group C^* -algebras of the groups $\mathbb{A}_{\mathbb{Q}}$ and $\mathbb{R} \times \mathbb{Q}_p$ as explained in [24]. In this way, we answer the question raised in [30, Remark 5.8]. This work is related to the investigation of noncommutative solenoids by Latrémolière and Packer [30–32] and the work of Larsen and Li [29].

We also mention that after submitting the preprint of this paper, we have discovered a simultaneous generalization of the main result to Gabor frames over $L^2(\mathbb{R} \times \Omega_a)$, where Ω_a denotes the group of a -adic integers [20, p. 184], [21, Section 10]. These groups generalize both the p -adic numbers and the finite adèles over the rationals, but do not appear often in the literature. An example of an application of these groups in operator algebras can be found in [26].

2. The p -adic numbers and adèles over the rationals

In this section, we give a brief exposition of the field of p -adic numbers and the adèle group $\mathbb{A}_{\mathbb{Q}}$ over the rational numbers. Given a prime number p , the p -adic absolute value on \mathbb{Q} is defined by

$$|x|_p = p^{-k}$$

where $x = p^k(a/b)$ and p divides neither a nor b . One also sets $|0|_p = 0$. The p -adic absolute value satisfies a strengthened version of the triangle inequality (the ultrametric triangle inequality), namely

$$|x + y|_p \leq \max\{|x|_p, |y|_p\}. \tag{1}$$

The completion of \mathbb{Q} with respect to the metric $d_p(x, y) = |x - y|_p$ is a field denoted by \mathbb{Q}_p and its elements are called p -adic numbers. The topology inherited from the metric makes \mathbb{Q}_p into a locally compact Hausdorff space. In particular, the p -adic numbers \mathbb{Q}_p form a (non-compact) locally compact abelian group with respect to the topology induced by the above metric and under addition. One can show that every p -adic number x has a p -adic expansion of the form

$$x = \sum_{k=-\infty}^{\infty} a_k p^k,$$

where $a_k \in \{0, \dots, p - 1\}$ for each k and there exists some $n \in \mathbb{Z}$ such that $a_k = 0$ for all $k < n$. The sequence $(a_k)_{k \in \mathbb{Z}}$ in this expansion is unique.

The p -adic integers The closed unit ball in \mathbb{Q}_p is denoted by \mathbb{Z}_p and its elements are called p -adic integers. Because of (1) and the multiplicativity of $|\cdot|_p$, \mathbb{Z}_p is a subring of \mathbb{Q}_p . In terms of p -adic expansions, a p -adic number $x = \sum_{k \in \mathbb{Z}} a_k p^k$ is a p -adic integer if and only if $a_k = 0$ for $k < 0$. The map $\{0, \dots, p - 1\}^{\mathbb{N}} \rightarrow \mathbb{Z}_p$ given by $(a_k)_k \mapsto \sum_k a_k p^k$ is a homeomorphism, which shows that \mathbb{Z}_p has the topology of a Cantor set. In particular, \mathbb{Z}_p is a compact subgroup of \mathbb{Q}_p . But \mathbb{Z}_p is also open in \mathbb{Q}_p . Indeed, if $x \in \mathbb{Z}_p$, then using (1) one shows that the open ball $B_{1/2}(x) = \{y \in \mathbb{Q}_p : |y - x|_p < 1/2\}$ is contained in \mathbb{Z}_p .

We take the Haar measure $\mu_{\mathbb{Q}_p}$ on \mathbb{Q}_p so that $\mu_{\mathbb{Q}_p}(\mathbb{Z}_p) = 1$. The Haar measure on \mathbb{Z}_p is the one on \mathbb{Q}_p restricted to \mathbb{Z}_p .

The dual groups of \mathbb{Q}_p and \mathbb{Z}_p Denote by $\mathbb{Z}[1/p]$ the subring of \mathbb{Q} consisting of rational numbers of the form a/p^k where $k, a \in \mathbb{Z}$. Then $\mathbb{Q}_p = \mathbb{Z}_p + \mathbb{Z}[1/p]$ and $\mathbb{Z}_p \cap \mathbb{Z}[1/p] = \mathbb{Z}$, so that

$$\mathbb{Q}_p/\mathbb{Z}_p = \frac{\mathbb{Z}_p + \mathbb{Z}[1/p]}{\mathbb{Z}_p} \cong \frac{\mathbb{Z}[1/p]}{\mathbb{Z}_p \cap \mathbb{Z}[1/p]} = \mathbb{Z}[1/p]/\mathbb{Z}$$

as abelian groups. Denote the quotient map $\mathbb{Q}_p \rightarrow \mathbb{Z}[1/p]/\mathbb{Z}$ by $x \mapsto \{x\}_p$. In terms of p -adic expansions, we have $\{\sum_{k \in \mathbb{Z}} a_k p^k\}_p = \sum_{k=-\infty}^{-1} a_k p^k + \mathbb{Z}$ (observe that for any p -adic number x , only finitely many of the a_k are nonzero). Every character $\omega \in \widehat{\mathbb{Q}_p}$ is of the form

$$\omega : \mathbb{Q}_p \rightarrow \mathbb{C}, \omega(x) = e^{2\pi i \{xy\}_p}, x \in \mathbb{Q}_p,$$

for some $y \in \mathbb{Q}_p$. In fact, the map $\mathbb{Q}_p \rightarrow \widehat{\mathbb{Q}_p}$ given by mapping y to the ω defined above is a topological isomorphism. Hence the Pontryagin dual of \mathbb{Q}_p , $\widehat{\mathbb{Q}_p}$, can be identified with \mathbb{Q}_p itself. We will use this particular identification for the rest of the paper.

Recall that the *annihilator* of a closed subgroup H of a locally compact abelian group G is defined by

$$H^\perp = \{\omega \in \widehat{G} : \omega(h) = 1 \text{ for all } h \in H\}.$$

In general, we have topological isomorphisms $\widehat{H} \cong \widehat{G}/H^\perp$ and $\widehat{G/H} \cong H^\perp$. Under the identification of $\widehat{\mathbb{Q}_p}$ with \mathbb{Q}_p , the annihilator \mathbb{Z}_p^\perp of the p -adic integers \mathbb{Z}_p is identified with \mathbb{Z}_p itself. Hence the dual group $\widehat{\mathbb{Z}_p}$ is isomorphic to $\widehat{\mathbb{Q}_p}/\mathbb{Z}_p^\perp \cong \mathbb{Q}_p/\mathbb{Z}_p \cong \mathbb{Z}[1/p]/\mathbb{Z}$. There is another interesting description of $\widehat{\mathbb{Z}_p}$, namely it is isomorphic to the Prüfer p -group $\mathbb{Z}(p^\infty)$, which is the subgroup of \mathbb{T} consisting of all p^n -th roots of unity as n ranges over all $n = 0, 1, 2, \dots$,

$$\mathbb{Z}(p^\infty) = \{z \in \mathbb{C} : z = e^{2\pi i k p^{-n}}, k = 1, \dots, p^n, n \in \mathbb{N}_0\}. \tag{2}$$

Observe that any $z \in \mathbb{Z}(p^\infty)$ can be uniquely written as $z = \exp(2\pi i \sum_{k=1}^\infty a_k p^{-k})$, where each $a_k \in \{0, 1, \dots, p-1\}$ and only finitely many a_k are non-zero. This allows us to identify every element $z \in \mathbb{Z}(p^\infty)$ with a unique p -adic number, also to be denoted by z , given by $z = \sum_{k=1}^\infty a_k p^{-k}$. Every element $z \in \mathbb{Z}(p^\infty)$ defines a character $\omega_z \in \widehat{\mathbb{Z}_p}$ via

$$\omega_z : \mathbb{Z}_p \rightarrow \mathbb{C}, \omega_z(x) = e^{2\pi i \{xz\}_p}, x \in \mathbb{Z}_p. \tag{3}$$

Conversely, every element in $\widehat{\mathbb{Z}_p}$ is given in this way for some $z \in \mathbb{Z}(p^\infty)$.

Restricted products Let $(G_i)_{i \in I}$ be a collection of locally compact abelian groups for some index set I . If for each $i \in I$, the group H_i is a compact and open subgroup of G_i , then the *restricted product* of the groups $(G_i)_{i \in I}$ (with respect to the H_i) is defined to be

$$G = \prod_{i \in I}^{H_i} G_i = \left\{ (x_i)_{i \in I} \in \prod_{i \in I} G_i : x_i \in H_i \text{ for all but finitely many } i \in I \right\}.$$

The correct topology in order to turn G into a locally compact group is the one with a basis of open sets of the form $\prod_{i \in I} U_i$, where each U_i is open in G_i and $U_i = H_i$ for all but finitely many $i \in I$. This is called the *restricted product topology*. If each G_i is locally compact and each H_i is compact, then the restricted product is a locally compact group [38, Proposition 5-1(i)]. Moreover, every character $\omega \in \widehat{G}$ is of the form

$$\omega((x_i)_{i \in I}) = \prod_{i \in I} \omega_i(x_i),$$

where $(\omega_i)_{i \in I}$ is an element of the restricted product $\prod_{i \in I}^{H_i^\perp} \widehat{G}_i$, [38, Theorem 5-4]. This gives us an identification

$$\left(\prod_{i \in I}^{H_i} G_i \right)^\wedge \cong \prod_{i \in I}^{H_i^\perp} \widehat{G}_i.$$

If for each $i \in I$, the Haar measure on G_i is μ_i , then the product measure $\mu = \prod_{i \in I} \mu_i$ gives a Haar measure on the restricted product of the G_i . Typically, the Haar measure on each G_i is normalized so that $\mu_i(H_i) = 1$.

The adèles The group of *finite adèles* is the restricted product

$$\mathbb{A}_{\mathbb{Q}, \text{fin}} = \prod_{p \in \mathbb{P}}^{\mathbb{Z}_p} \mathbb{Q}_p.$$

As $\widehat{\mathbb{Q}_p} \cong \mathbb{Q}_p$ and $\mathbb{Z}_p^\perp \cong \mathbb{Z}_p$, it follows from the previous paragraph that $\mathbb{A}_{\mathbb{Q}, \text{fin}}$ is a self-dual locally compact abelian group. The group of *adèles* is

$$\mathbb{A}_{\mathbb{Q}} = \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}}.$$

It is also a self dual locally compact abelian group. We write elements of the adèles as $(x_\infty, (x_p)_p)$, where $x_\infty \in \mathbb{R}$ and $(x_p)_p \in \mathbb{A}_{\mathbb{Q}, \text{fin}}$. The Haar measure on $\mathbb{A}_{\mathbb{Q}}$ is the product measure of the Lebesgue measures on \mathbb{R} and the measure on all the p -adics \mathbb{Q}_p (normalized for each p so that $\mu_{\mathbb{Q}_p}(\mathbb{Z}_p) = 1$).

Every element $y = (y_\infty, (y_p)_p) \in \mathbb{A}_{\mathbb{Q}}$ defines a character $\omega_y \in \widehat{\mathbb{A}_{\mathbb{Q}}}$ via

$$\omega_y : \mathbb{A}_{\mathbb{Q}} \rightarrow \mathbb{T}, x = (x_\infty, (x_p)_p) \mapsto e^{2\pi i x_\infty y_\infty} \cdot \prod_{p \in \mathbb{P}} e^{-2\pi i \{x_p y_p\}_p}. \tag{4}$$

Moreover, every character on $\mathbb{A}_{\mathbb{Q}}$ is given in such a way by some $y \in \mathbb{A}_{\mathbb{Q}}$. The minus in the exponential for the p -adics is not necessary for this identification. It is however required for the following neat fact: we will see in a moment that the rationals \mathbb{Q} can be embedded into $\mathbb{A}_{\mathbb{Q}}$ as a discrete and co-compact subgroup. By the identification of $\mathbb{A}_{\mathbb{Q}}$ with $\widehat{\mathbb{A}_{\mathbb{Q}}}$ above, one has $\mathbb{Q}^\perp = \mathbb{Q}$. The identification in (4) is the same as that used in [7, Satz 5.4.2]. In [37, Section 4.3.7], the identification of $\mathbb{A}_{\mathbb{Q}}$ with $\widehat{\mathbb{A}_{\mathbb{Q}}}$ is done with the minus in the exponential for the real part.

Lattices in the adèles A subgroup H of an abelian topological group G is called a *lattice* if it is discrete and the quotient group G/H is compact. A *fundamental domain* for H in G is a Borel set $B \subseteq G$ such that every $x \in G$ can be uniquely written as $x = b + h$, where $b \in B$ and $h \in H$. Fundamental domains always exist for lattices in abelian groups [28, Lemma 2]. The size or *covolume* of a lattice H in G , denoted by $s(H)$, is the measure $\mu(B)$ of any fundamental domain B for H in G .

Naturally, there is an abundance of lattices in \mathbb{R} (they are all of the form $H = \alpha\mathbb{Z}$, $\alpha \in \mathbb{R} \setminus \{0\}$). The p -adic numbers contain *no* lattices (the only co-compact subgroup in \mathbb{Q}_p is \mathbb{Q}_p itself, and the only discrete subgroup of \mathbb{Q}_p is the trivial subgroup $\{0\}$). However, both $\mathbb{A}_{\mathbb{Q}}$ and $\mathbb{R} \times \mathbb{Q}_p$ do contain lattices.

Proposition 2.1. *For any $\alpha \in \mathbb{R} \setminus \{0\}$ the mapping*

$$\varphi_{\alpha} : \mathbb{Q} \rightarrow \mathbb{A}_{\mathbb{Q}}, \varphi_{\alpha}(q) = (\alpha q, (q)_p)$$

embeds \mathbb{Q} as a lattice into $\mathbb{A}_{\mathbb{Q}}$. The set $B_{\alpha} = [0, |\alpha|) \times \prod_{p \in \mathbb{P}} \mathbb{Z}_p$ is a fundamental domain for $\varphi_{\alpha}(\mathbb{Q})$ in $\mathbb{A}_{\mathbb{Q}}$ and $s(\varphi_{\alpha}(\mathbb{Q})) = |\alpha|$. Moreover, under the identification of $\mathbb{A}_{\mathbb{Q}}$ with $\widehat{\mathbb{A}}_{\mathbb{Q}}$ as in (4), the group $\varphi_{\alpha}(\mathbb{Q})^{\perp}$ can be identified with $\varphi_{1/\alpha}(\mathbb{Q})$.

Proof. As described in, e.g., [38, Theorem 5-11], [7, Satz 5.2.1], and [37, Section 4.3.7], $\varphi_1(\mathbb{Q})$ is a lattice in $\mathbb{A}_{\mathbb{Q}}$ with fundamental domain $[0, 1) \times \prod_{p \in \mathbb{P}} \mathbb{Z}_p$ and $s(\varphi_1(\mathbb{Q})) = 1$. Moreover, it is also shown that $\varphi_1(\mathbb{Q})^{\perp} = \varphi_1(\mathbb{Q})$. The embedding of \mathbb{Q} into $\mathbb{A}_{\mathbb{Q}}$ via φ_{α} for $\alpha \neq 1$ corresponds to an application of the topological group automorphism $x \mapsto \alpha \cdot x$, $x \in \mathbb{R}$ on the real component of $\mathbb{A}_{\mathbb{Q}}$. It is immediate that the desired results hold. \square

Lattices in $\mathbb{R} \times \mathbb{Q}_p$ In a fashion similar to the construction given in equation (4), every $y = (y_{\infty}, y_p) \in \mathbb{R} \times \mathbb{Q}_p$ defines a character $\omega_y \in \widehat{\mathbb{R} \times \mathbb{Q}_p}$ via

$$\omega_y : \mathbb{R} \times \mathbb{Q}_p, x = (x_{\infty}, x_p) \mapsto e^{2\pi i(x_{\infty}y_{\infty} - \{x_p y_p\}_p)}. \tag{5}$$

One can show that every character in $\widehat{\mathbb{R} \times \mathbb{Q}_p}$ is given as in (5) for some $y \in \mathbb{R} \times \mathbb{Q}_p$.

As for the adèles, there is an abundance of lattices in $\mathbb{R} \times \mathbb{Q}_p$. This is well known and can be found in, e.g., [31].

Proposition 2.2. *Let p be a prime number. For any $\alpha \in \mathbb{R} \setminus \{0\}$, the mapping*

$$\psi_{\alpha} : \mathbb{Z}[1/p] \rightarrow \mathbb{R} \times \mathbb{Q}_p, \psi_{\alpha}(q) = (\alpha q, q)$$

embeds $\mathbb{Z}[1/p]$ as a lattice into $\mathbb{R} \times \mathbb{Q}_p$. The set $B_{\alpha} = [0, |\alpha|) \times \mathbb{Z}_p$ is a fundamental domain for $\psi_{\alpha}(\mathbb{Z}[1/p])$ in $\mathbb{R} \times \mathbb{Q}_p$ and $s(\psi_{\alpha}(\mathbb{Z}[1/p])) = |\alpha|$. Moreover, under the identification of $\mathbb{R} \times \mathbb{Q}_p$ with $\widehat{\mathbb{R} \times \mathbb{Q}_p}$ as in (5), the group $\psi_{\alpha}(\mathbb{Z}[1/p])^{\perp}$ can be identified with $\psi_{1/\alpha}(\mathbb{Z}[1/p])$.

The proof is very similar in nature to that of Proposition 2.1 and it is therefore omitted.

3. Feichtinger’s algebra for p -adic groups and the adèles

For any locally compact abelian group, the *Feichtinger algebra* \mathbf{S}_0 [8,22,34] (sometimes denoted by \mathbf{M}^1) is a Banach space of functions that behaves very much like the Schwartz–Bruhat space \mathcal{S} (in fact, \mathcal{S} is a dense subspace of \mathbf{S}_0 , see [8, Theorem 9]). For example, \mathbf{S}_0 is invariant under the Fourier transform and the Poisson formula holds pointwise. In this section, we describe \mathbf{S}_0 for the following concrete groups: the real line \mathbb{R} , the p -adic integers \mathbb{Z}_p , the p -adic numbers \mathbb{Q}_p , the group $\mathbb{R} \times \mathbb{Q}_p$, the finite adèles $\mathbb{A}_{\mathbb{Q}, \text{fin}}$, and the adèles $\mathbb{A}_{\mathbb{Q}}$.

The description below of \mathbf{S}_0 on \mathbb{Q}_p and $\mathbb{A}_{\mathbb{Q}}$ makes it clear that \mathbf{S}_0 is a far larger and more interesting space of functions than the Schwartz–Bruhat space on these groups. The latter consists “only” of the collection of all locally constant functions.

\mathbf{S}_0 on the real line, \mathbb{R} The Feichtinger algebra on the real line, $\mathbf{S}_0(\mathbb{R})$, is described in detail in, e.g., [12] and [14]. Here we only mention the following definition. If we let g be the Gaussian function $g(x) = e^{-x^2}$, $x \in \mathbb{R}$, then

$$\mathbf{S}_0(\mathbb{R}) = \left\{ f \in L^1(\mathbb{R}) : \int_{\mathbb{R}} \|f * E_{\omega} g\|_1 d\omega < \infty \right\}.$$

Here $*$ is the usual convolution of functions and $E_{\omega} : f(x) \mapsto e^{2\pi i \omega x} f(x)$, $x \in \mathbb{R}$ is the modulation operator. The \mathbf{S}_0 -norm of f is given by $\|f\|_{\mathbf{S}_0(\mathbb{R})} = \int_{\mathbb{R}} \|f * E_{\omega} g\|_1 d\omega$.

\mathbf{S}_0 on the p -adic integers, \mathbb{Z}_p Since \mathbb{Z}_p is a compact group, it follows from, e.g., [8, Remark 3] or [22, Lemma 4.11], that $\mathbf{S}_0(\mathbb{Z}_p)$ consists exactly of all continuous functions on \mathbb{Z}_p that have an absolutely summable Fourier series. With the identification of the Prüfer p -group $\mathbb{Z}(p^\infty)$ as the dual group of \mathbb{Z}_p (see (2) and (3)), we can describe $\mathbf{S}_0(\mathbb{Z}_p)$ as follows:

$$\mathbf{S}_0(\mathbb{Z}_p) = \left\{ f \in \mathbf{C}(\mathbb{Z}_p) : f(x) = \sum_{z \in \mathbb{Z}(p^\infty)} c(z) e^{2\pi i \{xz\}_p}, x \in \mathbb{Z}_p \text{ and where } c \in \ell^1(\mathbb{Z}(p^\infty)) \right\}.$$

Moreover, the norm on $\mathbf{S}_0(\mathbb{Z}_p)$ is defined by $\|f\|_{\mathbf{S}_0(\mathbb{Z}_p)} = \|c\|_1$, where f and c are related as above.

\mathbf{S}_0 on the p -adic numbers, \mathbb{Q}_p The p -adic numbers \mathbb{Q}_p contain the p -adic integers \mathbb{Z}_p as a compact open subgroup. A set of coset representatives Y_p of $\mathbb{Q}_p/\mathbb{Z}_p$ is given by

$$Y_p = \{y \in \mathbb{Q}_p : y = \sum_{j=1}^{\infty} c_j p^{-j}, c_j \in \{0, 1, p-1\}, c_j = 0 \text{ for all but finitely many } j\}. \tag{6}$$

It follows by, e.g., [8, Lemma 8(ii)], [22, Theorem 7.7], or [39, §2.9] that $\mathbf{S}_0(\mathbb{Q}_p)$ consists exactly of all continuous functions f for which their restrictions to each of the cosets of $\mathbb{Q}_p/\mathbb{Z}_p$, the collection of functions

$$\{f_y : \mathbb{Z}_p \rightarrow \mathbb{C}, x \mapsto f(x+y), x \in \mathbb{Z}_p\}_{y \in Y_p},$$

belongs to $\mathbf{S}_0(\mathbb{Z}_p)$ and such that $\|f\|_{\mathbf{S}_0(\mathbb{Q}_p)} := \sum_{y \in Y_p} \|f_y\|_{\mathbf{S}_0(\mathbb{Z}_p)} < \infty$.

The characterization of $\mathbf{S}_0(\mathbb{Z}_p)$ in terms of the functions with absolutely convergent Fourier series allows us to describe $\mathbf{S}_0(\mathbb{Q}_p)$ as those functions that are exactly of the form

$$f(x) = \sum_{z \in \mathbb{Z}(p^\infty)} c(y, z) e^{2\pi i \{(x-y)z\}_p} \text{ for all } x \in y + \mathbb{Z}_p \tag{7}$$

and for all $y \in Y_p$, and where $c \in \ell^1(Y_p \times \mathbb{Z}(p^\infty))$. Moreover, the norm on $\mathbf{S}_0(\mathbb{Q}_p)$ is equivalently defined by $\|f\|_{\mathbf{S}_0(\mathbb{Q}_p)} = \|c\|_{\ell^1(Y_p \times \mathbb{Z}(p^\infty))}$, where f and c are related as in (7).

\mathbf{S}_0 on $\mathbb{R} \times \mathbb{Q}_p$ It follows from, e.g., [8, Theorem 7] or [22, Theorem 7.4], that the functions in $\mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$ are exactly those of the form

$$f = \sum_{j \in \mathbb{N}} f_j^{(\mathbb{R})} \otimes f_j^{(\mathbb{Q}_p)} \text{ where } f_j^{(\mathbb{R})} \in \mathbf{S}_0(\mathbb{R}), f_j^{(\mathbb{Q}_p)} \in \mathbf{S}_0(\mathbb{Q}_p) \tag{8}$$

for all $j \in \mathbb{N}$ and such that $\sum_{j \in \mathbb{N}} \|f_j^{(\mathbb{R})}\|_{\mathbf{S}_0(\mathbb{R})} \|f_j^{(\mathbb{Q}_p)}\|_{\mathbf{S}_0(\mathbb{Q}_p)} < \infty$. The norm on $\mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$ is given by

$$\|f\|_{\mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)} = \inf \left\{ \sum_{j \in \mathbb{N}} \|f_j^{(\mathbb{R})}\|_{\mathbf{S}_0(\mathbb{R})} \|f_j^{(\mathbb{Q}_p)}\|_{\mathbf{S}_0(\mathbb{Q}_p)} \right\},$$

where the functions $f, \{f_j^{(\mathbb{R})}\}_{j \in \mathbb{N}}$ and $\{f_j^{(\mathbb{Q}_p)}\}_{j \in \mathbb{N}}$ are related as in (8) and the infimum is taken over all possible representations of f as in (8).

\mathbf{S}_0 on $\mathbb{A}_{\mathbb{Q}, \text{fin}}$ The finite adeles $\mathbb{A}_{\mathbb{Q}, \text{fin}}$ contain $H = \pi_{p \in \mathbb{P}} \mathbb{Z}_p$ as a compact open subgroup. A set of coset representatives Y of $\mathbb{A}_{\mathbb{Q}, \text{fin}}/H$ is given by

$$Y = \left\{ (y_2, y_3, \dots) \in \pi_{p \in \mathbb{P}} Y_p : \text{where } Y_p \text{ is as in (6) and } y_p = 0 \text{ for all but finitely many } p \right\}.$$

By [8, Lemma 8(ii)], [22, Theorem 7.7], or [39, §2.9] the Banach space $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q}, \text{fin}})$ consists exactly of all continuous functions f on $\mathbb{A}_{\mathbb{Q}, \text{fin}}$ for which their restriction to each of the cosets of $\mathbb{A}_{\mathbb{Q}, \text{fin}}/H$, the functions

$$\{f_y : H \mapsto \mathbb{C}, x \mapsto f(x+y), x \in H\}_{y \in Y},$$

belong to $\mathbf{S}_0(H)$ and are such that

$$\|f\|_{\mathbf{S}_0(\mathbb{A}_{\mathbb{Q}, \text{fin}})} = \sum_{y \in Y} \|f_y\|_{\mathbf{S}_0(H)} < \infty. \tag{9}$$

Here $\mathbf{S}_0(H)$ is the Banach space of continuous functions over the compact group $H = \pi_{p \in \mathbb{P}} \mathbb{Z}_p$ with absolutely convergent Fourier series.

Another characterization of $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q},\text{fin}})$ is given in [39, §9.4] in the following way: any function f in $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q},\text{fin}})$ is exactly of the form

$$f = \sum_{j \in \mathbb{N}} \bigotimes_{p \in \mathbb{P}} f_j^{(\mathbb{Q}_p)} = \sum_{j \in \mathbb{N}} f_j^{(\mathbb{Q}_2)} \otimes f_j^{(\mathbb{Q}_3)} \otimes \dots, \text{ where } f_j^{(\mathbb{Q}_p)} \in \mathbf{S}_0(\mathbb{Q}_p), p \in \mathbb{P}, j \in \mathbb{N}, \tag{10}$$

and for each $j \in \mathbb{N}$ only finitely many of the functions $f_j^{(\mathbb{Q}_p)}$, $p \in \mathbb{P}$ are not equal to $\mathbb{1}_{\mathbb{Z}_p}$, and such that

$$\sum_{j \in \mathbb{N}} \prod_{p \in \mathbb{P}} \|f_j^{(\mathbb{Q}_p)}\|_{\mathbf{S}_0(\mathbb{Q}_p)} < \infty.$$

The norm

$$\|f\|_{\mathbf{S}_0(\mathbb{A}_{\mathbb{Q},\text{fin}})} = \inf \left\{ \sum_{j \in \mathbb{N}} \prod_{p \in \mathbb{P}} \|f_j^{(\mathbb{Q}_p)}\|_{\mathbf{S}_0(\mathbb{Q}_p)} \right\},$$

where the infimum is taken over all possible representations of f as in (10), is a norm on $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q},\text{fin}})$ that is equivalent to the norm in (9).

\mathbf{S}_0 on the adèle group, $\mathbb{A}_{\mathbb{Q}}$ By definition, $\mathbb{A}_{\mathbb{Q}} = \mathbb{R} \times \mathbb{A}_{\mathbb{Q},\text{fin}}$. It follows from [8, Theorem 7] or [22, Theorem 7.4] that a function f belongs to $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})$ if and only if

$$f = \sum_{j \in \mathbb{N}} \bigotimes_{p \in \{\infty\} \cup \mathbb{P}} f_j^{(\mathbb{Q}_p)} = \sum_{j \in \mathbb{N}} f_j^{(\mathbb{R})} \otimes f_j^{(\mathbb{Q}_2)} \otimes f_j^{(\mathbb{Q}_3)} \otimes \dots, \tag{11}$$

($\mathbb{Q}_{\infty} = \mathbb{R}$) where for each $j \in \mathbb{N}$ only finitely many of the functions $f_j^{(\mathbb{Q}_p)}$, $p \in \mathbb{P}$ are not equal to $\mathbb{1}_{\mathbb{Z}_p}$, and such that

$$\sum_{j \in \mathbb{N}} \prod_{p \in \{\infty\} \cup \mathbb{P}} \|f_j^{(\mathbb{Q}_p)}\|_{\mathbf{S}_0(\mathbb{Q}_p)} < \infty.$$

Moreover, the $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})$ -norm is given by

$$\|f\|_{\mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})} = \inf \left\{ \sum_{j \in \mathbb{N}} \prod_{p \in \{\infty\} \cup \mathbb{P}} \|f_j^{(\mathbb{Q}_p)}\|_{\mathbf{S}_0(\mathbb{Q}_p)} \right\},$$

where the infimum is taken over all possible representations of f as in (11).

4. Gabor frames

In this section, we describe how one can construct Gabor frames for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ and $L^2(\mathbb{A}_{\mathbb{Q}})$ from existing Gabor frames for $L^2(\mathbb{R})$. The theory of Gabor frames is well understood, see, e.g., the books [4,14] and the recent paper [23] that develops the theory of Gabor frames for general LCA groups. We give a very brief account of Gabor frames for general LCA groups before we state our main result, Theorem 4.2.

The theory of Gabor frames for general LCA groups For a moment, let G be a general locally compact abelian group. We denote the dual group by \widehat{G} . The action that an element $\omega \in \widehat{G}$ has on $x \in G$ is written as $\omega(x)$. For any $x \in G$ and $\omega \in \widehat{G}$, we define the translation T_x and the modulation operator E_{ω} as follows:

$$T_x f(t) = f(t - x), E_{\omega} f(t) = \omega(t) f(t), t \in G.$$

The translation and modulation operators are unitary operators on $L^2(G)$ and isometries on $\mathbf{S}_0(G)$. For convenience we define the time-frequency shift operator for any $\lambda = (x, \omega) \in G \times \widehat{G}$ to be

$$\pi(\lambda) = \pi(x, \omega) = E_{\omega} T_x.$$

Let Λ be a lattice (a discrete and co-compact subgroup) of the time-frequency domain $G \times \widehat{G}$ and let g be a function in $L^2(G)$. We let $\langle \cdot, \cdot \rangle$ denote the L^2 -inner product with the linearity in the first entry. The collection of functions $\{\pi(\lambda)g\}_{\lambda \in \Lambda} \subset L^2(G)$ is a Gabor system. Such a system is a frame for $L^2(G)$ if there exist constants $A, B > 0$ such that

$$A \|f\|_2^2 \leq \sum_{\lambda \in \Lambda} |\langle f, \pi(\lambda)g \rangle|^2 \leq B \|f\|_2^2 \text{ for all } f \in L^2(G).$$

Equivalently, the associated Gabor frame operator

$$S_{g,\Lambda} : L^2(G) \rightarrow L^2(G), S_{g,\Lambda} f = \sum_{\lambda \in \Lambda} \langle f, \pi(\lambda)g \rangle \pi(\lambda)g$$

is well defined, linear, bounded, and invertible. The usefulness of (Gabor) frames lies in the following. If $g \in L^2(G)$ and the lattice Λ in $G \times \widehat{G}$ are such that $\{\pi(\lambda)g\}_{\lambda \in \Lambda}$ is a frame for $L^2(G)$, then there exists a (in general non-unique) function $h \in L^2(G)$ such that

$$f = \sum_{\lambda \in \Lambda} \langle f, \pi(\lambda)g \rangle \pi(\lambda)h \text{ for all } f \in L^2(G). \tag{12}$$

If g and h are such that (12) holds, then they are called a dual pair of Gabor frame generators. For a given Gabor frame $\{\pi(\lambda)g\}_{\lambda \in \Lambda}$, the canonical choice of the function h such that (12) holds is the *canonical dual generator* $h = S_{g,\Lambda}^{-1}g$. It is a celebrated result of Gabor analysis that, if $g \in \mathbf{S}_0(G)$ generates a Gabor frame, then the canonical dual generator also belongs to $\mathbf{S}_0(G)$ [16].

For our purposes, we mention only the following result of Gabor analysis:

Lemma 4.1 ([23, Theorem 6.1]). *Let G be an LCA group and let Λ be a lattice of $G \times \widehat{G}$. Two functions $g, h \in \mathbf{S}_0(G)$ are a dual pair of Gabor frame generators for $L^2(G)$ with respect to time–frequency shifts from Λ (i.e., (12) holds) if and only if*

$$\langle h, \pi(\lambda^\circ)g \rangle = s(\Lambda) \delta_{\lambda^\circ, 0} \text{ for all } \lambda^\circ \in \Lambda^\circ,$$

where Λ° is the adjoint lattice of Λ ,

$$\Lambda^\circ = \{\lambda^\circ \in G \times \widehat{G} : \pi(\lambda) \pi(\lambda^\circ) = \pi(\lambda^\circ) \pi(\lambda) \text{ for all } \lambda \in \Lambda\}.$$

Gabor systems in $L^2(\mathbb{R})$ Recall that for every $\omega \in \mathbb{R}$ the modulation operator E_ω is given by $E_\omega f(t) = e^{2\pi i \omega t} f(t)$. A Gabor system in $L^2(\mathbb{R})$ generated by a function $g \in L^2(\mathbb{R})$ with time–frequency shifts from the lattice $\Lambda = \alpha\mathbb{Z} \times \beta\mathbb{Z}$, $\alpha, \beta > 0$ is thus of the form

$$\{\pi(\lambda)g\}_{\lambda \in \alpha\mathbb{Z} \times \beta\mathbb{Z}} = \{E_{m\beta} T_{n\alpha} g\}_{m,n \in \mathbb{Z}} = \{t \mapsto e^{2\pi i m \beta t} g(t - n\alpha)\}_{m,n \in \mathbb{Z}}.$$

Celebrated results in time–frequency analysis include the following: (A) the Gaussian function $g(x) = e^{-\pi x^2}$ [36,41] and all totally positive functions [18,17] generate a Gabor frame for $L^2(\mathbb{R})$ whenever α and β are such that $\alpha\beta < 1$; (B) the values of α , β and γ such that the Gabor system $\{E_{m\beta} T_{n\alpha} \mathbb{1}_{[0,\gamma]}\}_{m,n \in \mathbb{Z}}$ is a frame for $L^2(\mathbb{R})$ is much more difficult to describe [25,6].

Gabor systems in $L^2(\mathbb{R} \times \mathbb{Q}_p)$ For every $\omega = (\omega_\infty, \omega_p) \in \mathbb{R} \times \mathbb{Q}_p$, the modulation operator $E_\omega \equiv E_{\omega_\infty, \omega_p}$ on functions f over $\mathbb{R} \times \mathbb{Q}_p$ is defined by

$$E_\omega f(t_\infty, t_p) \equiv E_{\omega_\infty, \omega_p} f(t_\infty, t_p) = e^{2\pi i (\omega_\infty t_\infty - \{\omega_p t_p\}_p)} f(t_\infty, t_p), (t_\infty, t_p) \in \mathbb{R} \times \mathbb{Q}_p.$$

A Gabor system generated by a function $g \in L^2(\mathbb{R} \times \mathbb{Q}_p)$ and the lattice

$$\Lambda = \psi_\alpha(\mathbb{Z}[1/p]) \times \psi_\beta(\mathbb{Z}[1/p]) = \{(\alpha q, q, \beta r, r) : q, r \in \mathbb{Z}[1/p]\}, \alpha, \beta > 0$$

is thus of the form

$$\{\pi(\lambda)g\}_{\lambda \in \Lambda} = \{(t_\infty, t_p) \mapsto e^{2\pi i (\beta r t_\infty - \{r t_p\}_p)} g(t_\infty - \alpha q, t_p - q)\}_{q,r \in \mathbb{Z}[1/p]}.$$

Gabor systems in $L^2(\mathbb{A}_\mathbb{Q})$ For every $\omega = (\omega_\infty, (\omega_p)_p) \in \mathbb{A}_\mathbb{Q}$ the modulation operator $E_\omega \equiv E_{\omega_\infty, (\omega_p)_p}$ on functions f over $\mathbb{A}_\mathbb{Q}$ is defined by

$$E_{\omega_\infty, (\omega_p)_p} f(t_\infty, (t_p)_p) = e^{2\pi i \omega_\infty t_\infty} \prod_{p \in \mathbb{P}} e^{-2\pi i \{\omega_p t_p\}_p} f(t_\infty, (t_p)_p), (t_\infty, (t_p)_p) \in \mathbb{A}_\mathbb{Q}.$$

A Gabor system generated by a function $g \in L^2(\mathbb{A}_\mathbb{Q})$ and a lattice

$$\Lambda = \varphi_\alpha(\mathbb{Q}) \times \varphi_\beta(\mathbb{Q}) = \{(\alpha q, (q)_p, \beta r, (r)_p) : q, r \in \mathbb{Q}\}, \alpha, \beta > 0$$

is thus of the form

$$\{\pi(\lambda)g\}_{\lambda \in \Lambda} = \{(t_\infty, (t_p)_p) \mapsto e^{2\pi i \beta r t_\infty} \prod_{p \in \mathbb{P}} e^{-2\pi i \{r t_p\}_p} g(t_\infty - \alpha q, (t_p - q)_p)\}_{q,r \in \mathbb{Q}}.$$

Gabor frames in $L^2(\mathbb{R} \times \mathbb{Q}_p)$ and $L^2(\mathbb{A}_{\mathbb{Q}})$ The following result describes that the construction of a Gabor frame in $L^2(\mathbb{R})$ implies that certain functions generate Gabor frames for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ and $L^2(\mathbb{A}_{\mathbb{Q}})$.

Theorem 4.2. Let $\alpha, \beta > 0$. For any two functions $g^{(\mathbb{R})}$ and $h^{(\mathbb{R})}$ in $\mathbf{S}_0(\mathbb{R})$ the following statements are equivalent.

- (i) $g^{(\mathbb{R})}$ and $h^{(\mathbb{R})}$ generate dual Gabor frames for $L^2(\mathbb{R})$ with respect to time–frequency shifts from the lattice $\alpha\mathbb{Z} \times \beta\mathbb{Z}$.
- (ii) For any $p \in \mathbb{P}$, the two functions $g = g^{(\mathbb{R})} \otimes \mathbb{1}_{\mathbb{Z}_p}$ and $h = h^{(\mathbb{R})} \otimes \mathbb{1}_{\mathbb{Z}_p}$ in $\mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$ generate dual Gabor frames for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ with respect to the lattice

$$\Lambda = \psi_{\alpha}(\mathbb{Z}[1/p]) \times \psi_{\beta}(\mathbb{Z}[1/p]) = \{(\alpha q, q, \beta r, r) : q, r \in \mathbb{Z}[1/p]\} \subset \mathbb{R} \times \mathbb{Q}_p \times \mathbb{R} \times \mathbb{Q}_p.$$

- (iii) The two functions g and h in $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})$, defined by

$$g = g^{(\mathbb{R})} \otimes \mathbb{1}_{\mathbb{Z}_2} \otimes \mathbb{1}_{\mathbb{Z}_3} \otimes \dots \text{ and } h = h^{(\mathbb{R})} \otimes \mathbb{1}_{\mathbb{Z}_2} \otimes \mathbb{1}_{\mathbb{Z}_3} \otimes \dots,$$

generate dual Gabor frames for $L^2(\mathbb{A}_{\mathbb{Q}})$ with respect to the lattice

$$\Lambda = \varphi_{\alpha}(\mathbb{Q}) \times \varphi_{\beta}(\mathbb{Q}) = \{(\alpha q, (q)_p, \beta r, (r)_p) : q, r \in \mathbb{Q}\} \subset \mathbb{R} \times \mathbb{A}_{\mathbb{Q},\text{fin}} \times \mathbb{R} \times \mathbb{A}_{\mathbb{Q},\text{fin}}.$$

Corollary 4.3. For any $g^{(\mathbb{R})} \in \mathbf{S}_0(\mathbb{R})$ and $\alpha, \beta > 0$ the following statements are equivalent.

- (i) The function $g^{(\mathbb{R})}$ generates a Gabor frame for $L^2(\mathbb{R})$ with respect to the lattice $\alpha\mathbb{Z} \times \beta\mathbb{Z}$.
- (ii) For any $p \in \mathbb{P}$, the function $g = g^{(\mathbb{R})} \otimes \mathbb{1}_{\mathbb{Z}_p}$ generates a Gabor frame for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ with respect to the lattice

$$\Lambda = \psi_{\alpha}(\mathbb{Z}[1/p]) \times \psi_{\beta}(\mathbb{Z}[1/p]) = \{(\alpha q, q, \beta r, r) : q, r \in \mathbb{Z}[1/p]\} \subset \mathbb{R} \times \mathbb{Q}_p \times \mathbb{R} \times \mathbb{Q}_p.$$

- (iii) The function $g \in \mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})$ defined by

$$g = g^{(\mathbb{R})} \otimes \mathbb{1}_{\mathbb{Z}_2} \otimes \mathbb{1}_{\mathbb{Z}_3} \otimes \dots,$$

generates a Gabor frame for $L^2(\mathbb{A}_{\mathbb{Q}})$ with respect to the lattice

$$\Lambda = \varphi_{\alpha}(\mathbb{Q}) \times \varphi_{\beta}(\mathbb{Q}) = \{(\alpha q, (q)_p, \beta r, (r)_p) : q, r \in \mathbb{Q}\} \subset \mathbb{R} \times \mathbb{A}_{\mathbb{Q},\text{fin}} \times \mathbb{R} \times \mathbb{A}_{\mathbb{Q},\text{fin}}.$$

Proof of Theorem 4.2. It follows from the description of \mathbf{S}_0 in Section 3 that the functions g and h in (ii) and (iii) belong to $\mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$ and $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})$, respectively. We only proof the equivalence between (i) and (iii) as the proof of the equivalence between (i) and (ii) is almost identical. Observe that $s(\Lambda) = \alpha\beta$ and that the adjoint lattice to the lattice Λ in (iii) is the discrete and co-compact subgroup of $\mathbb{A}_{\mathbb{Q}} \times \mathbb{A}_{\mathbb{Q}}$ given by

$$\Lambda^{\circ} = \varphi_{1/\beta}(\mathbb{Q}) \times \varphi_{1/\alpha}(\mathbb{Q}) = \{(\beta^{-1}q, (q)_p, \alpha^{-1}r, (r)_p) : q, r \in \mathbb{Q}\} \subset \mathbb{R} \times \mathbb{A}_{\mathbb{Q},\text{fin}} \times \mathbb{R} \times \mathbb{A}_{\mathbb{Q},\text{fin}}.$$

By Lemma 4.1, the two functions g and h generate dual Gabor frames for $L^2(\mathbb{A}_{\mathbb{Q}})$ if and only if they satisfy

$$\langle h, \pi(\lambda^{\circ})g \rangle = \alpha\beta \delta_{\lambda^{\circ}, 0} \text{ for all } \lambda^{\circ} \in \Lambda^{\circ}. \tag{13}$$

The tensor product form of g and h implies that (13) takes the form

$$\langle h^{(\mathbb{R})}, E_{\alpha^{-1}r} T_{\beta^{-1}q} g^{(\mathbb{R})} \rangle \prod_{p \in \mathbb{P}} \langle \mathbb{1}_{\mathbb{Z}_p}, E_r T_q \mathbb{1}_{\mathbb{Z}_p} \rangle = \alpha\beta \delta_{(q,r), (0,0)} \text{ for all } q, r \in \mathbb{Q}. \tag{14}$$

Observe that a fraction $q \in \mathbb{Q}$ belongs to \mathbb{Z}_p for all $p \in \mathbb{P}$ if and only if $q \in \mathbb{Z}$. Hence, if $q \in \mathbb{Q} \setminus \mathbb{Z}$, then, for some $p \in \mathbb{P}$ the support of the function $\mathbb{1}_{\mathbb{Z}_p}$ and $T_q \mathbb{1}_{\mathbb{Z}_p}$ is disjoint. This implies that (14) is satisfied for all $q \in \mathbb{Q} \setminus \mathbb{Z}$. Since modulation E_r is turned into translation T_r by the Fourier transform on \mathbb{Q}_p , it follows from Parseval's identity that also, for any given $r \in \mathbb{Q} \setminus \mathbb{Z}$, the inner product

$$\langle \mathbb{1}_{\mathbb{Z}_p}, E_r \mathbb{1}_{\mathbb{Z}_p} \rangle_{L^2(\mathbb{Q}_p)} = 0$$

for some $p \in \mathbb{P}$. These two observations imply that we have verified (14) for all $q, r \in \mathbb{Q} \setminus \mathbb{Z}$. It remains to show that (14) holds for all $q, r \in \mathbb{Z}$. If $q, r \in \mathbb{Z}$, then $\langle \mathbb{1}_{\mathbb{Z}_p}, E_r T_q \mathbb{1}_{\mathbb{Z}_p} \rangle = 1$ for all $p \in \mathbb{P}$ (we have normalized the Haar measure on each \mathbb{Q}_p such that $\int_{\mathbb{Q}_p} \mathbb{1}_{\mathbb{Z}_p} = 1$). This implies that we only need to verify

$$\langle h^{(\mathbb{R})}, E_{\alpha^{-1}r} T_{\beta^{-1}q} g^{(\mathbb{R})} \rangle = \alpha\beta \delta_{(q,r), (0,0)} \text{ for all } q, r \in \mathbb{Z}. \tag{15}$$

Lemma 4.1 states that (15) is satisfied if and only if the two Gabor systems $\{E_{m\beta} T_{n\alpha} g^{(\mathbb{R})}\}_{m,n \in \mathbb{Z}}$ and $\{E_{m\beta} T_{n\alpha} h^{(\mathbb{R})}\}_{m,n \in \mathbb{Z}}$ are dual Gabor frames, which is (i). \square

A Balian–Low theorem The classical Balian–Low theorem for the Feichtinger algebra on the real numbers states that if $g \in \mathbf{S}_0(\mathbb{R})$ and $\alpha\beta = 1$, then g cannot generate a Gabor frame for $L^2(\mathbb{R})$ over the lattice $\alpha\mathbb{Z} \times \beta\mathbb{Z} \subseteq \mathbb{R} \times \widehat{\mathbb{R}}$. It is a natural question to ask whether the following general statement for locally compact abelian groups holds: suppose that $g \in \mathbf{S}_0(G)$ and that Λ is a lattice in $G \times \widehat{G}$ of the form $\Gamma \times \Gamma^\perp$ with Γ a lattice in G . Then g should not generate a Gabor frame for $L^2(G)$ over Λ ?

It turns out that this does not hold for general LCA groups G and lattices Γ , see [13]. However, it has been shown to hold for all lattices in second countable compactly generated LCA groups with noncompact component of the identity [28]. The groups $\mathbb{A}_\mathbb{Q}$ and $\mathbb{R} \times \mathbb{Q}_p$ are both second countable with noncompact component of the identity, but not compactly generated, so the result in [28] does not cover these groups. Note that the lattice $\Gamma = \varphi_\alpha(\mathbb{Q})$ in $\mathbb{A}_\mathbb{Q}$ has annihilator $\Gamma^\perp = \varphi_{1/\alpha}(\mathbb{Q})$. Thus, if $\alpha\beta = 1$, then $\varphi_\alpha(\mathbb{Q}) \times \varphi_\beta(\mathbb{Q}) \subseteq \mathbb{A}_\mathbb{Q} \times \mathbb{A}_\mathbb{Q}$ is a lattice of the form $\Gamma \times \Gamma^\perp$ (and similarly for the lattice $\psi_\alpha(\mathbb{Z}[1/p]) \times \psi_\beta(\mathbb{Z}[1/p]) \subseteq (\mathbb{R} \times \mathbb{Q}_p) \times (\mathbb{R} \times \mathbb{Q}_p)$).

Combining Corollary 4.3 and the classical Balian–Low theorem for the Feichtinger algebra on \mathbb{R} , we obtain the following restricted Balian–Low type theorem for the groups $\mathbb{A}_\mathbb{Q}$ and $\mathbb{R} \times \mathbb{Q}_p$:

Proposition 4.4. *Let $g \in \mathbf{S}_0(\mathbb{R})$, and let $\alpha\beta = 1$. Then the following hold:*

- (i) *The function $g \otimes \mathbb{1}_{\mathbb{Z}_2} \otimes \mathbb{1}_{\mathbb{Z}_3} \otimes \dots \in \mathbf{S}_0(\mathbb{A}_\mathbb{Q})$ does not generate a Gabor frame for $L^2(\mathbb{A}_\mathbb{Q})$ over the lattice $\varphi_\alpha(\mathbb{Q}) \times \varphi_\beta(\mathbb{Q})$ in $\mathbb{A}_\mathbb{Q} \times \mathbb{A}_\mathbb{Q}$.*
- (ii) *The function $g \otimes \mathbb{1}_{\mathbb{Z}_p} \in \mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$ does not generate a Gabor frame for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ over the lattice $\psi_\alpha(\mathbb{Z}[1/p]) \times \psi_\beta(\mathbb{Z}[1/p])$ in $(\mathbb{R} \times \mathbb{Q}_p) \times (\mathbb{R} \times \mathbb{Q}_p)$.*

The above result only holds for functions in $\mathbf{S}_0(\mathbb{A}_\mathbb{Q})$ of the restricted form $g \otimes \mathbb{1}_{\mathbb{Z}_2} \otimes \mathbb{1}_{\mathbb{Z}_3} \otimes \dots$ with $g \in \mathbf{S}_0(\mathbb{R})$ (and analogously for $\mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$). An interesting question is whether one can obtain the same conclusion for all elements of $\mathbf{S}_0(\mathbb{A}_\mathbb{Q})$, and the first author is presently working on this.

Modulation spaces Modulation spaces were invented by Feichtinger in the early 1980s and can be defined on any locally compact abelian group, see, e.g., [9–11] and [14].

It is well known that the modulation spaces can be described using Gabor frames that are constructed with windows in \mathbf{S}_0 [14]. This characterization of the modulation spaces and the construction of the Gabor frames for $L^2(\mathbb{R} \times \mathbb{Q}_p)$ and $L^2(\mathbb{A}_\mathbb{Q})$ in Theorem 4.2 lead to the following.

Lemma 4.5. *Let $g \in \mathbf{S}_0(\mathbb{A}_\mathbb{Q})$ and Λ be as in Theorem 4.2(iii) such that $\{\pi(\lambda)g\}_{\lambda \in \Lambda}$ is a Gabor frame for $L^2(\mathbb{A}_\mathbb{Q})$. The modulation space $M^{s,t}(\mathbb{A}_\mathbb{Q})$, $s, t \in [1, \infty]$ consists exactly of all elements $\sigma \in \mathbf{S}'_0(\mathbb{A}_\mathbb{Q})$ such that*

$$\|\sigma\|_{M^{s,t}(\mathbb{A}_\mathbb{Q})} := \left(\sum_{r \in \mathbb{Q}} \left(\sum_{q \in \mathbb{Q}} |\sigma(E_{\beta r, (r)_p} T_{\alpha q, (q)_p} g)|^s \right)^{t/s} \right)^{1/t},$$

with the obvious modification if s or t equal ∞ .

In a similar way, the modulation spaces on $\mathbb{R} \times \mathbb{Q}_p$ can be defined using the Gabor frames constructed as in Theorem 4.2(ii).

It is well known that $\mathbf{S}_0 \cong M^{1,1}$, $L^2 \cong M^{2,2}$, and that $\mathbf{S}'_0 = M^{\infty,\infty}$.

In recent years, the modulation spaces have been used successfully as spaces of symbols in the theory of pseudo-differential operators. For example, the space $M^{\infty,1}$ coincides with the *Sjöstrand class*. Among many we refer to, e.g., [15, 19].

4.1. Heisenberg modules

As described in [24], the construction of Gabor frames for $L^2(G)$ with time-frequency shifts from a closed subgroup $\Lambda \subset G \times \widehat{G}$ (where G is any locally compact abelian group and \widehat{G} its dual group) is equivalent to the construction of certain projections in the twisted group C^* -algebra $C^*(\Lambda, c)$ where c denotes the cocycle coming from the Heisenberg representation [40]. We state briefly some of the theory of [24] for the case of the Gabor frames for $L^2(\mathbb{A}_\mathbb{Q})$ and $L^2(\mathbb{R} \times \mathbb{Q}_p)$ constructed in Theorem 4.2. This is the first example of a singly-generated Heisenberg module beyond the case of elementary locally compact abelian groups and it is not covered by the recent results in [24]. The equivalence bimodule that we use here is a suitable completion of the Feichtinger algebra on the respective groups. In contrast, the theory presented in [40] and [30,31] use completions of the Schwartz–Bruhat space $\mathcal{S}(G)$ and $C_c(G)$, respectively, to construct equivalence bimodules between twisted group C^* -algebras of lattices in $G \times \widehat{G}$.

Heisenberg modules over $\mathbb{R} \times \mathbb{Q}_p$ For $\alpha, \beta > 0$ we define the following two Banach algebras:

$$\mathcal{A} = \left\{ \mathbf{a} \in \mathbf{B}(L^2(\mathbb{R} \times \mathbb{Q}_p)) : \mathbf{a} = \sum_{q,r \in \mathbb{Z}[1/p]} a(q,r) E_{\beta r,r} T_{\alpha q,q}, a \in \ell^1(\mathbb{Z}[1/p]^2) \right\},$$

$$\mathcal{B} = \left\{ \mathbf{b} \in \mathbf{B}(L^2(\mathbb{R} \times \mathbb{Q}_p)) : \mathbf{b} = \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Z}[1/p]} b(q,r) (E_{\alpha^{-1}r,r} T_{\beta^{-1}q,q})^*, b \in \ell^1(\mathbb{Z}[1/p]^2) \right\}.$$

Indeed, the norms $\|\mathbf{a}\|_{\mathcal{A}} = \|a\|_1$, $\|\mathbf{b}\|_{\mathcal{B}} = \|b\|_1$ (where $\mathbf{a}, a, \mathbf{b}$ and b are related as above) turn \mathcal{A} and \mathcal{B} into involutive Banach algebras with respect to the composition of operators and where the involution is the L^2 -adjoint.

Observe that \mathcal{A} and \mathcal{B} are *not* generated by finitely many unitaries as is the case of the noncommutative 2-torus studied in relation to the construction of Gabor frames in $L^2(\mathbb{R})$ [35].

Elements in \mathcal{A} and \mathcal{B} act on functions in $L^2(\mathbb{R} \times \mathbb{Q}_p)$ from the left and the right, respectively, by

$$\mathbf{a} \cdot f := \sum_{q,r \in \mathbb{Z}[1/p]} a(q,r) E_{\beta r,r} T_{\alpha q,q} f, f \in L^2(\mathbb{R} \times \mathbb{Q}_p), \mathbf{a} \in \mathcal{A},$$

$$f \cdot \mathbf{b} := \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Z}[1/p]} b(q,r) (E_{\alpha^{-1}r,r} T_{\beta^{-1}q,q})^* f, f \in L^2(\mathbb{R} \times \mathbb{Q}_p), \mathbf{b} \in \mathcal{B}.$$

We define an \mathcal{A} - and \mathcal{B} -valued inner product in the following way:

$$\mathcal{A}(\cdot, \cdot) : \mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p) \times \mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p) \rightarrow \mathcal{A},$$

$$\mathcal{A}(f, g) = \sum_{q,r \in \mathbb{Z}[1/p]} \langle f, E_{\beta r,r} T_{\alpha q,q} g \rangle E_{\beta r,r} T_{\alpha q,q},$$

$$\langle \cdot, \cdot \rangle_{\mathcal{B}} : \mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p) \times \mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p) \rightarrow \mathcal{B},$$

$$\langle f, g \rangle_{\mathcal{B}} = \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Z}[1/p]} \langle g, (E_{\alpha^{-1}r,r} T_{\beta^{-1}q,q})^* f \rangle (E_{\alpha^{-1}r,r} T_{\beta^{-1}q,q})^*.$$

One can show that

$$\mathcal{A}(f, g) \cdot h = f \cdot \langle g, h \rangle_{\mathcal{B}} \text{ for all } f, g, h \in \mathbf{S}_0(\mathbb{R} \times \mathbb{A}_{\mathbb{Q}}).$$

Denote by A and B the C^* -closures of \mathcal{A} and \mathcal{B} inside $\mathbf{B}(L^2(\mathbb{R} \times \mathbb{Q}_p))$, respectively. The actions and algebra-valued inner products defined give $\mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$, the structure of a pre-imprimitivity \mathcal{A} - \mathcal{B} -bimodule. It can thus be completed into an imprimitivity A - B bimodule, a *Heisenberg module* in the sense of Rieffel [40], which sets up a Morita equivalence between A and B . It is worth noting that, in this case, A is a twisted group C^* -algebra on the group $\mathbb{Z}[1/p] \times \mathbb{Z}[1/p]$. These have been termed *noncommutative solenoids* by F. Latrémolière and J. Packer and are studied in [30–32], where they prove that B is also a noncommutative solenoid.

Proposition 4.6 ([24, Theorem 3.14]). *Let g, h be two functions in $\mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$ and consider the lattices*

$$\Lambda = \{(\alpha q, q, \beta r, r) : q, r \in \mathbb{Z}[1/p]\} \text{ and } \Lambda^\circ = \{(\beta^{-1}q, q, \alpha^{-1}r, r) : q, r \in \mathbb{Z}[1/p]\}$$

in $(\mathbb{R} \times \mathbb{Q}_p)^2$ with $\alpha, \beta > 0$ as in Theorem 4.2. The following statements are equivalent:

- (i) $f = \mathcal{A}(f, g) \cdot h$ for all $f \in \mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$;
- (ii) $\langle g, h \rangle_{\mathcal{B}}$ is the identity operator on $L^2(\mathbb{R} \times \mathbb{Q}_p)$;
- (iii) g and h generate dual Gabor frames with respect to Λ for $L^2(\mathbb{R} \times \mathbb{Q}_p)$;
- (iv) $\mathcal{A}(g, h)$ is an idempotent operator from $L^2(\mathbb{R} \times \mathbb{Q}_p)$ onto $V := \overline{\text{span}}\{\pi(\lambda^\circ)g\}_{\lambda^\circ \in \Lambda^\circ}$;
- (v) $f = g \cdot \langle h, f \rangle_{\mathcal{B}}$ for all $f \in \mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p) \cap V$.

We close with a result on projections in \mathcal{A} which follows from Theorem 4.2 by choosing for $g^{(\mathbb{R})}$ the Gaussian $g_0^{(\mathbb{R})}(t) = e^{-\pi t^2}$.

Proposition 4.7. *Let $g \in \mathbf{S}_0(\mathbb{R} \times \mathbb{Q}_p)$ be the function defined by $g = g_0^{(\mathbb{R})} \otimes \mathbb{1}_{\mathbb{Z}_p}$, where $g_0^{(\mathbb{R})}$ is the Gaussian, and consider the lattice*

$$\Lambda = \varphi_\alpha(\mathbb{Z}[1/p]) \times \varphi_\beta(\mathbb{Z}[1/p]) = \{(\alpha q, q, \beta r, r) : q, r \in \mathbb{Z}[1/p]\} \subset \mathbb{R} \times \mathbb{Q}_p \times \mathbb{R} \times \mathbb{Q}_p.$$

Then $\mathcal{A}(S_{g,\Lambda}^{-1/2}g, S_{g,\Lambda}^{-1/2}g)$ is a projection in \mathcal{A} if and only if $\alpha\beta < 1$.

Proof. As shown in [35] the construction of projections of the form $\mathcal{A}\langle g, g \rangle$ is equivalent to the construction of tight Gabor frames. Since $S_{g,\Lambda}^{-1/2}g$ generates the canonical tight Gabor frame the result follows from the result of Lyubarskii-Seip that $\{\pi(\alpha k, \beta l)g_0^{(\mathbb{R})}\}_{k,l \in \mathbb{Z}}$ is a Gabor frame if and only if $\alpha\beta < 1$. \square

Heisenberg modules over $\mathbb{A}_{\mathbb{Q}}$ For $\alpha, \beta > 0$ we define the following two Banach algebras:

$$\mathcal{A} = \left\{ \mathbf{a} \in \mathbf{B}(L^2(\mathbb{A}_{\mathbb{Q}})) : \mathbf{a} = \sum_{q,r \in \mathbb{Q}} a(q,r) E_{\beta r, (r)_p} T_{\alpha q, (q)_p}, a \in \ell^1(\mathbb{Q}^2) \right\},$$

$$\mathcal{B} = \left\{ \mathbf{b} \in \mathbf{B}(L^2(\mathbb{A}_{\mathbb{Q}})) : \mathbf{b} = \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Q}} b(q,r) (E_{\alpha^{-1}r, (r)_p} T_{\beta^{-1}q, (q)_p})^*, b \in \ell^1(\mathbb{Q}^2) \right\}.$$

The norms $\|\mathbf{a}\|_{\mathcal{A}} = \|a\|_1, \|\mathbf{b}\|_{\mathcal{B}} = \|b\|_1$ turn \mathcal{A} and \mathcal{B} into involutive Banach algebras with respect to the composition of operators and where the involution is the L^2 -adjoint. As in the case of $\mathbb{R} \times \mathbb{Q}_p$ described before, \mathcal{A} and \mathcal{B} are not generated by finitely many unitaries. Elements in \mathcal{A} and \mathcal{B} act on functions in $L^2(\mathbb{A}_{\mathbb{Q}})$ from the left and the right, respectively, by

$$\mathbf{a} \cdot f := \sum_{q,r \in \mathbb{Q}} a(q,r) E_{\beta r, (r)_p} T_{\alpha q, (q)_p} f, f \in L^2(\mathbb{A}_{\mathbb{Q}}), \mathbf{a} \in \mathcal{A},$$

$$f \cdot \mathbf{b} := \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Q}} b(q,r) (E_{\alpha^{-1}r, (r)_p} T_{\beta^{-1}q, (q)_p})^* f, f \in L^2(\mathbb{A}_{\mathbb{Q}}), \mathbf{b} \in \mathcal{B}.$$

We define an \mathcal{A} - and \mathcal{B} -valued inner product in the following way:

$$\mathcal{A}\langle \cdot, \cdot \rangle : \mathbf{S}_0(\mathbb{A}_{\mathbb{Q}}) \times \mathbf{S}_0(\mathbb{A}_{\mathbb{Q}}) \rightarrow \mathcal{A},$$

$$\mathcal{A}\langle f, g \rangle = \sum_{q,r \in \mathbb{Q}} \langle f, E_{\beta r, (r)_p} T_{\alpha q, (q)_p} g \rangle E_{\beta r, (r)_p} T_{\alpha q, (q)_p},$$

$$\langle \cdot, \cdot \rangle_{\mathcal{B}} : \mathbf{S}_0(\mathbb{A}_{\mathbb{Q}}) \times \mathbf{S}_0(\mathbb{A}_{\mathbb{Q}}) \rightarrow \mathcal{B},$$

$$\langle f, g \rangle_{\mathcal{B}} = \frac{1}{\alpha\beta} \sum_{q,r \in \mathbb{Q}} \langle g, (E_{\alpha^{-1}r, (r)_p} T_{\beta^{-1}q, (q)_p})^* f \rangle (E_{\alpha^{-1}r, (r)_p} T_{\beta^{-1}q, (q)_p})^*.$$

One can show that

$$\mathcal{A}\langle f, g \rangle \cdot h = f \cdot \langle g, h \rangle_{\mathcal{B}} \text{ for all } f, g, h \in \mathbf{S}_0(\mathbb{A}_{\mathbb{Q}}).$$

In this case, as in the case with the group $\mathbb{R} \times \mathbb{Q}_p$, we obtain from $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})$ an imprimitivity A - B bimodule, where A and B denote the C^* -closures of \mathcal{A} and \mathcal{B} in $\mathbf{B}(L^2(\mathbb{A}_{\mathbb{Q}}))$, respectively.

Proposition 4.8 ([24, Theorem 3.14]). *Let g, h be two functions in $\mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})$ and consider the lattices*

$$\Lambda = \{(\alpha q, (q)_p, \beta r, (r)_p) : q, r \in \mathbb{Q}\} \text{ and } \Lambda^\circ = \{(\beta^{-1}q, (q)_p, \alpha^{-1}r, (r)_p) : q, r \in \mathbb{Q}\}$$

in $\mathbb{A}_{\mathbb{Q}}^2$ with $\alpha, \beta > 0$ as in Theorem 4.2. The following statements are equivalent:

- (i) $f = \mathcal{A}\langle f, g \rangle \cdot h$ for all $f \in \mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})$;
- (ii) $\langle g, h \rangle_{\mathcal{B}}$ is the identity operator on $L^2(\mathbb{A}_{\mathbb{Q}})$;
- (iii) g and h generate dual Gabor frames with respect to Λ for $L^2(\mathbb{A}_{\mathbb{Q}})$;
- (iv) $\mathcal{A}\langle g, h \rangle$ is an idempotent operator from $L^2(\mathbb{A}_{\mathbb{Q}})$ onto $V := \overline{\text{span}}\{\pi(\lambda^\circ)g\}_{\lambda^\circ \in \Lambda^\circ}$;
- (v) $f = g \cdot \langle h, f \rangle_{\mathcal{B}}$ for all $f \in \mathbf{S}_0(\mathbb{A}_{\mathbb{Q}}) \cap V$.

We close with a result on projections in \mathcal{A} which follows from Theorem 4.2 by choosing for $g^{(\mathbb{R})}$ the Gaussian $g_0^{(\mathbb{R})}(t) = e^{-\pi t^2}$. The proof is analogous to the one of Proposition 4.7.

Proposition 4.9. *Let $g \in \mathbf{S}_0(\mathbb{A}_{\mathbb{Q}})$ be the function defined by*

$$g = g_0^{(\mathbb{R})} \otimes \mathbb{1}_{\mathbb{Z}_2} \otimes \mathbb{1}_{\mathbb{Z}_3} \otimes \dots,$$

where $g_0^{(\mathbb{R})}$ denotes the Gaussian, and consider the lattice

$$\Lambda = \varphi_\alpha(\mathbb{Q}) \times \varphi_\beta(\mathbb{Q}) = \{(\alpha q, (q)_p, \beta r, (r)_p) : q, r \in \mathbb{Q}\} \subset \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}} \times \mathbb{R} \times \mathbb{A}_{\mathbb{Q}, \text{fin}}.$$

Then $\mathcal{A}\langle S_{g,\Lambda}^{-1/2}g, S_{g,\Lambda}^{-1/2}g \rangle$ is a projection in \mathcal{A} if and only if $\alpha\beta < 1$.

Acknowledgements

The work of M.S.J. was carried out during the tenure of the ERCIM “Alain Bensoussan” Fellowship Programme at NTNU. This work was finished while M.S.J. and F.L. were visiting NuHAG at the Faculty of Mathematics at the University of Vienna. We are thankful for their hospitality. The first author thanks Nadia Larsen and Tron Omland for valuable discussions about the groups $\mathbb{A}_{\mathbb{Q}}$ and $\mathbb{R} \times \mathbb{Q}_p$ and for pointing out that these groups admit lattices.

References

- [1] O. Ahmad, F.A. Shah, N.A. Sheik, Gabor frames on non-Archimedean fields, *Int. J. Geom. Methods Mod. Phys.* 15 (05) (2018) 1850079.
- [2] S. Albeverio, S. Evdokimov, M. Skopina, p -adic multiresolution analysis and wavelet frames, *Fourier Anal. Appl.* 16 (5) (2010) 693–714.
- [3] J.J. Benedetto, R.L. Benedetto, *Frames of translates for number-theoretic groups*, ArXiv e-prints, 2018.
- [4] O. Christensen, *An Introduction to Frames and Riesz Bases*, Applied and Numerical Harmonic Analysis, second edition, Birkhäuser, Basel, 2016.
- [5] O. Christensen, S.S. Goh, Fourier-like frames on locally compact Abelian groups, *J. Approx. Theory* 192 (2015) 82–101.
- [6] X.-R. Dai, Q. Sun, The abc -problem for Gabor systems, *Mem. Amer. Math. Soc.* 244 (1152) (2016), ix+99.
- [7] A. Deitmar, *Automorphe Formen*, Springer, Heidelberg, Germany, 2010.
- [8] H.G. Feichtinger, On a new Segal algebra, *Monatshefte Math.* 92 (1981) 269–289.
- [9] H.G. Feichtinger, *Modulation Spaces on Locally Compact Abelian Groups*, Technical report, University of Vienna, January 1983.
- [10] H.G. Feichtinger, Modulation spaces of locally compact Abelian groups, in: R. Radha, M. Krishna, S. Thangavelu (Eds.), *Proc. Internat. Conf. on Wavelets and Applications*, Chennai, January 2002, Allied Publishers, New Delhi, 2003, pp. 1–56.
- [11] H.G. Feichtinger, Modulation spaces: looking back and ahead, *Sampl. Theory Signal Image Process.*, *Int. J.* 5 (2) (2006) 109–140.
- [12] H.G. Feichtinger, G. Zimmermann, A Banach space of test functions for Gabor analysis, in: H.G. Feichtinger, T. Strohmer (Eds.), *Gabor Analysis and Algorithms: Theory and Applications*, Applied and Numerical Harmonic Analysis, Birkhäuser, Boston, 1998, pp. 123–170.
- [13] K. Gröchenig, Aspects of Gabor analysis on locally compact Abelian groups, in: H.G. Feichtinger, T. Strohmer (Eds.), *Gabor Analysis and Algorithms: Theory and Applications*, Applied and Numerical Harmonic Analysis, Birkhäuser, Boston, MA, USA, 1998, pp. 123–170.
- [14] K. Gröchenig, *Foundations of Time-Frequency Analysis*, Appl. Numer. Harmon. Anal., Birkhäuser, Boston, MA, USA, 2001.
- [15] K. Gröchenig, A pedestrian’s approach to pseudodifferential operators, in: C. Heil (Ed.), *Harmonic Analysis and Applications*, Volume in Honor of John J. Benedetto’s 65th Birthday, Birkhäuser Boston, Boston, MA, 2006, pp. 139–169.
- [16] K. Gröchenig, M. Leinert, Wiener’s lemma for twisted convolution and Gabor frames, *J. Amer. Math. Soc.* 17 (2004) 1–18.
- [17] K. Gröchenig, J.L. Romero, J. Stöckler, Sampling theorems for shift-invariant spaces, Gabor frames, and totally positive functions, *Invent. Math.* 211 (3) (2018) 1119–1148.
- [18] K. Gröchenig, J. Stöckler, Gabor frames and totally positive functions, *Duke Math. J.* 162 (6) (2013) 1003–1031.
- [19] K. Gröchenig, T. Strohmer, Pseudodifferential operators on locally compact abelian groups and Sjöstrand’s symbol class, *J. Reine Angew. Math.* 613 (2007) 121–146.
- [20] V.P. Havin, N.K. Nikol’skij, *Commutative Harmonic Analysis II. Group Methods in Commutative Harmonic Analysis*, Springer, Berlin, 1998, Translated from Russian.
- [21] E. Hewitt, K.A. Ross, *Abstract Harmonic Analysis I*, Grundlehren Math. Wiss., vol. 115, Springer, Berlin, 1963.
- [22] M.S. Jakobsen, On a (no longer) new Segal algebra: a review of the Feichtinger algebra, *J. Fourier Anal. Appl.* 24 (6) (2018) 1579–1660.
- [23] M.S. Jakobsen, J. Lemvig, Density and duality theorems for regular Gabor frames, *J. Funct. Anal.* 270 (1) (2016) 229–263.
- [24] M.S. Jakobsen, F. Luef, *Duality of Gabor frames and Heisenberg modules*, ArXiv e-prints, 2018.
- [25] A.J.E.M. Janssen, Zak transforms with few zeros and the tie, in: H.G. Feichtinger, T. Strohmer (Eds.), *Advances in Gabor Analysis*, Applied and Numerical Harmonic Analysis, Birkhäuser, Basel, 2003, pp. 31–70.
- [26] S. Kaliszewski, T. Omland, J. Quigg, Cuntz–Li algebras from a -adic numbers, *Rev. Roum. Math. Pures Appl.* 59 (3) (2014) 331–370.
- [27] E. King, M. Skopina, Quincunx multiresolution analysis for $L^2(\mathbb{Q}_2^2)$, p -Adic Numbers Ultrametric Anal. Appl. 2 (3) (2010) 222–231.
- [28] G. Kutyniok, E. Kaniuth, Zeros of the Zak transform on locally compact abelian groups, *Proc. Amer. Math. Soc.* 126 (12) (1998) 3561–3569.
- [29] N. Larsen, X. Li, The 2-adic ring C^* -algebra of the integers and its representations, *J. Funct. Anal.* 262 (4) (2012) 1392–1426.
- [30] F. Latrémolière, J. Packer, Noncommutative solenoids and their projective modules, *Commu. Noncommut. Harmon. Anal. Appl.* 603 (2013) 35–53.
- [31] F. Latrémolière, J. Packer, Explicit construction of equivalence bimodules between noncommutative solenoids, *Contemp. Math.*, Amer. Math. Soc. 650 (2015) 111–140.
- [32] F. Latrémolière, J. Packer, Noncommutative solenoids and the Gromov–Hausdorff propinquity, *Proc. Amer. Math. Soc.* 145 (2017) 2043–2057.
- [33] D. Li, H.K. Jjiang, Basic results of Gabor frame on local fields, *Chin. Ann. Math.* 28 (2) (2007) 165–176.
- [34] V. Losert, A characterization of the minimal strongly character invariant Segal algebra, *Ann. Inst. Fourier (Grenoble)* 30 (1980) 129–139.
- [35] F. Luef, Projections in noncommutative tori and Gabor frames, *Proc. Amer. Math. Soc.* 139 (2) (2011) 571–582.
- [36] Y.I. Lyubarskii, Frames in the Bargmann space of entire functions, in: *Entire and Subharmonic Functions*, in: *Advances in Soviet Mathematics*, vol. 11, American Mathematical Society (AMS), Providence, RI, 1992, pp. 167–180.
- [37] Y. Manin, A. Panchishkin, *Introduction to Modern Number Theory*, second edition, Encyclopaedia Math. Sci., vol. 49, Springer-Verlag, Berlin, 2005.
- [38] D. Ramakrishnan, R.J. Valenza, *Fourier Analysis on Number Fields*, Springer, New York, 1999.
- [39] H. Reiter, *Metaplectic Groups and Segal Algebras*, Lecture Notes in Mathematics, Springer, Berlin, 1989.
- [40] M.A. Rieffel, Projective modules over higher-dimensional noncommutative tori, *Can. J. Math.* 40 (2) (1988) 257–338.
- [41] K. Seip, Density theorems for sampling and interpolation in the Bargmann–Fock space. I, *J. Reine Angew. Math.* 429 (1992) 91–106.
- [42] F.A. Shah, Gabor frames on local fields of positive characteristic, *Tbil. Math. J.* 9 (2) (2016) 129–139.
- [43] F.A. Shah, Gabor-type expansions on local fields, *SeMa J.* 75 (3) (2018) 485–498.
- [44] V. Shelkovich, M. Skopina, p -adic Haar multiresolution analysis and pseudo-differential operators, *J. Fourier Anal. Appl.* 15 (3) (2009) 366–393.