



INSTITUT DE FRANCE
Académie des sciences

Comptes Rendus

Mathématique


Arnaud Beauville

A non-hyperelliptic curve with torsion Ceresa class

Volume 359, issue 7 (2021), p. 871-872

<<https://doi.org/10.5802/crmath.226>>

© Académie des sciences, Paris and the authors, 2021.
Some rights reserved.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



Les Comptes Rendus. Mathématique sont membres du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org



Algebraic geometry / *Geometrie algébrique*

A non-hyperelliptic curve with torsion Ceresa class

Arnaud Beauville*, ^a

^a Université Côte d'Azur, CNRS – Laboratoire J.-A. Dieudonné, Parc Valrose, F-06108

Nice cedex 2, France.

E-mail: arnaud.beauville@unice.fr

Abstract. We exhibit a non-hyperelliptic curve C of genus 3 such that the class of the Ceresa cycle $[C] - [-C]$ in the intermediate Jacobian of J_C is torsion.

Manuscript received 21st May 2021, accepted 24th May 2021.

1. Introduction

Let C be a complex curve of genus $g \geq 3$, and p a point of C . We embed C into its Jacobian J by the Abel–Jacobi map $x \mapsto [x] - [p]$. The *Ceresa cycle* $\mathfrak{z}_p(C)$ is the cycle $[C] - [(-1_J)^*C]$ in the Chow group $CH_1(J)_{\text{hom}}$ of homologically trivial 1-cycles. The *Ceresa class* $c_p(C)$ is the image of $\mathfrak{z}_p(C)$ in the intermediate Jacobian $\mathfrak{J}_1(J)$ parameterizing 1-cycles under the Abel–Jacobi map $CH_1(J)_{\text{hom}} \rightarrow \mathfrak{J}_1(J)$.

When C is general, $\mathfrak{z}_p(C)$ is not algebraically trivial [2]. On the other hand, if C is hyperelliptic $\mathfrak{z}_p(C)$ is algebraically trivial – in fact it is zero if one chooses for p a Weierstrass point. Not much is known besides these two extreme cases. There are few curves for which $\mathfrak{z}_p(C)$ is known to be not algebraically trivial: Fermat curves of degree ≤ 1000 [4], and the Klein quartic [5]. An essential ingredient of these results is the fact that $c_p(C)$ is not a torsion class.

It is an open question whether there are non-hyperelliptic curves with $\mathfrak{z}_p(C)$ algebraically trivial. As observed in [3, Remark 2.4], this condition is equivalent to a number of interesting properties: in particular the existence of a *multiplicative Chow–Künneth decomposition* modulo algebraic equivalence, or the fact that the class $[C] \in CH_1(J) \otimes \mathbb{Q}$ is algebraically equivalent to the minimal class $\frac{\theta^{g-1}}{(g-1)!}$, where $\theta \in CH^1(J)$ is the class of the principal polarization.

In this note we exhibit a curve C of genus 3 with the weaker property that the Ceresa class $c_p(C)$ is torsion (under the Bloch–Beilinson conjectures, this actually implies the algebraic triviality of $\mathfrak{z}_p(C)$ up to torsion). The construction is very simple: the curve C has an automorphism σ which

* Corresponding author.

fixes a point p , and therefore preserves $c_p(C)$; we just have to check that the fixed point set of σ acting on $\mathfrak{J}_1(J)$ is finite.

A similar example, based on a much more sophisticated approach, appears in [1, Remark 3.6].

2. The result

Proposition 1. *Let $C \subset \mathbb{P}^2$ be the genus 3 curve defined by $X^4 + XZ^3 + Y^3Z = 0$, and let $p = (0, 0, 1)$. The Ceresa class $c_p(C)$ is torsion.*

Proof. Let ω be a primitive 9th root of unity. We consider the automorphism σ of C defined by $\sigma(X, Y, Z) = (X, \omega^2 Y, \omega^3 Z)$. We have $\sigma(p) = p$; therefore σ preserves the Ceresa cycle $\mathfrak{z}_p(C)$, and also its class $c_p(C)$ in $\mathfrak{J} := \mathfrak{J}_1(J)$.

Thus it suffices to prove that σ has finitely many fixed points on \mathfrak{J} ; equivalently, that the eigenvalues of σ acting on the tangent space $T_0(\mathfrak{J})$ are $\neq 1$.

Now $T_0(\mathfrak{J})$ is identified with $H^{0,3}(J) \oplus H^{1,2}(J) = \wedge^3 V^* \oplus (\wedge^2 V^* \otimes V)$, where $V = H^{1,0}(J) = H^0(C, K_C)$. We first compute the eigenvalues of σ on V . The elements of V are of the form $L \cdot \frac{XdZ - ZdX}{Y^2Z}$, with $L \in H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}}(1))$; it follows that the eigenvalues of σ on V are $\omega^5, \omega^7, \omega^8$. Therefore the eigenvalue on $\wedge^3 V^*$ is ω^7 , and the eigenvalues on $\wedge^2 V^*$ are $\omega^3, \omega^5, \omega^6$. Thus each product of an eigenvalue on $\wedge^2 V^*$ and one on V is $\neq 1$, hence the Proposition. \square

References

- [1] D. Bisogno, W. Li, D. Litt, P. Srinivasan, “Group-theoretic Johnson classes and non-hyperelliptic curves with torsion Ceresa class”, <https://arxiv.org/abs/2004.06146>, 2020.
- [2] G. Ceresa, “ C is not algebraically equivalent to C^- in its Jacobian”, *Ann. Math.* **117** (1983), no. 2, p. 285-291.
- [3] L. Fu, R. Laterveer, C. Vial, “Multiplicative Chow–Künneth decompositions and varieties of cohomological K3 type”, *Ann. Mat. Pura Appl. (4)* **200** (2021), no. 5, p. 2085-2126.
- [4] N. Otsubo, “On the Abel–Jacobi maps of Fermat Jacobians”, *Math. Z.* **270** (2012), no. 1-2, p. 423-444.
- [5] Y. Tadokoro, “A nontrivial algebraic cycle in the Jacobian variety of the Klein quartic”, *Math. Z.* **260** (2008), no. 2, p. 265-275.