



INSTITUT DE FRANCE  
Académie des sciences

# *Comptes Rendus*

---

# *Mathématique*


Sergio A. Carrillo

**A quick proof of the regularity of the flow of analytic vector fields**

Volume 359, issue 9 (2021), p. 1155-1159

<<https://doi.org/10.5802/crmath.271>>

© Académie des sciences, Paris and the authors, 2021.  
*Some rights reserved.*

 This article is licensed under the  
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.  
<http://creativecommons.org/licenses/by/4.0/>



*Les Comptes Rendus. Mathématique* sont membres du  
Centre Mersenne pour l'édition scientifique ouverte  
[www.centre-mersenne.org](http://www.centre-mersenne.org)



Complex analysis and geometry, Dynamical systems / *Analyse et géométrie complexes, Systèmes dynamiques*

# A quick proof of the regularity of the flow of analytic vector fields

Sergio A. Carrillo<sup>a</sup>

<sup>a</sup> Programa de matemáticas, Universidad Sergio Arboleda, Calle 74 # 14-14, Bogotá, Colombia.

E-mail: [sergio.carrillo@usa.edu.co](mailto:sergio.carrillo@usa.edu.co)

**Abstract.** We offer a new and elementary proof of the convergence of the Lie series giving the flow of an analytic vector field as well as a natural deduction of such series.

**2020 Mathematics Subject Classification.** 32M25.

**Funding.** The author is supported by Ministerio de Economía y Competitividad from Spain, under the Project “Métodos asintóticos, algebraicos y geométricos en foliaciones singulares y sistemas dinámicos” (Ref.: PID2019-105621GB-I00) and Univ. Sergio Arboleda project IN.BG.086.20.002.

*Manuscript received 29th July 2021, accepted 8th September 2021.*

## 1. Introduction

Consider a germ  $X$  of an analytic vector field defined at a point  $p$  on a  $n$ -complex analytic manifold  $M$ . In local coordinates  $z = (z_1, \dots, z_n)$  such that  $z(p) = 0 \in \mathbb{C}^n$ ,  $X$  is a vector field in  $(\mathbb{C}^n, 0)$ , i.e., a  $n$ -tuple of convergent power series

$$X = (X_1, \dots, X_n) \in \mathbb{C}\{z\}^n,$$

or the derivation

$$X(f)(z) = \sum_{j=1}^n X_j(z) \frac{\partial f}{\partial z_j}(z), \quad \text{where } f \in \mathbb{C}\{z\}.$$

The dynamics of  $X$  is determined through its flow  $\phi_X(t, z)$ , which is the solution of the differential equation

$$\partial_t \phi_X(t, z) = X(\phi_X(t, z)), \quad \phi_X(0, z) = z. \quad (1)$$

Geometrically, for  $z_0 \in \mathbb{C}^n$  near 0, the map  $t \in (\mathbb{C}, 0) \mapsto \phi_X(t, z_0)$  is a curve passing through  $z_0$  at  $t = 0$ , having at the point  $\phi_X(t, z_0)$  the tangent vector  $X(\phi_X(t, z_0))$ .

A classical result in the general theory of ordinary differential equations is that (1) has a unique solution which is as regular as  $X$  is. In our case, if  $X$  is analytic, the flow will be analytic in both variables  $(t, z)$ . More specifically, we have:

**Theorem 1.** *Given a germ  $X$  of an analytic vector field at  $0 \in \mathbb{C}^n$ , the differential equation (1) admits a unique analytic solution  $\phi_X(t, z) \in \mathbb{C}\{t, z\}^n$  which is given by the Lie series*

$$\phi_X(t, z) = \sum_{m=0}^{\infty} (X^m(z_1), \dots, X^m(z_n)) \frac{t^m}{m!}. \tag{2}$$

Here  $X^m$  denotes the iterations of  $X$  as a derivation, i.e.,

$$X^0(f) = f, \quad X^{m+1}(f) := X(X^m(f)), \quad m \geq 0, \quad \text{for } f \in \mathbb{C}\{z\}. \tag{3}$$

The proof of the analyticity of  $\phi_X$  is usually done using the Cauchy majorant series technique [3, Chapter VII] or by the contraction map principle [5, Chapter I.1] which extends to less regular systems. Then, the expansion (2) is obtained as a consequence of analyticity writing  $\phi_X$  as its Taylor series at  $t = 0$ . In particular, once we have an analytic solution of (1) at  $(t, z) = (0, 0)$ , it must be series (2). On the other hand, the series (2) can be shown to be convergent employing majorizing series as explained in Gröbner’s book [4, Satz 2]. Due to his contribution this series is also known as a Gröbner–Lie series, having applications in the inversion of a mapping of power series [9, Section 2], in the resolution of systems of autonomous and non-autonomous differential equations [4, 8], and in the classification of tangent to the identity biholomorphisms [1, 6].

We will prove Theorem 1 finding the series (2) as a solution to an auxiliary *linear* partial differential equation, proving its convergence with the aid of Nagumo norms, and finally checking that (2) solves (1) using the analyticity of this map. This proof is even more elementary than previous approaches due to the linear character of the auxiliary problem. In fact, it only uses Cauchy’s formulas and Taylor expansions.

## 2. The proof of the theorem

We need some preliminaries. First, we extend (3) to maps  $F = (f_1, \dots, f_n) \in \mathbb{C}\{z\}^n$  by

$$X^m(F) := (X^m(f_1), \dots, X^m(f_n)), \quad m \geq 0.$$

In particular,  $X^m(\text{id}) = (X^m(z_1), \dots, X^m(z_n))$  where  $\text{id}(z) = z$ . Moreover,

$$X^{m+1}(F)(z) = D_z[X^m(F)](z) \cdot X(z), \tag{4}$$

where  $D_z$  denotes the Jacobian matrix with respect to  $z$  and  $\cdot$  stands for the usual product of a matrix and a vector.

Second, if  $\phi_X(s, \phi_X(\tau, z_0))$  and  $\phi_X(\tau + s, z_0)$  are defined, the 1-parametric group law asserts that

$$\phi_X(s, \phi_X(\tau, z_0)) = \phi_X(\tau + s, z_0),$$

because both sides of the equality are solutions of (1) at the point  $\phi_X(\tau, z_0)$ , but such solution is unique. In particular,  $\phi_X(t, \phi_X(-t, z_0)) = z_0$ , for small  $z_0$  and  $t$ . If we differentiate this equation with respect to  $t$ , we find that

$$\partial_t \phi_X(t, \phi_X(-t, z_0)) + D_z \phi_X(t, \phi_X(-t, z_0)) \partial_t \phi_X(-t, z_0)(-1) = 0.$$

Using (1) at  $-t$  and setting  $z = \phi_X(-t, z_0)$  it turns out that  $w = \phi_X$  satisfies

$$\partial_t w(t, z) = D_z w(t, z) \cdot X(z), \tag{5}$$

see [1, p. 546]. This is a linear PDE in  $w \in \mathbb{C}^n$  that will lead to the series (2), as we shall see.

Finally, to establish the convergence of (2) we use the Nagumo norms. These were introduced by M. Nagumo in his work [7] on power series solutions of analytic PDEs. These norms have numerous applications, for instance, in singularly perturbed problems, see [2].

Let  $D_r^n = \{z \in \mathbb{C}^n : |z_j| < r, j = 1, \dots, n\}$  be the polydisc centered at the origin with polyradius  $(r, \dots, r)$ , for a common  $r > 0$ . We also write  $\mathcal{O}(D_r^n)$  for the set of complex-valued holomorphic functions on  $D_r^n$ .

**Definition 2.** If  $r > 0$ ,  $f \in \mathcal{O}(D_r^n)$ , and  $m \in \mathbb{N}$  the  $m$ th Nagumo norm of  $f$  is

$$\|f\|_m := \sup_{z \in D_r^n} |f(z)|(r - |z_1|)^m \cdots (r - |z_n|)^m.$$

If  $m = 0$  this is just the supremum norm which is finite in case  $f$  is bounded on  $D_r^n$ . In this situation  $\|f\|_m \leq r^{mn} \|f\|_0$ . Moreover, we have:

**Lemma 3.** If  $f, g \in \mathcal{O}(D_r^n)$  and  $m, k \in \mathbb{N}$ , then

- (1)  $\|f + g\|_m \leq \|f\|_m + \|g\|_m$  and  $\|fg\|_{m+k} \leq \|f\|_m \|g\|_k$ .
- (2)  $\|\frac{\partial f}{\partial z_j}\|_{m+1} \leq e(m+1)r^{n-1} \|f\|_m$ ,  $j = 1, \dots, n$ .

For the sake of completeness we include the proof at the end of the paper. Since we work with vector- and matrix-valued maps, we extend these norms for  $F = (f_1, \dots, f_n) \in \mathcal{O}(D_r^n)^n$  and  $A = (A_{ij})_{i,j=1,\dots,n} \in \mathcal{O}(D_r^n)^{n \times n}$  by the rules

$$\|F\|_m := \max_{1 \leq j \leq n} \|f_j\|_m \quad \text{and} \quad \|A\|_m := \max_{1 \leq i \leq n} \sum_{j=1}^n \|A_{ij}\|_m.$$

Then Lemma 3 shows that

$$\|A \cdot F\|_{m+k} \leq \|A\|_m \|F\|_k, \quad \|D_z F\|_{m+1} \leq ner^{n-1} (m+1) \|F\|_m. \tag{6}$$

We are now ready to prove the result.

**Proof of Theorem 1.** We assume  $X$  is not identically zero. Let us fix a closed polydisc  $\overline{D_r^n}$  where each  $X_j$  is bounded and holomorphic in a neighborhood of this domain. Taking into account equation (5), we search for a solution  $w(t, z)$  of the initial-value problem

$$\partial_t w(t, z) = D_z w(t, z) \cdot X(z), \quad w(0, z) = z. \tag{7}$$

Setting  $w(t, z) = \sum_{m=0}^\infty \varphi_m(z) t^m / m!$ , after equating coefficients in common powers of  $t^m$ , we find

$$\varphi_0(z) = w(0, z) = z, \quad \varphi_{m+1}(z) = D_z \varphi_m(z) \cdot X(z), \quad m \geq 0. \tag{8}$$

Due to (4),  $\varphi_m(z) = X^m(\text{id})(z)$ , and thus  $\varphi_m$  is holomorphic and bounded on  $D_r^n$ , for all  $m \geq 0$ . To prove that  $w(t, z)$  converges we study the series

$$W(\tau) := \sum_{m=0}^\infty \|\varphi_m\|_m \frac{\tau^m}{m!},$$

where  $\|\cdot\|_m$  denotes the vector  $m$ th Nagumo norm on the polydisc  $D_r^n$ . First, since  $X$  is bounded, then  $\|X\|_0 > 0$  is finite. Now, thanks to (6) we find that

$$\frac{\|\varphi_{m+1}\|_{m+1}}{(m+1)!} \leq \frac{\|D_z \varphi_m\|_{m+1}}{(m+1)!} \|X\|_0 \leq ner^{n-1} \|X\|_0 \frac{\|\varphi_m\|_m}{m!}.$$

Since  $\|\varphi_0\|_0 = \|z\|_0 = r$ , the previous inequality shows that

$$\frac{\|\varphi_m\|_m}{m!} \leq r\alpha^m, \quad \alpha := ner^{n-1} \|X\|_0.$$

This proves that  $W(\tau)$  is convergent. Finally, if  $|z_1|, \dots, |z_n| \leq \rho < r$ , then

$$\max_{1 \leq j \leq n} \frac{|X^m(z_j)|}{m!} \leq \frac{\|\varphi_m\|_m / m!}{(r - |z_1|)^m \cdots (r - |z_n|)^m} \leq r\alpha^m (r - \rho)^{-mn},$$

showing that  $w(t, z)$  converges in the polydisc  $D_{(r-\rho)^n/\alpha} \times D_\rho^n$ .

To conclude we prove that  $w(t, z)$  solves (1). For this, since  $X \circ w$  is analytic, it is equal to its Taylor series around  $t = 0$ :

$$X(w(t, z)) = \sum_{m=0}^\infty \partial_t^m (X \circ w)(0, z) \frac{t^m}{m!}.$$

Therefore, we need to check that

$$\partial_t^m (X \circ w)(0, z) = \varphi_{m+1}(z), \quad \text{for all } m \geq 0, \tag{9}$$

since then

$$X(w(t, z)) = \sum_{m=0}^{\infty} \varphi_{m+1}(z) \frac{t^m}{m!} = \partial_t \left( \sum_{m=0}^{\infty} \varphi_m(z) \frac{t^m}{m!} \right) = \partial_t w(t, z),$$

as required. We proceed by induction on  $m$ . If  $m = 0$ ,  $X(w(0, z)) = X(z) = \varphi_1(z)$  and (9) is valid. If we assume it holds for some  $m$ , note first that

$$\begin{aligned} \partial_t^{m+1}(X \circ w) &= \partial_t^m \partial_t(X \circ w) = \partial_t^m ((D_z X \circ w) \cdot \partial_t w) = \partial_t^m ((D_z X \circ w) \cdot D_z w \cdot X) \\ &= \partial_t^m D_z(X \circ w) \cdot X = D_z \partial_t^m(X \circ w) \cdot X. \end{aligned}$$

Setting  $t = 0$  and using the induction hypothesis we conclude that

$$\begin{aligned} \partial_t^{m+1}(X \circ w)(0, z) &= D_z \partial_t^m(X \circ w)(0, z) \cdot X(z) \\ &= D_z (\partial_t^m(X \circ w)(0, z)) \cdot X(z) = D_z \varphi_{m+1}(z) \cdot X(z) = \varphi_{m+2}(z), \end{aligned}$$

as desired. □

Theorem 1 gives an alternative to establish some elementary observations:

(i). If we change the initial condition in (7) and consider instead

$$\partial_t w(t, z) = D_z w(t, z) \cdot X(z), \quad w(0, z) = F(z),$$

for  $F \in \mathbb{C}\{z\}^n$ , this system has as unique solution  $\Phi(t, z) = \sum_{m=0}^{\infty} X^m(F)(z) \frac{t^m}{m!}$ , whose form and convergence are determined in exactly the same way as before. But  $\Psi(t, z) = F(\phi_X(t, z))$  also satisfies this system since  $\Psi(0, z) = F(z)$  and

$$\begin{aligned} \partial_t \Psi(t, z) &= D_z[F](\phi_X(t, z)) \partial_t \phi_X(t, z) \\ &= D_z[F](\phi_X(t, z)) D_z[\phi_X](t, z) X(z) = D_z \Psi(t, z) X(z), \end{aligned}$$

where we used (5). In conclusion, we recover the classical formula

$$F(\phi_X(t, z)) = \sum_{m=0}^{\infty} X^m(F)(z) \frac{t^m}{m!},$$

for the Lie series associated to  $X$  and  $F$ .

(ii). If  $X(0) = 0$  and  $X(z) = Az + \dots$ , where  $A = D_z X(0)$ , then

$$D_z[X^m(\text{id})](0) = A^m.$$

In fact, differentiating the second equation in (8) and taking  $z = 0$  shows that  $D_z \varphi_{m+1}(0) = D_z \varphi_m(0) \cdot A$ , which proves the claim. Writing  $\phi_X(t, z)$  as a power series in  $z$  with coefficients series in  $t$  we get

$$\phi_X(t, z) = e^{At} z + \dots$$

In particular,  $\phi_X(t, \cdot) : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n, 0)$  is a biholomorphism. Moreover, if  $A = 0$ , then  $\phi_X(t, \cdot)$  is tangent to the identity.

We conclude this note with the proof of Lemma 3.

**Proof of Lemma 3.** We prove (2) for  $m > 0$  and  $j = 1$ . The remaining statements are easily supplied. Fix  $z = (z_1, z') \in D_r^n$ ,  $z' \in \mathbb{C}^{n-1}$ , and take  $0 < \rho < r - |z_1|$  so that  $D = \{y \in \mathbb{C} : |y - z_1| \leq \rho\}$  is contained in  $D_r$ . Cauchy's formulas show that

$$\left| \frac{\partial f}{\partial z_1}(z) \right| \leq \frac{1}{\rho} \sup_{|y-z_1|=\rho} |f(y, z')|, \tag{10}$$

By Definition 2 we have that

$$|f(y, z')| \leq \frac{\|f\|_m}{(r - |y|)^m R}, \quad \text{where } R = \prod_{j=2}^n (r - |z_j|)^m,$$

and taking the supremum for  $y$  on the boundary of the disc  $D$ , we obtain

$$\sup_{|y-z_1|=\rho} |f(y, z')| \leq \frac{\|f\|_m}{(r - (\rho + |z_1|))^m R}. \quad (11)$$

Putting together (10) and (11) we find

$$\left| \frac{\partial f}{\partial z_1}(z) \prod_{j=1}^n (r - |z_j|)^{m+1} \right| \leq r^{n-1} \frac{\|f\|_m}{\rho} \frac{(r - |z_1|)^{m+1}}{(r - |z_1| - \rho)^m}.$$

The result follows taking  $\rho = \frac{r - |z_1|}{m+1}$  and using the inequality  $(1 - \frac{1}{m+1})^{-m} < e$ .  $\square$

## References

- [1] P. Ahern, J.-P. Rosay, "Entire functions, in the classification of differentiable germs tangent to the identity, in one or two variables", *Trans. Am. Math. Soc.* **347** (1995), no. 2, p. 543-572.
- [2] M. Canalis-Durand, J.-P. Ramis, R. Schäfke, Y. Sibuya, "Gevrey solutions of singularly perturbed differential equations", *J. Reine Angew. Math.* **518** (2000), no. 2, p. 95-129.
- [3] H. Cartan, *Elementary theory of analytic functions of one or several complex variables*, Hermann, 1973.
- [4] W. Gröbner, *Die Lie-Reihen und ihre Anwendungen*, VEB Deutscher Verlag der Wissenschaften, 1960.
- [5] Y. Ilyashenko, S. Yakovenko, *Lectures on analytic differential equations*, Graduate Studies in Mathematics, vol. 86, American Mathematical Society, 2008.
- [6] J. Martinet, J.-P. Ramis, "Classification analytique des équations différentielles non linéaires résonnantes du premier ordre", *Ann. Sci. Éc. Norm. Supér.* **16** (1983), no. 4, p. 571-621.
- [7] M. Nagumo, "Über das Anfangswertproblem partieller Differentialgleichungen", *Jap. J. Math.* **18** (1942), p. 41-47.
- [8] S. Steinberg, "Lie series and nonlinear ordinary differential equations", *J. Math. Anal. Appl.* **101** (1984), p. 39-63.
- [9] R. Winkel, "An exponential formula for polynomial vector fields: II. Lie series, exponential substitution, and rooted trees", *Adv. Math.* **147** (1999), p. 260-303.