



ACADÉMIE
DES SCIENCES
INSTITUT DE FRANCE

Comptes Rendus

Mathématique


Xuan Zhao and Zhenliang Zhang

Some lacunarity properties of partial quotients of real numbers

Volume 362 (2024), p. 1089-1095

Online since: 5 November 2024

<https://doi.org/10.5802/crmath.594>

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*The Comptes Rendus. Mathématique are a member of the
Mersenne Center for open scientific publishing*
www.centre-mersenne.org — e-ISSN : 1778-3569



Research article / *Article de recherche*
Number theory / *Théorie des nombres*

Some lacunarity properties of partial quotients of real numbers

Quelques propriétés de lacunarité des quotients partiels de nombres réels

Xuan Zhao ^a and Zhenliang Zhang ^{*,b}

^a National Education Examinations Authority, 100084 Beijing, China

^b School of Mathematical Sciences, Chongqing Normal University, Chongqing, 401331, P. R. China

E-mails: zhaox@mail.neea.edu.cn, zhliang_zhang@163.com

Abstract. We consider lacunarity properties of sequence of partial quotients for real numbers in their continued fraction expansions. Hausdorff dimension of the sets of points with different lacunarity conditions on their partial quotients are calculated.

Résumé. Nous considérons des propriétés de lacunarité de la suite des quotients partiels du développement en fraction continue de nombres réels. Nous calculons la dimension de Hausdorff d'ensembles de points dont la suite des quotients partiels satisfait à différentes conditions de lacunarité.

Keywords. Hausdorff dimension, Continued fraction expansion.

Mots-clés. Dimension de Hausdorff, développement en fraction continue.

Mathematical subject classification (2010). 11K50, 28A78.

Funding. Zhenliang Zhang is supported by project of the Science and Technology Research Program of Chongqing Municipal Education Commission (No. KJQN202100528), Natural Science Foundation of Chongqing (No. CSTB2022NSCQ-MSX1255).

Manuscript received 21 July 2023, revised 26 September 2023 and 13 October 2023, accepted 27 November 2023.

1. Introduction

Continued fraction expansion is induced by the Gauss transformation $T : [0, 1) \rightarrow [0, 1)$ given by

$$T(0) := 0, \quad T(x) = \frac{1}{x} \bmod 1, \quad x \in (0, 1).$$

*Corresponding author

Then every irrational number $x \in [0, 1)$ can be uniquely expanded into an infinite continued fraction:

$$x = \frac{1}{a_1(x) + \frac{1}{a_2(x) + \frac{1}{\ddots}}}} := [a_1(x), a_2(x), \dots],$$

where $a_1(x) = [1/x]$ and $a_n(x) = a_1(T^{n-1}(x))$ for $n \geq 2$ are called the partial quotients of x . The finite truncation

$$\frac{p_n(x)}{q_n(x)} = [a_1(x), \dots, a_n(x)]$$

is called the n th convergent of x .

It is well-known that continued fraction expansion plays an important role in Diophantine approximation and dynamical systems:

- In Diophantine approximation, how well an irrational number can be approximated by rationals depends on the growth rate of the partial quotients. For example, the classic Jarník set [12] can be expressed as

$$\mathcal{X}(\psi) = \left\{ x \in [0, 1) : a_{n+1}(x) \geq \psi(q_n(x)), \text{ for infinitely many } n \in \mathbb{N} \right\}.$$

- In dynamical system, continued fraction is a classic dynamics with infinitely many branches [17].

As said above, the growth rate of the partial quotient is tightly related to the Diophantine properties of an irrational number. Many metric results have been achieved in this aspect, such as the Borel–Bernstein theorem that deals with Lebesgue measure theory on the growth rate of $\{a_n(x)\}_{n \geq 1}$. Hausdorff dimension of sets obeying some restrictions on the partial quotients have been well established in Good [6], Łuczak [16], Wang & Wu [18], etc.

Very recently, it was found by Kleinbock and Wadleigh [14] that the Dirichlet improvable set is highly related to the growth rate of the product of two consecutive partial quotients in the following sense. Let $\psi: [t_0, \infty) \rightarrow \mathbb{R}_+$ be non-increasing with $t_0 > 1$ is fixed and $t\psi(t) < 1$ for all $t \geq t_0$. The ψ -Dirichlet improvable set is defined as

$$\mathcal{D}(\psi) = \left\{ x \in [0, 1) : \min_{1 \leq q < Q} \|qx\| < \psi(Q), \text{ for all } Q \gg 1 \right\}.$$

Then by taking $\Psi(q) = \frac{q\psi(q)}{1-q\psi(q)}$, one has

$$G(\Psi) \subset [0, 1) \setminus \mathcal{D}(\psi) \subset G\left(\frac{\Psi}{4}\right)$$

where

$$G(\Psi) = \left\{ x \in [0, 1) : a_{n+1}(x)a_n(x) \geq \Psi(q_n(x)), \text{ for infinitely many } n \in \mathbb{N} \right\}.$$

Since then, the metric theory relating the growth of two consecutive partial quotients are extensively studied in [1–3, 7–10, 15].

In this note, we take another turn by studying the relative growth rate of two consecutive partial quotients. All the above mentioned works are concerning the size of limsup sets, however, the sets we will consider below are of liminf nature. So the method used here is different from the above works (see [10] for a unified way of dealing with the Hausdorff dimension of the above mentioned works). These liminf sets concerns the points with partial quotients increasing very fast, so can be uniformly well approximated by their convergents. At first, we give some notations. Let $\{s_n\}_{n \geq 1}$ be a sequence of strictly increasing sequence of integers.

- call it a sub-lacunary sequence, if

$$\lim_{n \rightarrow \infty} s_{n+1}/s_n = 1;$$

- call it lacunary if $\exists c > 1$ such that

$$s_{n+1}/s_n \geq c, \text{ for all } n \gg 1;$$

- log-lacunary if for some $t > 0$,

$$\frac{s_{n+1}}{s_n} \geq (\log n)^t, \text{ for all } n \gg 1;$$

- polynomial-lacunary if for some $t > 0$,

$$\frac{s_{n+1}}{s_n} \geq n^t, \text{ for all } n \gg 1;$$

- exponential-lacunary if for some $b > 1$,

$$\frac{s_{n+1}}{s_n} \geq b^n, \text{ for all } n \gg 1.$$

At first, we give an auxiliary set to be compliant with the set of points with strictly increasing partial quotients.

$$G = \left\{ x \in [0, 1) : \{a_n(x)\} \text{ is a strictly increasing sequence} \right\}. \tag{1}$$

Then we define

$$\begin{aligned} \mathcal{S}_G &= \left\{ x \in G : \{a_n(x)\} \text{ is sub-lacunary} \right\}, \\ \mathcal{L}_G &= \left\{ x \in G : \{a_n(x)\} \text{ is log-lacunary} \right\}, \\ \mathcal{P}_G &= \left\{ x \in G : \{a_n(x)\} \text{ is polynomial-lacunary} \right\}, \\ \mathcal{E}_G &= \left\{ x \in G : \{a_n(x)\} \text{ is exponential-lacunary} \right\}. \end{aligned}$$

In this note, we show that

Theorem 1. *By denoting \dim_H the Hausdorff dimension, we have*

$$\dim_H \mathcal{S}_G = \dim_H \mathcal{L}_G = \dim_H \mathcal{P}_G = \dim_H \mathcal{E}_G = 1/2.$$

2. Preliminaries

Recall that $p_n(x)/q_n(x)$ is the n th convergent of x . The numerator and denominator of $p_n(x)/q_n(x)$ can be determined recursively: for any $k \geq 1$

$$p_k(x) = a_k(x)p_{k-1}(x) + p_{k-2}(x), \quad q_k(x) = a_k(x)q_{k-1}(x) + q_{k-2}(x) \tag{2}$$

with the conventions $p_0 = 0, q_0 = 1, p_{-1} = 1, q_{-1} = 0$.

For simplicity, we write

$$p_n(x) = p_n(a_1, \dots, a_n) = p_n, \quad q_n(x) = q_n(a_1, \dots, a_n) = q_n \tag{3}$$

when the partial quotients a_1, \dots, a_n are clear.

For any positive integers a_1, \dots, a_n , define

$$I_n(a_1, \dots, a_n) := \{x \in [0, 1) : a_1(x) = a_1, \dots, a_n(x) = a_n\}$$

and call it a *cylinder of order n* . We use $I_n(x)$ to denote the n th order cylinder containing x .

Proposition 2 (Khinchin [13]). For any $n \geq 1$ and $(a_1, \dots, a_n) \in \mathbb{N}^n$, p_k, q_k are defined recursively by (2) for $0 \leq k \leq n$. Then

$$I_n(a_1, \dots, a_n) = \begin{cases} \left(\frac{p_n}{q_n}, \frac{p_n + p_{n-1}}{q_n + q_{n-1}} \right) & \text{if } n \text{ is even} \\ \left(\frac{p_n + p_{n-1}}{q_n + q_{n-1}}, \frac{p_n}{q_n} \right) & \text{if } n \text{ is odd.} \end{cases} \tag{4}$$

Therefore, the length of a cylinder of order n is given by

$$|I_n(a_1, \dots, a_n)| = \frac{1}{q_n(q_n + q_{n-1})}.$$

The next lemma relates a ball with the cylinders, basically following from Proposition 2 on the distribution of cylinders [11].

Proposition 3. Let $x \in I_n(a_1, \dots, a_n)$ with $a_n \geq 2$. Then

$$B(x, |I_n(a_1, \dots, a_n)|) \subset \bigcup_{i=-1}^2 I_n(a_1, \dots, a_n + i).$$

Next, we introduce the mass distribution principle which is a classic method in estimating the Hausdorff dimension of a set from below.

Proposition 4 ([4]). Let E be a Borel set and μ be a measure with $\mu(E) > 0$. Suppose that for any $x \in E$,

$$\liminf_{r \rightarrow 0} \frac{\log \mu(B(x, r))}{\log r} \geq s \tag{5}$$

where $B(x, r)$ denotes an open ball centered at x and radius r , then $\dim_H E \geq s$.

3. Proof of Theorem 1

Good [6] showed that the Hausdorff dimension of G defined in (1) is one-half, so by the simple inclusion

$$G \supset \mathcal{S}_G, \text{ and } G \supset \mathcal{L}_G \supset \mathcal{P}_G \supset \mathcal{E}_G,$$

it is sufficient to show that

$$\dim_H \mathcal{S}_G \geq 1/2 \text{ and } \dim_H \mathcal{E}_G \geq 1/2.$$

Lemma 5. Let $\alpha > 1$. Define

$$E_\alpha = \left\{ x \in [0, 1) : (2n - 1)^\alpha \leq a_n(x) < (2n)^\alpha, \text{ for all } n \geq 1 \right\}.$$

Then

$$\dim_H E_\alpha = \frac{\alpha - 1}{2\alpha}.$$

Proof.

- (I) For the upper bound of $\dim_H E_\alpha$, we consider a natural cover of E_α . It is clear that for any $N \geq 1$, the family of intervals

$$\bigcup_{a_1, \dots, a_N : (2n-1)^\alpha \leq a_n(x) < (2n)^\alpha, 1 \leq n \leq N} I_N(a_1, \dots, a_N)$$

covers E_α . Therefore, for any $s > 0$, the s -Hausdorff measure of E_α can be estimated from above by

$$\begin{aligned} \mathcal{H}^s(E_\alpha) &\leq \liminf_{N \rightarrow \infty} \sum_{a_1, \dots, a_N: (2n-1)^\alpha \leq a_n(x) < (2n)^\alpha, 1 \leq n \leq N} |I_N(a_1, \dots, a_n)|^s \\ &\leq \liminf_{N \rightarrow \infty} \sum_{a_1, \dots, a_N: (2n-1)^\alpha \leq a_n < (2n)^\alpha, 1 \leq n \leq N} \prod_{n=1}^N \frac{1}{a_n^{2s}} \\ &= \liminf_{N \rightarrow \infty} \prod_{n=1}^N \sum_{(2n-1)^\alpha \leq a_n < (2n)^\alpha} \frac{1}{a_n^{2s}}. \end{aligned}$$

For any $s > \frac{\alpha-1}{2\alpha}$, one can choose n_o such that for all $n \geq n_o$,

$$\sum_{(2n-1)^\alpha \leq a_n < (2n)^\alpha} \frac{1}{a_n^{2s}} \leq \frac{(2n)^\alpha - (2n-1)^\alpha}{(2n-1)^{2\alpha s}} \leq \frac{2\alpha \cdot (2n)^{\alpha-1}}{(2n-1)^{2\alpha s}} < 1,$$

and thus it follows that

$$\mathcal{H}^s(E_\alpha) \leq \prod_{n=1}^{n_o} \sum_{(2n-1)^\alpha \leq a_n < (2n)^\alpha} \frac{1}{a_n^{2s}} < \infty.$$

As a result,

$$\dim_H E_\alpha \leq \frac{\alpha-1}{2\alpha}.$$

(II) For the lower bound of $\dim_H E_\alpha$, notice that the set E_α has a nice Cantor structure. For each $n \geq 1$, let

$$\mathcal{E}_n = \left\{ I_n(a_1, \dots, a_n) : (2k-1)^\alpha \leq a_k < (2k)^\alpha, \text{ for all } 1 \leq k \leq n \right\}.$$

Then

$$E = \bigcap_{n=1}^{\infty} \bigcup_{I_n(a_1, \dots, a_n) \in \mathcal{E}_n} I_n(a_1, \dots, a_n),$$

and every element in $I_{n-1}(a_1, \dots, a_{n-1})$ in \mathcal{E}_{n-1} contains exactly

$$c_1 n^{\alpha-1} \leq D_n := (2n)^\alpha - (2n-1)^\alpha \leq c_2 n^{\alpha-1}.$$

elements $I_n(a_1, \dots, a_n)$ in \mathcal{E}_n .

Then, we distribute a mass supported on E_α by setting

$$\mu(I_n(a_1, \dots, a_n)) = \frac{1}{D_1 \cdots D_n},$$

for all $n \geq 1$ and $I_n(a_1, \dots, a_n)$ in \mathcal{E}_n .

For any $x \in E_\alpha$, for each $r > 0$ small enough, let n be the integer such that

$$|I_{n+1}(x)| < r \leq |I_n(x)|.$$

Then, by Proposition 3, the ball $B(x, r)$ can intersect at most three cylinders of order n , thus

$$\liminf_{r \rightarrow \infty} \frac{\log \mu(B(x, r))}{\log r} \geq \liminf_{n \rightarrow \infty} \frac{\log[3 \cdot (D_1 \cdots D_n)^{-1}]}{\log |I_{n+1}(x)|}.$$

By the recursive relation of $q_n(x)$, it follows that

$$q_{n+1}(x) \leq \prod_{k=1}^{n+1} (a_k(x) + 1) \leq \prod_{k=1}^{n+1} (2k)^\alpha \leq c^n \prod_{k=1}^{n+1} k^\alpha;$$

on the other hand,

$$\prod_{k=1}^n D_k \geq c_1^n \prod_{k=1}^n k^{\alpha-1}.$$

Thus it follows that

$$\liminf_{r \rightarrow 0} \frac{\log \mu((x, r))}{\log r} \geq \liminf_{n \rightarrow \infty} \frac{\log \prod_{k=1}^n k^{\alpha-1}}{\log (\prod_{k=1}^{n+1} k^\alpha)^2} = \frac{\alpha-1}{2\alpha}.$$

Finally by Proposition 4, one has

$$\dim_{\text{H}} E_\alpha \geq \frac{\alpha-1}{2\alpha}. \quad \square$$

By a result in [5] or the same line of the argument as above, except minor modifications on notation, one can show the following.

Lemma 6. *Let $b > 1$. Define*

$$F_b = \left\{ x \in [0, 1) : b^{1+2+\dots+n} \leq a_n(x) < 2 \cdot b^{1+2+\dots+n}, \text{ for all } n \geq 1 \right\}.$$

Then $\dim_{\text{H}} F_b = 1/2$.

Proof of Theorem 1. It is clear that for any $\alpha > 1$,

$$E_\alpha \subset \mathcal{S}_G, \text{ and so, } \dim_{\text{H}} \mathcal{S}_G \geq \lim_{\alpha \rightarrow \infty} \frac{\alpha-1}{2\alpha} = \frac{1}{2}.$$

It is also clear that for any $b > 1$,

$$F_b \subset \mathcal{E}_G, \text{ and so, } \dim_{\text{H}} \mathcal{E}_G \geq \frac{1}{2}. \quad \square$$

Declaration of interests

The authors do not work for, advise, own shares in, or receive funds from any organization that could benefit from this article, and have declared no affiliations other than their research organizations.

References

- [1] A. Bakhtawar, P. Bos and M. Hussain, “Hausdorff dimension of an exceptional set in the theory of continued fractions”, *Nonlinearity* **33** (2020), no. 6, pp. 2615–2639.
- [2] A. Bakhtawar, P. Bos and M. Hussain, “The sets of Dirichlet non-improvable numbers versus well-approximable numbers”, *Ergodic Theory Dyn. Syst.* **40** (2020), no. 12, pp. 3217–3235.
- [3] P. Bos, M. Hussain and D. Simmons, “The generalised Hausdorff measure of sets of Dirichlet non-improvable numbers”, *Proc. Am. Math. Soc.* **151** (2023), no. 5, pp. 1823–1838.
- [4] K. Falconer, *Fractal geometry*, John Wiley & Sons, 1990, pp. xxii+288. Mathematical foundations and applications.
- [5] A. H. Fan, L. M. Liao, B. W. Wang and J. Wu, “On Khintchine exponents and Lyapunov exponents of continued fractions”, *Ergodic Theory Dyn. Syst.* **29** (2009), no. 1, pp. 73–109.
- [6] I. J. Good, “The fractional dimensional theory of continued fractions”, *Proc. Camb. Philos. Soc.* **37** (1941), pp. 199–228.
- [7] L. L. Huang, J. Wu and J. Xu, “Metric properties of the product of consecutive partial quotients in continued fractions”, *Isr. J. Math.* **238** (2020), no. 2, pp. 901–943.
- [8] M. Hussain, D. Kleinbock, N. Wadleigh and B. W. Wang, “Hausdorff measure of sets of Dirichlet non-improvable numbers”, *Mathematika* **64** (2018), no. 2, pp. 502–518.
- [9] M. Hussain, B. Li and N. Shulga, “Hausdorff dimension analysis of sets with the product of consecutive vs single partial quotients in continued fractions”, *Discrete Contin. Dyn. Syst.* **44** (2024), no. 1, pp. 154–181.

- [10] M. Hussain and N. Shulga, “Metrical properties of exponentially growing partial quotients”, 2023. <https://arxiv.org/abs/2309.10529>.
- [11] M. Iosifescu and C. Kraaikamp, *Metrical theory of continued fractions*, Kluwer Academic Publishers, 2002, pp. xx+383.
- [12] V. Jarník, “A contribution to the metric theory of Diophantine approximations”, *Prace Mat.-Fiz.* **36** (1929), pp. 91–106.
- [13] A. Y. Khinchin, *Continued fractions*, The University of Chicago Press, 1964, pp. xi+95.
- [14] D. Kleinbock and N. Wadleigh, “A zero-one law for improvements to Dirichlet’s Theorem”, *Proc. Am. Math. Soc.* **146** (2018), no. 5, pp. 1833–1844.
- [15] B. Li, B. W. Wang and J. Xu, “Hausdorff dimension of Dirichlet non-improvable set versus well-approximable set”, *Ergodic Theory Dyn. Syst.* **43** (2023), no. 8, pp. 2707–2731.
- [16] T. Łuczak, “On the fractional dimension of sets of continued fractions”, *Mathematika* **44** (1997), no. 1, pp. 50–53.
- [17] R. D. Mauldin and M. Urbański, “Conformal iterated function systems with applications to the geometry of continued fractions”, *Trans. Am. Math. Soc.* **351** (1999), no. 12, pp. 4995–5025.
- [18] B. W. Wang and J. Wu, “Hausdorff dimension of certain sets arising in continued fraction expansions”, *Adv. Math.* **218** (2008), no. 5, pp. 1319–1339.