



ACADÉMIE
DES SCIENCES
INSTITUT DE FRANCE

Comptes Rendus

Mathématique


Khadijeh Baghaei

Global boundedness of solutions to a chemotaxis consumption model with signal dependent motility and logistic source

Volume 362 (2024), p. 1131-1145

Online since: 5 November 2024

<https://doi.org/10.5802/crmath.605>

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*The Comptes Rendus. Mathématique are a member of the
Mersenne Center for open scientific publishing*
www.centre-mersenne.org — e-ISSN : 1778-3569



Research article / *Article de recherche*

Partial differential equations / *Équations aux dérivées partielles*

Global boundedness of solutions to a chemotaxis consumption model with signal dependent motility and logistic source

Limite globale des solutions d'un modèle de consommation de chimiotaxie avec motilité dépendante du signal et source logistique

Khadijeh Baghaei

Abstract. This paper deals with the following chemotaxis system:

$$\begin{cases} u_t = \nabla \cdot (\gamma(v) \nabla u - u \xi(v) \nabla v) + \mu u(1 - u), & x \in \Omega, t > 0, \\ v_t = \Delta v - uv, & x \in \Omega, t > 0, \end{cases}$$

under homogeneous Neumann boundary conditions in a bounded domain $\Omega \subset \mathbb{R}^n$, $n \geq 2$, with smooth boundary. Here, the functions $\gamma(v)$ and $\xi(v)$ are as:

$$\gamma(v) = (1 + v)^{-k} \quad \text{and} \quad \xi(v) = -(1 - \alpha) \gamma'(v),$$

where $k > 0$ and $\alpha \in (0, 1)$.

For the above system, we prove that the corresponding initial boundary value problem admits a unique global classical solution which is uniformly-in-time bounded. This result is obtained under some conditions on initial value v_0 and μ and without any restriction on k and α . The obtained result extends the recent results obtained for this problem.

Résumé. Cet article porte sur le système de chimiotaxie suivant :

$$\begin{cases} u_t = \nabla \cdot (\gamma(v) \nabla u - u \xi(v) \nabla v) + \mu u(1 - u), & x \in \Omega, t > 0, \\ v_t = \Delta v - uv, & x \in \Omega, t > 0, \end{cases}$$

sous des conditions aux limites homogènes de Neumann dans un domaine borné $\Omega \subset \mathbb{R}^n$, $n \geq 2$, avec une frontière lisse. Ici, les fonctions $\gamma(v)$ et $\xi(v)$ sont les suivantes :

$$\gamma(v) = (1 + v)^{-k} \quad \text{and} \quad \xi(v) = -(1 - \alpha) \gamma'(v),$$

où $k > 0$ and $\alpha \in (0, 1)$.

Pour le système ci-dessus, nous prouvons que le problème de valeur limite initiale correspondant admet une unique solution classique globale qui est uniformément bornée en temps. Ce résultat est obtenu sous certaines conditions sur la valeur initiale v_0 et μ et sans restriction sur k et α . Le résultat obtenu étend les résultats récents obtenus pour ce problème.

Funding. This research was supported by a grant from PIAIS (No. 1402-10108).

Manuscript received 9 August 2023, revised 20 December 2023, accepted 1 January 2024.

1. Introduction

In this paper, we consider the following initial boundary value problem:

$$\begin{cases} u_t = \nabla \cdot (\gamma(v) \nabla u - u \xi(v) \nabla v) + \mu u(1 - u), & x \in \Omega, t > 0, \\ v_t = \Delta v - uv, & x \in \Omega, t > 0, \\ \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0, & x \in \partial\Omega, t > 0, \\ u(x, 0) = u_0, v(x, 0) = v_0, & x \in \Omega, \end{cases} \quad (1.1)$$

where $\Omega \subset \mathbb{R}^n$, $n \geq 2$, is a bounded domain with smooth boundary, ν denotes the unit outward normal vector to $\partial\Omega$ and u_0 and v_0 are initial functions. In the above problem, $u = u(x, t)$ is the cell density and $v = v(x, t)$ denotes the nutrient consumed chemical concentrations.

In mathematical biology, systems such as (1.1) describe the mechanism of chemotaxis. The chemotaxis is the movement of cells towards a higher concentration a chemical signal substance produced by cells. We first state the results related to the classical chemotaxis system, which has been introduced by Keller and Segel in [15]. The classical chemotaxis system can be written as follows:

$$\begin{cases} u_t = \nabla \cdot (\gamma(v) \nabla u - u \xi(v) \nabla v) + \mu u(1 - u), & x \in \Omega, t > 0, \\ \tau v_t = \Delta v - v + u, & x \in \Omega, t > 0, \end{cases} \quad (1.2)$$

where $\tau \in \{0, 1\}$. In the following, we write some important results for this problem in the absence and presence of the logistic source, respectively. In the absence of logistic source, when $\gamma \in C^3((0, \infty))$ and $\xi = -\gamma'$ as well as

$$\limsup_{s \rightarrow \infty} \gamma(s) < \frac{1}{\tau},$$

then the problem (1.2) admits a unique global classical solution in any dimension [33]. Also, for $n \geq 1$, when the function γ has strictly positive upper and lower bounds, then the classical solutions are uniformly-in-time bounded [33]. This result is also proved when $n \geq 2$ and the function γ decays at a certain slow rate at infinity [33]. Among the special functions that have been studied as γ are:

$$\gamma(v) = c_0 v^{-k} \quad \text{with } k > 0 \quad \text{and } c_0 > 0$$

and

$$\gamma(v) = e^{-\chi v} \quad \text{with } \chi > 0.$$

For $\gamma(v) = c_0 v^{-k}$, for all $k > 0$, the global existence and boundedness of the solution is proved under a smallness assumption on c_0 in any dimension [34]. Also, for this function, by removing the assumption on c_0 , when $n \geq 2$ and $k \in (0, \frac{2}{n-2})$, the same result is proved in cases $\tau = 0$ [1] and $\tau = 1$ [10]. In the other special case $\gamma(v) = e^{-\chi v}$, if $n = 2$ and $\int_{\Omega} u_0 dx < \frac{4\pi}{\chi}$, then the classical solutions are global and bounded whereas for $\int_{\Omega} u_0 dx > \frac{4\pi}{\chi}$, blow up occurs either in finite or infinite time [14]. But, in the case of $\tau = 0$, the blow up occurs in infinite time [11]. Also, for $n = 2$, the classical solutions are globally bounded if the positive function γ decreases slower than an exponential speed at high signal concentrations [9]. For $n \geq 3$, the same result is true when γ decreases at certain algebraically speed [9]. We now state the result obtained in [8] which is the global existence of very weak solution to the problem (1.2) when $\gamma(v) = \frac{1}{c+v^k}$ with $c \geq 0$ and $k > 0$. This result was obtained without any smallness assumption on the initial data provided that

$$k \in \begin{cases} (0, \frac{7}{3}), & \text{if } n = 1, \\ (0, 2), & \text{if } n = 2, \\ (0, \frac{4}{3}), & \text{if } n = 3. \end{cases}$$

Also, for $n \geq 1$ with $\gamma(v) = e^{-v}$, the existence of weak-strong solutions is proved in [6]. We now state the results to the problem (1.2) in the presence of logistic source. If the decreasing function $\gamma \in C^3([0, \infty))$ satisfies $\lim_{s \rightarrow \infty} \gamma(s) = 0$ and $\lim_{s \rightarrow \infty} \frac{\gamma'(s)}{\gamma(s)}$ exists, then the classical solutions are global and bounded for $n = 2$ [13]. For $n \geq 3$, the same result is true when $\mu > 0$ is large and the last condition is replaced with $|\gamma'(s)| \leq m$, where m is some positive constant [28]. Also, in two dimensional case, the same result holds when $\gamma \in C^3((0, \infty))$ and $|\gamma'| + |\gamma''| \in L^\infty((0, \infty))$ [31]. Now, we write some important results related to the problem (1.1). The origin of the definition of this problem comes from the following chemotaxis-Navier–Stokes system which describes the motion of oxygen-driven swimming bacteria in an in-compressible fluid

$$\begin{cases} u_t + \omega \cdot \nabla u = \nabla \cdot (\nabla u - u \xi(v) \nabla v), & x \in \Omega, t > 0, \\ v_t + \omega \cdot \nabla v = \Delta v - u g(v), & x \in \Omega, t > 0, \\ \omega_t + (\omega \cdot \nabla) \omega = \Delta \omega - \nabla P + u \nabla \phi, & x \in \Omega, t > 0, t > 0, \\ \nabla \cdot \omega = 0, & x \in \Omega, t > 0, t > 0. \end{cases}$$

Here, u denotes the bacteria density and v is the oxygen concentration. Also, ω and P are the velocity and pressure of the fluid, respectively. The function ξ measures the chemotactic sensitivity, g is the consumption rate of the oxygen by the bacteria, and ϕ is a given potential function [27]. For the related results with the chemotaxis-Navier–Stokes systems, we refer the interested readers to [5, 7, 12, 30] and references therein. We see that the problem (1.1) can be obtained from the preceding chemotaxis-Navier–Stokes system in the case of $\gamma(v) \equiv 1$ with the choice $\omega \equiv 0$ and $g(v) = v$. For the problem (1.1), in the absence of logistic source, when $\gamma(v) \equiv 1$, $\xi(v) \equiv \chi$, where χ is some positive constant, for $n = 2$, the classical solutions are global and bounded in bounded convex domains with a smooth boundary [25]. Also, for $n \geq 3$, this result holds in bounded domains with a smooth boundary provided that $\|v_0\|_{L^\infty(\Omega)} \leq \frac{1}{6(n+1)\chi}$ [24]. Later, this condition was extended to $\|v_0\|_{L^\infty(\Omega)} \leq \frac{\pi}{\chi \sqrt{2(n+1)}}$ and the same result was obtained in the absence of logistic source [3] and the presence of logistic source [4].

We now state the related results to the problem (1.1) in the absence of logistic source, when the function γ is not constant. We begin by stating the results obtained in [20] which is for the positive function $\gamma \in C^0([0, \infty))$ and $\xi = -\gamma'$. For non-negative initial data from $(C^0(\overline{\Omega}))^* \times L^\infty(\Omega)$, this problem admits a global very weak solution in all dimensions, also, for $\gamma \in C^1([0, \infty))$, the solutions stabilize toward a semi-trivial spatially homogeneous steady state in the large time limit. This result is obtained for $n \leq 3$ in [20] and later for $n \geq 1$ in [18]. If the decreasing function γ belongs to $C^3([0, \infty))$ and $\xi = -\gamma'$, then for non-negative initial data from $C^0(\overline{\Omega}) \times W^{1,\infty}(\overline{\Omega})$ under a smallness assumption on $\|v_0\|_{L^\infty(\Omega)}$, there exists a unique global classical solution that is bounded [19]. If the initial data (u_0, v_0) belongs to $(W^{1,\infty}(\Omega))^2$ and $\gamma \in C^3([0, \infty))$, then classical solutions are globally bounded when $n \leq 2$, and weak solutions are global when $n \geq 3$, in particular, such weak solutions become eventually smooth if $n = 3$ [21]. If $\gamma \in C^0([0, \infty)) \cap C^3([0, \infty))$, $\gamma(s) > 0$ for all $s > 0$ as well as $\xi = -\gamma'$, and γ satisfies:

$$\liminf_{s \searrow 0} \frac{\gamma(s)}{s^\alpha} > 0 \quad \text{and} \quad \limsup_{s \searrow 0} s^\beta |\gamma'(s)| < \infty$$

with $\alpha > 0$ and $\beta > 0$, then this problem admits a global generalized solution for all reasonably regular initial data [32]. In [26], the function γ is assumed to satisfy:

$$c_1 s^{-k} \leq \gamma(s) \leq c_2 s^{-k} \quad \text{for all } s > 0, \quad (*)$$

where k, c_1 and c_2 are some positive constants. Under this assumption for $\gamma \in C^3([0, \infty))$ and the initial data belongs to $(W^{1,\infty}(\Omega))^2$, the problem (1.1) admits a global classical solution when

$n = 1$, and a global weak-strong solution when $n \geq 2$. Also, this result is true if $2 \leq n \leq 5$, $k > \frac{n-2}{6-n}$ and γ satisfies (*) and

$$|\gamma'(s)| \leq c_3 s^{-k-1} \quad \text{for all } s > 0,$$

where c_3 is some positive constant [26].

Finally, we state the important results related to the problem (1.1) in the presence of logistic source. If $\gamma \in C^3([0, \infty))$ and $\gamma'(s) < 0$ for all $s \geq 0$ as well as $\xi = -\gamma'$, then for $n = 2$ and $\mu > 0$, the solutions are global and bounded [22]. Also, the same result is obtained when $n \geq 3$ and μ is suitably large [22]. Moreover, the solution converges exponentially to $(1, 0)$ when t tends to infinity [22]. For this problem, when the logistic source is as $f(u) = au - \mu u^\kappa$ with $a > 0$, $\mu > 0$ and $\kappa > 1$, the classical solutions are global and bounded if one of the cases ($n \leq 2, \kappa > 1$; $n \geq 3, \kappa > 2$ or $n \geq 3, \kappa = 2$ and μ is large) holds [29]. If the positive decreasing function γ belongs to $C^2([0, \infty))$ and $\gamma'' \geq 0$ as well as $\xi = -(1 - \alpha)\gamma'$ with $\alpha \in (0, 1)$, in [23], it is proved that the problem (1.1) admits a unique global classical solution that is uniformly in time bounded provided that:

$$\frac{(\gamma'(s))^2}{\gamma''(s)} \leq \frac{n}{2(n+1)^3}, \quad 0 < \|v_0\|_{L^\infty(\Omega)} \leq \gamma^{-1}\left(\frac{1}{n+1}\right)$$

and

$$\mu > \max_{0 < s \leq \|v_0\|_{L^\infty(\Omega)}} \frac{-\gamma'(s) \|v_0\|_{L^\infty(\Omega)}}{\gamma(s)}.$$

We see that the above conditions in the case of $\gamma(s) = (1 + s)^{-k}$ ($k > 0$) are as follows:

$$k < \frac{n}{2(n+1)^3 - n}, \quad \mu > k \|v_0\|_{L^\infty(\Omega)} \quad \text{and} \quad \|v_0\|_{L^\infty(\Omega)} \leq \gamma^{-1}\left(\frac{1}{n+1}\right).$$

Because of $\gamma' < 0$, the last condition is written as:

$$0 < \|v_0\|_{L^\infty(\Omega)} \leq (n+1)^{\frac{1}{k}} - 1.$$

In [2], we studied the special case $\gamma(s) = (1 + s)^{-k}$ ($k > 0$) and we were able to improve the conditions in [23]. In fact, we proved the same result in [23] under the following conditions:

$$k(1 - \alpha) < \frac{4}{n+5}, \quad 0 < \|v_0\|_{L^\infty(\Omega)} \leq \left[\frac{4[l-1]}{n+1}\right]^{\frac{1}{k}} - 1$$

and

$$\mu > \frac{n \|v_0\|_{L^\infty(\Omega)}}{l(n+1)(1 + \|v_0\|_{L^\infty(\Omega)})}$$

with $l = \frac{1}{k(1-\alpha)}$. In this paper, we focus again on the functions γ and ξ as follows:

$$\gamma(s) = (1 + s)^{-k} \quad \text{and} \quad \xi(s) = -(1 - \alpha)\gamma'(s), \tag{1.3}$$

where $k > 0$ and $\alpha \in (0, 1)$ and extend our recent result. In fact, we remove the condition on $k(1 - \alpha)$ and prove the solutions are uniformly in time bounded under some conditions on $\|v_0\|_{L^\infty(\Omega)}$ and μ .

2. Our results

Here, we state the standard well-posedness and classical solvability result.

Lemma 1. *Let $u_0 \geq 0$ and $v_0 \geq 0$ satisfy $(u_0, v_0) \in (W^{1,r}(\Omega))^2$ for some $r > n$. Then problem (1.1) has a unique local in time classical solution*

$$(u, v) \in \left(C([0, T_{\max}); W^{1,r}(\Omega)) \cap C^{2,1}(\overline{\Omega} \times (0, T_{\max}))\right)^2$$

where T_{\max} denotes the maximal existence time. In addition, if $T_{\max} < +\infty$, then:

$$\limsup_{t \rightarrow T_{\max}} \|u(\cdot, t)\|_{L^\infty(\Omega)} = +\infty.$$

Moreover, u and v satisfy the following inequalities:

$$u \geq 0 \quad \text{and} \quad 0 \leq v \leq \|v_0\|_{L^\infty(\Omega)} \quad \text{in} \quad \Omega \times (0, T_{\max}), \tag{2.1}$$

also,

$$\int_{\Omega} u(\cdot, t) \, dx \leq c, \tag{2.2}$$

where c is some positive constant.

For details of the proof, we refer the reader to [13, 23].

Based on main ideas in [3, 4, 16, 17], we write the following key lemma similar to [2, Lemma 2.2].

Lemma 2. *Let (u, v) be the solution of problem (1.1). If there exists a smooth positive function $\varphi(s)$ such that for $0 \leq s \leq \|v_0\|_{L^\infty(\Omega)}$, the following inequality holds:*

$$(B(s))^2 - 4 A(s) C(s) \leq 0, \tag{2.3}$$

where for $p \geq 2$, the functions A, B and C are defined as:

$$\begin{cases} A(s) = (p-1) \varphi(s) \gamma(s), \\ B(s) = (p-1) \varphi(s) \xi(s) - \varphi'(s) (\gamma(s) + 1), \\ C(s) = \frac{1}{p} \varphi''(s) - \varphi'(s) \xi(s), \end{cases} \tag{2.4}$$

then:

$$\frac{1}{p} \frac{d}{dt} \int_{\Omega} u^p \varphi(v) \, dx \leq - \int_{\Omega} \left[\mu \varphi(v) + \frac{1}{p} v \varphi'(v) \right] u^{p+1} \, dx + \mu \int_{\Omega} u^p \varphi(v) \, dx.$$

Proof. We assume that there exists a smooth positive function $\varphi(s)$ such that for $0 \leq s \leq \|v_0\|_{L^\infty(\Omega)}$ and $p \geq 2$, (2.3) holds. We take this function and use (1.1) and integration by parts to write:

$$\begin{aligned} \frac{1}{p} \frac{d}{dt} \int_{\Omega} u^p \varphi(v) \, dx &= \int_{\Omega} u^{p-1} \varphi(v) u_t \, dx + \frac{1}{p} \int_{\Omega} u^p \varphi'(v) v_t \, dx \\ &= -(p-1) \int_{\Omega} u^{p-2} \varphi(v) \gamma(v) |\nabla u|^2 \, dx \\ &\quad + \int_{\Omega} u^{p-1} [(p-1) \varphi(v) \xi(v) - \varphi'(v) (\gamma(v) + 1)] (\nabla u \cdot \nabla v) \, dx \\ &\quad + \int_{\Omega} u^p \left[\varphi'(v) \xi(v) - \frac{1}{p} \varphi''(v) \right] |\nabla v|^2 \, dx \\ &\quad - \int_{\Omega} \left[\mu \varphi(v) + \frac{1}{p} v \varphi'(v) \right] u^{p+1} \, dx + \mu \int_{\Omega} u^p \varphi(v) \, dx. \end{aligned} \tag{2.5}$$

For convenience in calculations, we write (2.5) as follows:

$$\frac{1}{p} \frac{d}{dt} \int_{\Omega} u^p \varphi(v) \, dx = \int_{\Omega} J(u, v) \, dx - \int_{\Omega} \left[\mu \varphi(v) + \frac{1}{p} v \varphi'(v) \right] u^{p+1} \, dx + \mu \int_{\Omega} u^p \varphi(v) \, dx \tag{2.6}$$

with

$$\begin{aligned} J(u, v) &= -(p-1) u^{p-2} \varphi(v) \gamma(v) |\nabla u|^2 \\ &\quad + u^{p-1} [(p-1) \varphi(v) \xi(v) - \varphi'(v) (\gamma(v) + 1)] (\nabla u \cdot \nabla v) \\ &\quad + u^p \left[\varphi'(v) \xi(v) - \frac{1}{p} \varphi''(v) \right] |\nabla v|^2 \\ &= -u^{p-2} |\nabla u|^2 A(v) + u^{p-1} (\nabla u \cdot \nabla v) B(v) - u^p |\nabla v|^2 C(v), \end{aligned} \tag{2.7}$$

where A, B and C are defined in (2.4). Now, by considering (2.7), we can write

$$\begin{aligned}
 J(u, v) &= -\left(\sqrt{u^{p-2}A(v)}\nabla u - \frac{u^{p-1}B(v)}{2\sqrt{u^{p-2}A(v)}}\nabla v\right) \cdot \left(\sqrt{u^{p-2}A(v)}\nabla u - \frac{u^{p-1}B(v)}{2\sqrt{u^{p-2}A(v)}}\nabla v\right) \\
 &\quad + u^p \left[\frac{(B(v))^2}{4A(v)} - C(v)\right] |\nabla v|^2 \\
 &\leq u^p \left[\frac{(B(v))^2 - 4A(v)C(v)}{4A(v)}\right] |\nabla v|^2.
 \end{aligned}$$

In view of the condition (2.3), we see that $J \leq 0$. Thus, the equality (2.6) becomes

$$\frac{1}{p} \frac{d}{dt} \int_{\Omega} u^p \varphi(v) \, dx \leq - \int_{\Omega} \left[\mu \varphi(v) + \frac{1}{p} v \varphi'(v) \right] u^{p+1} \, dx + \mu \int_{\Omega} u^p \varphi(v) \, dx.$$

This completes our proof. □

Before presenting a smooth positive function φ such that the relation (2.3) holds, we present the following preliminary lemma.

Lemma 3. *Let the function γ for $s \geq 0$ is defined as $\gamma(s) = (1 + s)^{-k}$ with $k > 0$. If the function H for $z > 0$ is defined as*

$$H(z) = d^2 p z^{2\lambda} - 2 d^2 (p-2) z^{2\lambda-1} + d^2 p z^{2\lambda-2} - 2 d [2 d (p-1) - p + 2 l] z^{\lambda-1} - 2 d p z^{\lambda} + p, \quad (2.8)$$

where for $\alpha \in (0, 1)$ and $p > 2$, the parameters λ, l and d are as follows:

$$\lambda = d(p-1)(1-\alpha), \quad l = \frac{1}{k(1-\alpha)}$$

and

$$d > \frac{-l + \sqrt{l^2 + p(p-2)}}{2(p-2)}. \quad (2.9)$$

Then there exists $\delta_0 > 0$ such that for $0 < s \leq \delta_0$, $H(\gamma(s)) \leq 0$.

Proof. At first, we see that

$$\begin{aligned}
 H(1) &= d^2 p - 2 d^2 (p-2) + d^2 p - 2 d [2 d (p-1) - p + 2 l] - 2 d p + p \\
 &= -4(p-2) d^2 - 4 l d + p.
 \end{aligned}$$

It is not difficult to see that the choice of d as (2.9) implies that:

$$-4(p-2) d^2 - 4 l d + p < 0.$$

Thus, $H(1) < 0$. By considering the continuity of the function H on $(0, \infty)$ and γ on $[0, \infty)$, and also $H(\gamma(0)) = H(1) < 0$, we conclude that there exists $\delta_0 > 0$ such that for $0 < s \leq \delta_0$, $H(\gamma(s)) \leq 0$. □

We now present a smooth positive function φ and show that for this function, the relation (2.3) holds. We must state that the following lemma is the only place where the special choice of γ is used.

Lemma 4. *Assume that the initial values u_0 and v_0 are non-negative and satisfy:*

$$(u_0, v_0) \in (W^{1,r}(\Omega))^2 \quad \text{for some } r > n,$$

and

$$0 < \|v_0\|_{L^\infty(\Omega)} \leq \delta_0,$$

which δ_0 is introduced in Lemma 3. Also, assume that the functions $\gamma(v)$ and $\xi(v)$ are defined as (1.3) and for $p > n$, the following condition holds:

$$\mu > \frac{d(p-1) \|v_0\|_{L^\infty(\Omega)}}{l p (1 + \|v_0\|_{L^\infty(\Omega)})}, \quad (2.10)$$

where $l = \frac{1}{k(1-\alpha)}$ and d is chosen as (2.9). Then, there exists some positive constant c which also depends on δ_0 and p such that the following estimate holds:

$$\|u(\cdot, t)\|_{L^p(\Omega)} \leq c, \quad \forall t \in (0, T_{\max}). \tag{2.11}$$

Proof. We prove this lemma in two steps. In the first step, we present a smooth positive function φ and show that for this function (2.3) holds. In the second step, we prove (2.11) holds.

Step 1. As a starting point, in Lemma 2, we take $p > n$ and define the smooth positive function φ as follows:

$$\varphi(v) = e^{(\gamma(v))^\lambda}$$

with $\lambda = d(p-1)(1-\alpha)$, where d is taken as (2.9). For this function, we have:

$$\varphi'(v) = \lambda (\gamma(v))^{\lambda-1} \gamma'(v) \varphi(v) \tag{2.12}$$

and

$$\varphi''(v) = \lambda [(\lambda-1)(\gamma'(v))^2 + \gamma(v)\gamma''(v) + \lambda(\gamma(v))^\lambda (\gamma'(v))^2] (\gamma(v))^{\lambda-2} \varphi(v).$$

Because of

$$\gamma'(v) = -k(1+v)^{-k-1} \quad \text{and} \quad \gamma''(v) = k(k+1)(1+v)^{-k-2},$$

we see that $\gamma(v)\gamma''(v) = (1 + \frac{1}{k})(\gamma'(v))^2$. Thus,

$$\varphi''(v) = \lambda \left[\lambda + \frac{1}{k} + \lambda(\gamma(v))^\lambda \right] (\gamma(v))^{\lambda-2} (\gamma'(v))^2 \varphi(v). \tag{2.13}$$

By considering the values of A, B and C from (2.4), we can write:

$$\begin{aligned} (B(v))^2 - 4A(v)C(v) &= (p-1)^2(\varphi(v))^2(\xi(v))^2 + (\varphi'(v))^2(\gamma(v)+1)^2 \\ &\quad - \frac{4(p-1)}{p} \varphi(v)\varphi''(v)\gamma(v) - 2(p-1)\varphi(v)\varphi'(v)\xi(v)(1-\gamma(v)). \end{aligned}$$

Making use of (2.12), (2.13) and $\xi(v) = -(1-\alpha)\gamma'(v)$, we obtain:

$$\begin{aligned} (B(v))^2 - 4A(v)C(v) &= (\gamma'(v))^2(\varphi(v))^2 \left\{ (p-1)^2(1-\alpha)^2 + \lambda^2(\gamma(v))^{2\lambda-2}(\gamma(v)+1)^2 \right. \\ &\quad \left. - \frac{4\lambda(p-1)}{p} \left[\lambda + \frac{1}{k} + \lambda(\gamma(v))^\lambda \right] (\gamma(v))^{\lambda-1} + 2\lambda(p-1)(1-\alpha)(\gamma(v))^{\lambda-1}(1-\gamma(v)) \right\}. \end{aligned}$$

Because of $\lambda = d(p-1)(1-\alpha)$, we replace $(p-1)(1-\alpha)$ with $\frac{\lambda}{d}$ to have:

$$\begin{aligned}
 & (B(v))^2 - 4A(v)C(v) \\
 &= (\gamma'(v))^2(\varphi(v))^2 \left\{ \frac{\lambda^2}{d^2} + \lambda^2(\gamma(v))^{2\lambda-2}(\gamma(v)+1)^2 \right. \\
 &\quad \left. - \frac{4\lambda^2}{d p(1-\alpha)} \left[\lambda(1+(\gamma(v))^\lambda) + \frac{1}{k} \right] (\gamma(v))^{\lambda-1} + \frac{2\lambda^2}{d} (\gamma(v))^{\lambda-1}(1-\gamma(v)) \right\} \\
 &= \frac{\lambda^2(\gamma'(v))^2(\varphi(v))^2}{d^2 p} \left\{ p + d^2 p(\gamma(v))^{2\lambda-2}(\gamma(v)+1)^2 \right. \\
 &\quad \left. - 4d \left[d(p-1)(1+(\gamma(v))^\lambda) + \frac{1}{k(1-\alpha)} \right] (\gamma(v))^{\lambda-1} + 2d p(\gamma(v))^{\lambda-1}(1-\gamma(v)) \right\} \quad (2.14) \\
 &= \frac{\lambda^2(\gamma'(v))^2(\varphi(v))^2}{d^2 p} \left\{ p + d^2 p(\gamma(v))^{2\lambda} - 2d^2(p-2)(\gamma(v))^{2\lambda-1} \right. \\
 &\quad \left. + d^2 p(\gamma(v))^{2\lambda-2} - 2d[2d(p-1) - p + 2l](\gamma(v))^{\lambda-1} - 2d p(\gamma(v))^\lambda \right\} \\
 &:= \frac{\lambda^2}{d^2 p} (\gamma'(v))^2(\varphi(v))^2 H(\gamma(v)),
 \end{aligned}$$

where the function H is defined by (2.8). By considering Lemma 3, we see that there exists $\delta_0 > 0$ such that for $0 < v \leq \delta_0$, $H(\gamma(v)) \leq 0$. Because of $0 < \|v_0\|_{L^\infty(\Omega)} \leq \delta_0$, we conclude that $H(\gamma(v))$ is non-positive in the interval $(0, \|v_0\|_{L^\infty(\Omega)})$. Therefore, by combining this with (2.14), we conclude that the relation (2.3) holds.

Step 2. In order to prove (2.11), we apply Lemma 2 and write:

$$\begin{aligned}
 & \frac{1}{p} \frac{d}{dt} \int_{\Omega} u^p \varphi(v) \, dx + \mu \int_{\Omega} u^p \varphi(v) \, dx \\
 & \leq - \int_{\Omega} \left[\mu \varphi(v) + \frac{1}{p} v \varphi'(v) \right] u^{p+1} \, dx + 2\mu \int_{\Omega} u^p \varphi(v) \, dx. \quad (2.15)
 \end{aligned}$$

We now apply the Young inequality to the second term on the right hand side of (2.15) to have:

$$2\mu \int_{\Omega} u^p \varphi(v) \, dx \leq \epsilon \int_{\Omega} u^{p+1} \varphi(v) \, dx + c_\epsilon \int_{\Omega} \varphi(v) \, dx,$$

where ϵ is chosen as follows:

$$0 < \epsilon < \mu - \frac{k\lambda \|v_0\|_{L^\infty(\Omega)}}{p(1 + \|v_0\|_{L^\infty(\Omega)})} \quad (2.16)$$

and:

$$c_\epsilon = \frac{1}{p+1} \left[\frac{p}{\epsilon(p+1)} \right]^p (2\mu)^{p+1}.$$

Because of $\gamma'(v) < 0$, (2.12) implies that $\varphi'(v) < 0$. Thus,

$$\varphi(\|v_0\|_{L^\infty(\Omega)}) < \varphi(v) < \varphi(0). \quad (2.17)$$

Making use of this, we can write:

$$2\mu \int_{\Omega} u^p \varphi(v) \, dx \leq \epsilon \int_{\Omega} u^{p+1} \varphi(v) \, dx + c_0 \quad (2.18)$$

with $c_0 = c_\epsilon |\Omega| \varphi(0)$. We now combine the inequality (2.18) with (2.15) to obtain:

$$\frac{1}{p} \frac{d}{dt} \int_{\Omega} u^p \varphi(v) \, dx + \mu \int_{\Omega} u^p \varphi(v) \, dx \leq \int_{\Omega} \left[(\epsilon - \mu) \varphi(v) - \frac{1}{p} v \varphi'(v) \right] u^{p+1} \, dx + c_0. \quad (2.19)$$

We use $\gamma'(v) = -\frac{k}{1+v}\gamma(v)$, $\gamma(v) \leq 1$ and (2.16) to have

$$\begin{aligned} (\epsilon - \mu)\varphi(v) - \frac{1}{p}v\varphi'(v) &= \left[\epsilon - \mu - \frac{\lambda}{p}v(\gamma(v))^{\lambda-1}\gamma'(v) \right] \varphi(v) \\ &= \left[\epsilon - \mu + \frac{k\lambda v}{p(1+v)}(\gamma(v))^\lambda \right] \varphi(v) \\ &\leq \left[\epsilon - \mu + \frac{k\lambda\|v_0\|_{L^\infty(\Omega)}}{p(1+\|v_0\|_{L^\infty(\Omega)})} \right] \varphi(v) \\ &\leq 0. \end{aligned}$$

This along with (2.19) yields:

$$\frac{d}{dt} \int_{\Omega} u^p \varphi(v) \, dx + \mu p \int_{\Omega} u^p \varphi(v) \, dx \leq p c_0.$$

We put:

$$y(t) = \int_{\Omega} u^p \varphi(v) \, dx.$$

Thus,

$$y'(t) + \mu p y(t) \leq p c_0.$$

This yields:

$$y(t) \leq \max \left\{ y(0), \frac{c_0}{\mu} \right\}. \tag{2.20}$$

This along with (2.17) allows us to write:

$$\int_{\Omega} u^p \, dx \leq (\varphi(\|v_0\|_{L^\infty(\Omega)}))^{-1} \max \left\{ y(0), \frac{c_0}{\mu} \right\} := c.$$

By considering the value of c_0 , we see that c also depends on δ_0 and p . This completes our proof. \square

The proof of the following lemma is the same as [24, Lemma 3.2] or [2, Lemma 2.4]. But, we write it to complement our content.

Lemma 5. *Assume that for $p > n$, the following estimate holds:*

$$\|u(\cdot, t)\|_{L^p(\Omega)} \leq C_1, \quad \forall t \in (0, T_{\max}), \tag{2.21}$$

where C_1 is some positive constant, then there exists some positive constant C_2 which also depends on C_1 and $\|v_0\|_{L^\infty(\Omega)}$ such that

$$\|\nabla v(\cdot, t)\|_{L^\infty(\Omega)} \leq C_2 \tag{2.22}$$

for all $t \in (0, T_{\max})$.

Proof. By considering Lemma 1, we see that it is sufficient to prove for any $\tau \in (0, T_{\max})$,

$$\|\nabla v(\cdot, t)\|_{L^\infty(\Omega)} \leq C_2 \quad \text{for all } t \in (\tau, T_{\max}). \tag{2.23}$$

We use the representation formula for the second equation (1.1) to have:

$$v(\cdot, t) = e^{t(\Delta-1)} v_0 + \int_0^t e^{(t-s)(\Delta-1)} (1 - u(\cdot, s)) v(\cdot, s) \, ds, \quad t \in (0, T_{\max}).$$

We now take $p > n$ and use $0 \leq v \leq \|v_0\|_{L^\infty(\Omega)}$ to write:

$$\begin{aligned} \|(1 - u(\cdot, s))v(\cdot, s)\|_{L^p(\Omega)} &\leq \|v(\cdot, s)\|_{L^\infty(\Omega)} \left(\int_{\Omega} |1 - u(\cdot, s)|^p dx \right)^{\frac{1}{p}} \\ &\leq \|v(\cdot, s)\|_{L^\infty(\Omega)} \left(\int_{\Omega} (1 + |u(\cdot, s)|)^p dx \right)^{\frac{1}{p}} \\ &\leq \|v(\cdot, s)\|_{L^\infty(\Omega)} \left(2^{p-1} \int_{\Omega} (1 + |u(\cdot, s)|)^p dx \right)^{\frac{1}{p}} \\ &\leq 2^{\frac{p-1}{p}} \|v(\cdot, s)\|_{L^\infty(\Omega)} \left(|\Omega|^{\frac{1}{p}} + \|u(\cdot, s)\|_{L^p(\Omega)} \right) \\ &\leq c, \end{aligned} \tag{2.24}$$

where we have used the inequality $(a + b)^m \leq 2^{m-1}(a^m + b^m)$ with $a, b \geq 0$ and $m > 1$, also $(a + b)^{m'} \leq (a^{m'} + b^{m'})$ with $0 < m' < 1$. We note that the constant c in the last estimate, it also depends on C_1 and $\|v_0\|_{L^\infty(\Omega)}$. In order to prove (2.23), we take $\tau \in (0, \min\{1, T_{\max}\})$ and $\theta \in (\frac{p+n}{2p}, 1)$ and use the estimates (3.16) and (3.17) in [24], also (2.24) to obtain:

$$\begin{aligned} \|v(\cdot, t)\|_{W^{1,\infty}(\Omega)} &\leq c \|(-\Delta + 1)^\theta v(\cdot, t)\|_{L^p(\Omega)} \\ &\leq c t^{-\theta} e^{-\delta t} \|v_0\|_{L^p(\Omega)} + c \int_0^t (t-s)^{-\theta} e^{-\delta(t-s)} \|(1 - u(\cdot, s))v(\cdot, s)\|_{L^p(\Omega)} ds \\ &\leq c t^{-\theta} + c \int_0^t (t-s)^{-\theta} e^{-\delta(t-s)} ds \\ &\leq c t^{-\theta} + c \int_0^{+\infty} \sigma^{-\theta} e^{-\delta\sigma} d\sigma \\ &\leq c(\tau^{-\theta} + 1), \quad t \in (\tau, T_{\max}), \end{aligned}$$

where the constants can vary from line to line. In this estimate, from the third line on-wards, the constants are also dependent on C_1 and $\|v_0\|_{L^\infty(\Omega)}$. This completes our proof. \square

The proof of the following lemma is the same as [2, Lemma 2.5] and similar to [24, Lemma 3.2]. But, we write it to complement our content.

Lemma 6. Assume that the initial values u_0 and v_0 are non-negative and satisfy:

$$(u_0, v_0) \in (W^{1,r}(\Omega))^2 \quad \text{for some } r > n.$$

Also, assume that (2.22) holds. Then there exists some positive constant c such that for all $t \in (0, T_{\max})$, the following estimate holds

$$\|u(\cdot, t)\|_{L^\infty(\Omega)} \leq c.$$

Proof. We take $q \geq 2$ and use from (1.1) and integration by parts to obtain:

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} u^q dx &= q \int_{\Omega} u^{q-1} [\nabla \cdot (\gamma(v)\nabla u - u\xi(v)\nabla v) + \mu u(1 - u)] dx \\ &= -q(q-1) \int_{\Omega} \gamma(v) u^{q-2} |\nabla u|^2 dx + q(q-1) \int_{\Omega} u^{q-1} \xi(v) (\nabla u \cdot \nabla v) dx \\ &\quad + \mu q \int_{\Omega} u^q (1 - u) dx. \end{aligned} \tag{2.25}$$

Because of $0 \leq v \leq \|v_0\|_{L^\infty(\Omega)}$, we have:

$$\begin{aligned} \gamma(v) &= (1 + v)^{-k} \geq (1 + \|v_0\|_{L^\infty(\Omega)})^{-k} := c_1, \\ \xi(v) &= k(1 - \alpha)(1 + v)^{-k-1} \leq k(1 - \alpha) := c_2. \end{aligned}$$

Making use of these, (2.22) and Young’s inequality, we can write (2.25) as follows:

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} u^q dx &\leq -c_1 q (q-1) \int_{\Omega} u^{q-2} |\nabla u|^2 dx + C_2 c_2 q (q-1) \int_{\Omega} u^{q-1} |\nabla u| dx + \mu q \int_{\Omega} u^q dx \\ &= -\frac{4c_1(q-1)}{q} \int_{\Omega} |\nabla u^{\frac{q}{2}}|^2 dx + 2C_2 c_2 (q-1) \int_{\Omega} u^{\frac{q}{2}} \cdot |\nabla u^{\frac{q}{2}}| dx + \mu q \int_{\Omega} u^q dx \quad (2.26) \\ &\leq -\frac{2c_1(q-1)}{q} \int_{\Omega} |\nabla u^{\frac{q}{2}}|^2 dx + q \left(\frac{(q-1)(C_2 c_2)^2}{2c_1} + \mu \right) \int_{\Omega} u^q dx. \end{aligned}$$

We now add $q \int_{\Omega} u^q dx$ on both sides of (2.26) to have:

$$\frac{d}{dt} \int_{\Omega} u^q dx + q \int_{\Omega} u^q dx \leq -\frac{2c_1(q-1)}{q} \int_{\Omega} |\nabla u^{\frac{q}{2}}|^2 dx + c_3 \int_{\Omega} u^q dx \quad (2.27)$$

with

$$c_3 = q \left(\frac{(q-1)(C_2 c_2)^2}{2c_1} + \mu + 1 \right).$$

We first apply the Gagliardo–Nirenberg inequality and then use the Young inequality with exponents $s = \frac{n+2}{2}$ and $s' = \frac{n+2}{n}$ to obtain:

$$\begin{aligned} c_3 \int_{\Omega} u^q dx &= c_3 \|u^{\frac{q}{2}}\|_{L^2(\Omega)}^2 \leq c_3 (C_{GN})^2 \left(\|\nabla u^{\frac{q}{2}}\|_{L^2(\Omega)}^{\frac{n}{n+2}} \|u^{\frac{q}{2}}\|_{L^1(\Omega)}^{\frac{2}{n+2}} + \|u^{\frac{q}{2}}\|_{L^1(\Omega)} \right)^2 \\ &\leq 2c_3 (C_{GN})^2 \left(\|\nabla u^{\frac{q}{2}}\|_{L^2(\Omega)}^{\frac{2n}{n+2}} \|u^{\frac{q}{2}}\|_{L^1(\Omega)}^{\frac{4}{n+2}} + \|u^{\frac{q}{2}}\|_{L^1(\Omega)}^2 \right) \\ &\leq \frac{2c_1(q-1)}{q} \|\nabla u^{\frac{q}{2}}\|_{L^2(\Omega)}^2 + (c_4 + 2c_3(C_{GN})^2) \|u^{\frac{q}{2}}\|_{L^1(\Omega)}^2 \\ &= \frac{2c_1(q-1)}{q} \int_{\Omega} |\nabla u^{\frac{q}{2}}|^2 dx + c_5 \left(\int_{\Omega} u^{\frac{q}{2}} dx \right)^2 \end{aligned}$$

with

$$c_4 = \frac{1}{s} \left(\frac{2c_1 s' (q-1)}{q} \right)^{-\frac{s}{s'}} (2c_3(C_{GN})^2)^s \quad \text{and} \quad c_5 = c_4 + 2c_3(C_{GN})^2,$$

where C_{GN} is the constant in the Gagliardo–Nirenberg inequality. Combining the last inequality with (2.27) yields:

$$\frac{d}{dt} \int_{\Omega} u^q dx + q \int_{\Omega} u^q dx \leq c_5 \left(\int_{\Omega} u^{\frac{q}{2}} dx \right)^2.$$

For $0 \leq t \leq T_{\max}$, we can write:

$$\frac{d}{dt} \left(e^{qt} \int_{\Omega} u^q dx \right) \leq c_5 e^{qt} \left(\int_{\Omega} u^{\frac{q}{2}} dx \right)^2.$$

Now, we integrate and use $e^{-qt} \leq 1$ to get:

$$\begin{aligned} \int_{\Omega} u^q dx &\leq \int_{\Omega} u_0^q dx + \frac{c_5}{q} \sup_{0 \leq t \leq T_{\max}} \left(\int_{\Omega} u^{\frac{q}{2}} dx \right)^2 \\ &\leq |\Omega| \|u_0\|_{L^\infty(\Omega)}^q + \frac{c_5}{q} \sup_{0 \leq t \leq T_{\max}} \left(\int_{\Omega} u^{\frac{q}{2}} dx \right)^2. \end{aligned}$$

Thus,

$$\begin{aligned} \left(\int_{\Omega} u^q dx \right)^{\frac{1}{q}} &\leq \left[|\Omega| \|u_0\|_{L^\infty(\Omega)}^q + \frac{c_5}{q} \sup_{0 \leq t \leq T_{\max}} \left(\int_{\Omega} u^{\frac{q}{2}} dx \right)^2 \right]^{\frac{1}{q}} \\ &\leq |\Omega|^{\frac{1}{q}} \|u_0\|_{L^\infty(\Omega)} + \left(\frac{c_5}{q} \right)^{\frac{1}{q}} \sup_{0 \leq t \leq T_{\max}} \left(\int_{\Omega} u^{\frac{q}{2}} dx \right)^{\frac{2}{q}}. \quad (2.28) \end{aligned}$$

We note that

$$\begin{aligned}
 c_5 &= c_4 + 2 c_3 (C_{GN})^2 \\
 &= \frac{1}{s} \left(\frac{2 c_1 s' (q-1)}{q} \right)^{-\frac{s}{s'}} (2 c_3 (C_{GN})^2)^s + 2 c_3 (C_{GN})^2 \\
 &= \frac{1}{s} (2 c_1 s')^{-\frac{s}{s'}} (2 (C_{GN})^2)^s \left(\frac{q}{q-1} \right)^{\frac{n}{2}} (c_3)^s + 2 c_3 (C_{GN})^2 \\
 &\leq m \left[\left(\frac{q}{q-1} \right)^{\frac{n}{2}} (c_3)^s + c_3 \right] \\
 &\leq m \left[\left(\frac{q}{q-1} \right)^{\frac{n}{2}} + 1 \right] (c_3)^s
 \end{aligned}$$

with

$$m = \max \left\{ \frac{1}{s} (2 c_1 s')^{-\frac{s}{s'}} (2 (C_{GN})^2)^s, 2 (C_{GN})^2 \right\}.$$

Here, we have used from $c_3 > 1$ and $s > 1$. By inserting c_3 and using $q \geq 2$, we obtain:

$$\begin{aligned}
 \frac{c_5}{q} &\leq m \left[\left(\frac{q}{q-1} \right)^{\frac{n}{2}} + 1 \right] \left(\frac{(q-1) (C_2 c_2)^2}{2 c_1} + \mu + 1 \right)^{\frac{n}{2}+1} q^{\frac{n}{2}} \\
 &\leq 2 m \left(\frac{(C_2 c_2)^2}{2 c_1} + \mu + 1 \right)^{\frac{n}{2}+1} \left(\frac{q}{q-1} \right)^{\frac{n}{2}} (q-1)^{\frac{n}{2}+1} q^{\frac{n}{2}} \\
 &= c_6 (q-1) q^n \\
 &\leq c_6 q^{n+1}
 \end{aligned} \tag{2.29}$$

with

$$c_6 = 2 m \left(\frac{(C_2 c_2)^2}{2 c_1} + \mu + 1 \right)^{\frac{n}{2}+1}.$$

Making use of (2.29) and $q^{\frac{n+1}{q}} > 1$, we can write (2.28) as follows:

$$\begin{aligned}
 \left(\int_{\Omega} u^q dx \right)^{\frac{1}{q}} &\leq |\Omega|^{\frac{1}{q}} \|u_0\|_{L^\infty(\Omega)} + (c_6 q^{n+1})^{\frac{1}{q}} \sup_{0 \leq t \leq T_{\max}} \left(\int_{\Omega} u^{\frac{q}{2}} dx \right)^{\frac{2}{q}} \\
 &\leq c_7^{\frac{1}{q}} q^{\frac{n+1}{q}} \left(\|u_0\|_{L^\infty(\Omega)} + \sup_{0 \leq t \leq T_{\max}} \left(\int_{\Omega} u^{\frac{q}{2}} dx \right)^{\frac{2}{q}} \right)
 \end{aligned} \tag{2.30}$$

with $c_7 = |\Omega| + c_6$. We now define:

$$M(q) = \max \left\{ \|u_0\|_{L^\infty(\Omega)}, \sup_{0 \leq t \leq T_{\max}} \left(\int_{\Omega} u^q dx \right)^{\frac{1}{q}} \right\}.$$

This allows us to write (2.30) as:

$$M(q) \leq 2 c_7^{\frac{1}{q}} q^{\frac{n+1}{q}} M\left(\frac{q}{2}\right).$$

We now take $q = 2^i$ ($i \in \mathbb{N}$) to obtain:

$$\begin{aligned}
 M(2^i) &\leq 2 c_7^{2^{-i}} 2^{\frac{(n+1)i}{2^i}} M(2^{i-1}) \\
 &\leq 2 c_7^{2^{-i}+2^{-i+1}} 2^{(n+1)\left(\frac{i}{2^i} + \frac{i-1}{2^{i-1}}\right)} M(2^{i-2}) \\
 &\leq \dots \\
 &\leq 2 c_7^{2^{-i}+2^{-i+1}+\dots+2^{-1}} 2^{(n+1)\left(\frac{i}{2^i} + \frac{i-1}{2^{i-1}} + \dots + \frac{1}{2}\right)} M(1),
 \end{aligned} \tag{2.31}$$

and compute the following elementary series:

$$S := \sum_{i=1}^{\infty} \frac{i}{2^i} = \sum_{i=0}^{\infty} \frac{i+1}{2^{i+1}} = \sum_{i=0}^{\infty} \left(\frac{i}{2^{i+1}} + \frac{1}{2^{i+1}} \right) = \frac{1}{2} \sum_{i=1}^{\infty} \frac{i}{2^i} + \sum_{i=0}^{\infty} \frac{1}{2^{i+1}} = \frac{1}{2} S + 1.$$

Thus, $S = 2$. Making use of this, $\lim_{i \rightarrow \infty} \|u(\cdot, t)\|_{L^{2^i}(\Omega)} = \|u(\cdot, t)\|_{L^\infty(\Omega)}$ and (2.2), by letting $i \rightarrow \infty$ in (2.31), we obtain the desired result. \square

We now can write our main theorem.

Theorem 7. *Let the initial values u_0 and v_0 are non-negative and satisfy:*

$$(u_0, v_0) \in (W^{1,r}(\Omega))^2 \quad \text{for some } r > n$$

and

$$0 < \|v_0\|_{L^\infty(\Omega)} \leq \delta_0,$$

which δ_0 is introduced in Lemma 3. Assume that the condition (2.10) holds. Then, the solution of the problem (1.1) with the functions $\gamma(v)$ and $\xi(v)$ defined by (1.3) is global and bounded.

Proof. By considering the extensibility criterion provided by Lemma 1, the proof is a consequence of (2.3) and Lemma 6. \square

Declaration of interests

The authors do not work for, advise, own shares in, or receive funds from any organization that could benefit from this article, and have declared no affiliations other than their research organizations.

References

- [1] J. Ahn and C. Yoon, “Global well-posedness and stability of constant equilibria in parabolic-elliptic chemotaxis systems without gradient sensing”, *Nonlinearity* **32** (2019), no. 4, pp. 1327–1351.
- [2] K. Baghaei, “Boundedness of classical solutions to a chemotaxis consumption system with signal dependent motility and logistic source”, *C. R. Math. Acad. Sci. Paris* **361** (2023), pp. 1641–1652.
- [3] K. Baghaei and A. Khelghati, “Boundedness of classical solutions for a chemotaxis model with consumption of chemoattractant”, *C. R. Math. Acad. Sci. Paris* **355** (2017), no. 6, pp. 633–639.
- [4] K. Baghaei and A. Khelghati, “Global existence and boundedness of classical solutions for a chemotaxis model with consumption of chemoattractant and logistic source”, *Math. Methods Appl. Sci.* **40** (2017), no. 10, pp. 3799–3807.
- [5] T. Black and M. Winkler, “Global weak solutions and absorbing sets in a chemotaxis–Navier–Stokes system with prescribed signal concentration on the boundary”, *Math. Models Methods Appl. Sci.* **32** (2022), no. 1, pp. 137–173.
- [6] M. Burger, P. Laurençot and A. Trescases, “Delayed blow-up for chemotaxis models with local sensing”, *J. Lond. Math. Soc.* **103** (2021), no. 4, pp. 1596–1617.
- [7] X. Cao and J. Lankeit, “Global classical small-data solutions for a three-dimensional chemotaxis Navier–Stokes system involving matrix-valued sensitivities”, *Calc. Var. Partial Differ. Equ.* **55** (2016), no. 4, article no. 107 (39 pages).
- [8] L. Desvillettes, Y.-J. Kim, A. Trescases and C. Yoon, “A logarithmic chemotaxis model featuring global existence and aggregation”, *Nonlinear Anal., Real World Appl.* **50** (2019), pp. 562–582.
- [9] K. Fujie and J. Jiang, “Boundedness of classical solutions to a degenerate Keller–Segel type model with signal-dependent motilities”, *Acta Appl. Math.* **176** (2021), article no. 3 (36 pages).

- [10] K. Fujie and T. Senba, “Global boundedness of solutions to a parabolic-parabolic chemotaxis system with local sensing in higher dimensions”, *Nonlinearity* **35** (2022), no. 7, pp. 3777–3811.
- [11] K. Fujie and J. Jiang, “Global existence for a kinetic model of pattern formation with density-suppressed motilities”, *J. Differ. Equations* **269** (2020), no. 6, pp. 5338–5378.
- [12] J. Han and C. Liu, “Global weak solution for a chemotaxis Navier–Stokes system with p -Laplacian diffusion and singular sensitivity”, *Nonlinear Anal., Real World Appl.* **73** (2023), article no. 103898 (24 pages).
- [13] H.-Y. Jin, Y.-J. Kim and Z.-A. Wang, “Boundedness, stabilization, and pattern formation driven by density-suppressed motility”, *SIAM J. Appl. Math.* **78** (2018), no. 3, pp. 1632–1657.
- [14] H.-Y. Jin and Z.-A. Wang, “Critical mass on the Keller–Segel system with signal-dependent motility”, *Proc. Am. Math. Soc.* **148** (2020), no. 11, pp. 4855–4873.
- [15] E. F. Keller and L. A. Segel, “Initiation of slime mold aggregation viewed as an instability”, *J. Theor. Biol.* **26** (1970), no. 3, pp. 399–415.
- [16] A. Khelghati and K. Baghaei, “Boundedness of classical solutions for a chemotaxis model with rotational flux terms”, *Z. Angew. Math. Mech.* **98** (2018), no. 10, pp. 1864–1877.
- [17] A. Khelghati and K. Baghaei, “Boundedness of classical solutions for a chemotaxis system with general sensitivity function”, *Appl. Anal.* **98** (2019), no. 3, pp. 611–621.
- [18] P. Laurençot, “Long term spatial homogeneity for a chemotaxis model with local sensing and consumption”, *Commun. Math. Sci.* **21** (2023), no. 6, pp. 1743–1750.
- [19] D. Li and J. Zhao, “Global boundedness and large time behavior of solutions to a chemotaxis-consumption system with signal-dependent motility”, *Z. Angew. Math. Phys.* **72** (2021), no. 2, article no. 57 (20 pages).
- [20] G. Li and M. Winkler, “Relaxation in a Keller–Segel-consumption system involving signal-dependent motilities”, *Commun. Math. Sci.* **21** (2023), no. 2, pp. 299–322.
- [21] G. Li and M. Winkler, “Refined regularity analysis for a Keller–Segel-consumption system involving signal-dependent motilities”, *Appl. Anal.* **103** (2024), no. 1, pp. 45–64.
- [22] X. Li, L. Wang and X. Pan, “Boundedness and stabilization in the chemotaxis consumption model with signal-dependent motility”, *Z. Angew. Math. Phys.* **72** (2021), no. 4, article no. 170 (18 pages).
- [23] Y. Li and S. Lu, “Global solutions to a chemotaxis-growth system with signal-dependent motilities and signal consumption”, *J. Math. Anal. Appl.* **521** (2023), no. 1, article no. 126902 (17 pages).
- [24] Y. Tao, “Boundedness in a chemotaxis model with oxygen consumption by bacteria”, *J. Math. Anal. Appl.* **381** (2011), no. 2, pp. 521–529.
- [25] Y. Tao and M. Winkler, “Eventual smoothness and stabilization of large-data solutions in a three-dimensional chemotaxis system with consumption of chemoattractant”, *J. Differ. Equations* **252** (2012), no. 3, pp. 2520–2543.
- [26] Y. Tao and M. Winkler, “Global solutions to a Keller–Segel-consumption system involving singularly signal-dependent motilities in domains of arbitrary dimension”, *J. Differ. Equations* **343** (2023), pp. 390–418.
- [27] I. Tuval, L. Cisneros, C. Dombrowski, C. W. Wolgemuth, J. O. Kessler and R. E. Goldstein, “Bacterial swimming and oxygen transport near contact lines”, *Proc. Natl. Acad. Sci. USA* **102** (2005), no. 7, pp. 2277–2282.
- [28] J. Wang and M. Wang, “Boundedness in the higher-dimensional Keller–Segel model with signal-dependent motility and logistic growth”, *J. Math. Phys.* **60** (2019), no. 1, article no. 011507 (14 pages).
- [29] L. Wang, “Global dynamics for a chemotaxis consumption system with signal-dependent motility and logistic source”, *J. Differ. Equations* **348** (2023), pp. 191–222.

- [30] M. Winkler, “Stabilization in a two-dimensional chemotaxis-Navier–Stokes system”, *Arch. Ration. Mech. Anal.* **211** (2014), no. 2, pp. 455–487.
- [31] M. Winkler, “A result on parabolic gradient regularity in Orlicz spaces and application to absorption-induced blow-up prevention in a Keller–Segel-type cross-diffusion system”, *Int. Math. Res. Not.* **2023** (2023), no. 19, pp. 16336–16393.
- [32] M. Winkler, “Global generalized solvability in a strongly degenerate taxis-type parabolic system modeling migration-consumption interaction”, *Z. Angew. Math. Phys.* **74** (2023), no. 1, article no. 32 (20 pages).
- [33] Y. Xiao and J. Jiang, “Global existence and uniform boundedness in a fully parabolic Keller–Segel system with non-monotonic signal-dependent motility”, *J. Differ. Equations* **354** (2023), pp. 403–429.
- [34] C. Yoon and Y.-J. Kim, “Global existence and aggregation in a Keller–Segel model with Fokker–Planck diffusion”, *Acta Appl. Math.* **149** (2017), pp. 101–123.