



ACADÉMIE
DES SCIENCES
INSTITUT DE FRANCE

Comptes Rendus

Mathématique

Luc Illusie

Grothendieck and differential calculus

Volume 363 (2025), p. 829-841

Online since: 15 July 2025

<https://doi.org/10.5802/crmath.764>



This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*The Comptes Rendus. Mathématique are a member of the
Mersenne Center for open scientific publishing*
www.centre-mersenne.org — e-ISSN : 1778-3569



Historical Commentary / *Commentaire historique*
Algebra / *Algèbre*

Grothendieck and differential calculus

Grothendieck et le calcul différentiel

Luc Illusie^a

^a Institut de Mathématique d'Orsay, Bât. 307, Université Paris-Saclay, 91405 Orsay Cédex, France
E-mail: luc.illusie@wanadoo.fr

Abstract. I discuss some of the contributions that Grothendieck brought to differential calculus in the 1960s (infinitesimal neighborhoods, algebraic de Rham cohomology, crystalline cohomology) and sketch a few recent developments.

Résumé. Je présente quelques-unes des contributions que Grothendieck a apportées au calcul différentiel dans les années 1960 (voisinages infinitésimaux, cohomologie de de Rham algébrique, cohomologie cristalline), et esquisse certains développements récents.

Keywords. Infinitesimal neighborhood, de Rham complex, de Rham cohomology, crystalline cohomology, p -adic Hodge theory, cotangent complex, prismatic cohomology.

Mots-clés. Voisinage infinitésimal, complexe de de Rham, cohomologie de de Rham, cohomologie cristalline, théorie de Hodge p -adique, complexe cotangent, cohomologie prismatique.

2020 Mathematics Subject Classification. 14F20, 14F30, 14F40.

Manuscript received 11 February 2025, revised 12 June 2025, accepted 6 June 2025.

Introduction

Roughly speaking, differential calculus consists in this: given a good space X , a good function f on X , and points x, y in X , evaluate $f(y) - f(x)$ when x and y are *infinitesimally* close. What does one mean by “*infinitesimally* close”? If one just says “close”, one would mean that the point (x, y) belongs to a neighborhood of the diagonal in $X \times X$. These neighborhoods lead to the notion of “uniform structure” in Bourbaki, *Topologie générale*. But what would be an *infinitesimal neighborhood* of the diagonal? Grothendieck was the first one to give a rigorous definition of this notion. He did it in the context of analytic geometry, in two remarkable exposés at the Séminaire Cartan 1960-61 [31], which brought a totally new perspective on sheaves of differential forms and differential calculus in general.

Let me explain his construction in a simple case. Let X be a smooth complex analytic variety, of dimension n . Consider the diagonal map $i: X \rightarrow X \times X$. This is a closed immersion, defined by a coherent sheaf of ideals I fitting in the exact sequence

$$0 \longrightarrow I \longrightarrow \mathcal{O}_{X \times X} \longrightarrow \mathcal{O}_X \longrightarrow 0. \quad (1)$$

In local coordinates (x_i) on X , (x_i, y_i) on $X \times X$, it is locally generated by $(y_i - x_i)$, $1 \leq i \leq n$. It coincides with $\mathcal{O}_{X \times X}$ outside the diagonal. Grothendieck considered the exact sequence deduced from (1) by dividing $\mathcal{O}_{X \times X}$ by I^2 :

$$0 \longrightarrow I/I^2 \longrightarrow \mathcal{O}_{X \times X}/I^2 \longrightarrow \mathcal{O}_X \longrightarrow 0. \quad (2)$$

He observed that:

- (a) there is a canonical¹ isomorphism

$$I/I^2 \xrightarrow{\sim} \Omega_X^1, \quad (3)$$

where Ω_X^1 is the sheaf of 1-forms on X , a vector bundle of rank n , with local basis (dx_i) ($1 \leq i \leq n$);

- (b) if p_1^*, p_2^* are the two sections of $\mathcal{O}_{X \times X}/I^2 \rightarrow \mathcal{O}_X$ induced by the projections $X \times X \rightarrow X$, then the map

$$\mathcal{O}_X \longrightarrow I/I^2, f \longmapsto p_2^* f - p_1^* f,$$

coincides via (3) with the derivation $d_X: \mathcal{O}_X \rightarrow \Omega_X^1$.

The verification is immediate: for (a), we have the exact sequence

$$0 \longrightarrow I/I^2 \xrightarrow{d_{X \times X}} i^* \Omega_{X \times X}^1 \longrightarrow \Omega_X^1 \longrightarrow 0,$$

where the surjection is the sum map: $\Omega_X^1 \oplus \Omega_X^1 \rightarrow \Omega_X^1$, so that we get an isomorphism (3) from I/I^2 to Ω_X^1 , injected into $\Omega_X^1 \oplus \Omega_X^1$ by $(-\text{Id}, \text{Id})$; (b) expresses that locally $y_i - x_i \pmod{I^2}$ corresponds to dx_i , I/I^2 is free with basis the classes of $y_i - x_i \pmod{I^2}$, and

$$f(y) - f(x) = \sum \partial f / \partial x_i(x) (y_i - x_i) \pmod{I^2}.$$

Grothendieck defined the *first infinitesimal neighborhood of the diagonal* as the closed analytic subspace $\Delta_X^1(1) \hookrightarrow X \times X$ defined by the ideal I^2 . It has the same underlying space as X , and its ring of functions, $\mathcal{O}_{X \times X}/I^2$, also called sheaf of *principal parts* of order 1 and denoted \mathcal{P}_X^1 , is by (2) an extension of \mathcal{O}_X by the ideal of square zero $I/I^2 = \Omega_X^1$. By replacing I^2 by higher powers of I , one gets higher infinitesimal neighborhoods $\Delta_X^r(1)$ of the diagonal, leading to differential calculus of higher order.

At the same time, Grothendieck realized that one could reverse the steam, and *define* the sheaf of differentials Ω^1 by the I/I^2 formula, even if X is a singular analytic space, and even in a relative situation $X \rightarrow S$, and that in this way one obtained a theory with good functoriality properties. He also saw that one could develop a similar theory in algebraic geometry, for schemes, which he wrote up a couple of years later in EGA IV (see [24, 0] and [26, IV]).

If $X \rightarrow S$ is a morphism of schemes, Grothendieck defines $\Omega_{X/S}^1$ as I/I^2 , where I is the ideal of the diagonal immersion $X \rightarrow X \times_S X$, and the map $d_{X/S}: \mathcal{O}_X \rightarrow \Omega_{X/S}^1$ by $p_2^* - p_1^*$, where, as above, p_1^*, p_2^* are the two sections of $\mathcal{O}_{X \times_S X}/I^2 \rightarrow \mathcal{O}_X$ given by the two projections. The sheaf $\Omega_{X/S}^1$ is quasi-coherent (locally free of finite type when X/S is smooth), and $d_{X/S}$ is an \mathcal{O}_S -derivation. When X and S are affine, so that $X \rightarrow S$ corresponds to a homomorphism of rings $A \rightarrow B$, then $\Omega_{X/S}^1$ is the quasi-coherent sheaf defined by $\Omega_{B/A}^1 := J/J^2$, where J is the kernel of the surjection $B \otimes_A B \rightarrow B$, $b_1 \otimes b_2 \mapsto b_1 b_2$, and $d_{X/S}$ corresponds to the A -derivation $d_{B/A}: B \rightarrow \Omega_{B/A}^1$, $d_{B/A}(b) = 1 \otimes b - b \otimes 1$. One checks that composition with $d_{B/A}$ defines a functorial isomorphism

$$\text{Hom}_B(\Omega_{B/A}^1, M) \xrightarrow{\sim} \text{Der}_A(B, M)$$

where $\text{Der}_A(B, M)$ is the B -module of A -derivations of B in a B -module M , so that $\Omega_{B/A}^1$ thus defined is the module of *Kähler differentials* of B/A .²

¹As the referee points out, though being called “canonical”, this isomorphism involves in its description a choice of signs (or order on the factors of $X \times X$).

²This module, defined by Kähler in 1953 [37], didn’t attract much attention until Grothendieck used it extensively, already in [29].

One checks that $d: \mathcal{O}_X \rightarrow \Omega_{X/S}^1$ uniquely extends to a complex

$$\Omega_{X/S}^\bullet = (\mathcal{O}_X \xrightarrow{d} \Omega_{X/S}^1 \longrightarrow \cdots \longrightarrow \Omega_{X/S}^i \xrightarrow{d} \Omega_{X/S}^{i+1} \longrightarrow \cdots),$$

called the *de Rham complex* of X/S , where $\Omega_{X/S}^i := \bigwedge_{\mathcal{O}_X}^i \Omega_{X/S}^1$, such that $d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^i \alpha \wedge d\beta$ for α of degree i .

Grothendieck defined the *algebraic de Rham cohomology* of X/S by

$$H_{\mathrm{dR}}^*(X/S) = H^*(X, \Omega_{X/S}^\bullet)$$

(hypercohomology of X for the Zariski topology with coefficients in $\Omega_{X/S}^\bullet$). This object enjoys nice functoriality properties. Its calculation led to rich developments.

Let k be a field. For X/k a standard projective space, a projective smooth curve, or an Abelian variety, the de Rham cohomology groups $H_{\mathrm{dR}}^i(X/k)$ are finite dimensional k -vector spaces, with the same dimension (and same structure) as the transcendental Betti cohomology groups when $k = \mathbb{C}$. But in general, algebraic de Rham cohomology behaves in quite different ways in characteristic zero and in characteristic $p > 0$.

1. Algebraic de Rham cohomology in characteristic zero

Let X be a smooth scheme over \mathbb{C} . The set $X(\mathbb{C})$ of its complex points underlies a smooth complex analytic variety X^{an} . By the *holomorphic Poincaré lemma*³ its de Rham complex $\Omega_{X^{\mathrm{an}}}^\bullet$ (where we omit $/\mathbb{C}$ for short) is a resolution of the constant sheaf \mathbb{C}_X : the sequence

$$0 \longrightarrow \mathbb{C}_X \longrightarrow \mathcal{O}_{X^{\mathrm{an}}} \longrightarrow \Omega_{X^{\mathrm{an}}}^1 \longrightarrow \cdots$$

is exact. In other words, the augmentation

$$\mathbb{C}_X \longrightarrow \Omega_{X^{\mathrm{an}}}^\bullet$$

is a quasi-isomorphism, which thus induces an isomorphism

$$H^*(X^{\mathrm{an}}, \mathbb{C}) \xrightarrow{\sim} H_{\mathrm{dR}}^*(X^{\mathrm{an}}), \quad (4)$$

by which a class $\omega \in H_{\mathrm{dR}}^n(X^{\mathrm{an}})$ corresponds to the homomorphism

$$\mathrm{Hom}(H_n(X^{\mathrm{an}}, \mathbb{Z}), \mathbb{C}), \gamma \mapsto \int_\gamma \omega$$

(with a suitable sign convention, see [18, I, Theorem 1.2, p. 14]).

On the algebraic side, the Poincaré lemma fails: $\Omega_X^\bullet := \Omega_{X/\mathbb{C}}^\bullet$ is far from being a resolution of \mathbb{C}_X (as is already seen for X the affine line $\mathrm{Spec}(\mathbb{C}[t])$: the form $dt/(1+t)$ near zero has no algebraic primitive). However, we have a canonical map of ringed spaces

$$X^{\mathrm{an}} \longrightarrow X, \quad (5)$$

inducing a map $\Omega_X^\bullet \rightarrow \Omega_{X^{\mathrm{an}}}^\bullet$ on de Rham complexes, hence on de Rham cohomology

$$H_{\mathrm{dR}}^*(X) \longrightarrow H_{\mathrm{dR}}^*(X^{\mathrm{an}}). \quad (6)$$

Grothendieck proved the following surprising theorem.

Theorem 1 ([25]). *The map (6) is an isomorphism, and thus induces an isomorphism $H_{\mathrm{dR}}^*(X) \xrightarrow{\sim} H^*(X^{\mathrm{an}}, \mathbb{C})$.*

³According to de Rham [48, p. 646], the C^∞ analogue of this lemma, namely the exactness of the augmented de Rham complex of global C^∞ -forms on \mathbb{R}^n , attributed to Poincaré, was in fact first proved by Volterra [53, pp. 407–422] in 1889.

In particular, $\dim_{\mathbb{C}} H_{\text{dR}}^i(X) = b_i$, where $b_i = \dim_{\mathbb{C}} H^i(X^{\text{an}}, \mathbb{C})$ is the transcendental i th Betti number, and these algebraic de Rham cohomology spaces are homotopy invariants!

In the proper case, the proof is easy. By Serre's GAGA theorems, the map

$$H^j(X, \Omega_X^i) \longrightarrow H^j(X^{\text{an}}, \Omega_{X^{\text{an}}}^i)$$

is an isomorphism, which implies the result by the Hodge to de Rham spectral sequence $E_1^{ij} = H^j(X, \Omega_X^i) \Rightarrow H_{\text{dR}}^{i+j}(X)$. However, in the non-proper case, the proof is more difficult. Grothendieck uses Hironaka's resolution of singularities [32] to compactify X by a divisor with normal crossings at infinity, and concludes by a local calculation due to Atiyah–Hodge. This calculation, which involves differential forms with logarithmic poles, will be revisited by Deligne in his work on mixed Hodge theory [15,16], and later will be one of the sources of the theory of logarithmic geometry.

Example. Let $X := \text{Spec}(\mathbb{C}[t, t^{-1}])$ be the punctured affine line. Then $H_{\text{dR}}^0(X) = \mathbb{C}$, $H_{\text{dR}}^1(X) = \mathbb{C} \cdot (dt/t)$, and $H_{\text{dR}}^n(X) = 0$ for $n > 1$. The class dt/t is a basis of $H^1(X^{\text{an}}, \mathbb{C})$, sending the class of the circle in $H_1(X^{\text{an}}, \mathbb{Z})$ to $2\pi i$.

Let X_0 be a smooth scheme over \mathbb{Q} , and let $X = X_0 \otimes \mathbb{C}$. Since $H_{\text{dR}}^*(X_0/\mathbb{Q}) \otimes \mathbb{C} = H_{\text{dR}}^*(X/\mathbb{C})$, the isomorphism of Theorem 1 gives a *period isomorphism*

$$H_{\text{dR}}^*(X_0/\mathbb{Q}) \otimes \mathbb{C} \xrightarrow{\sim} H^*(X^{\text{an}}, \mathbb{Q}) \otimes \mathbb{C}. \quad (7)$$

As $H^*(X^{\text{an}}, \mathbb{Q}) = \text{Hom}(H_*(X^{\text{an}}, \mathbb{Z}), \mathbb{Q})$, the image by (7) of a class $\omega \in H_{\text{dR}}^n(X_0/\mathbb{Q})$ corresponds to the homomorphism sending $\gamma \in H_n(X^{\text{an}}, \mathbb{Z})$ to $\int_{\gamma} \omega \in \mathbb{C}$. These numbers are called *periods*. As the above example shows, they can be transcendental. Let $t(X)$ be the transcendence degree over \mathbb{Q} of the subfield of \mathbb{C} generated by those periods (for all n). Grothendieck asked for bounds for $t(X)$ and conjectured that $t(X)$ is the dimension of the image of the *motivic Galois group* of \mathbb{Q} into $\text{GL}(H^*(X^{\text{an}}, \mathbb{Q}))$ [3, 0.4]. See [3] for the state of the art on this conjecture and related ones in 2016.

For X/\mathbb{C} projective and smooth, X^{an} is Kähler, and the Hodge decomposition for $H^*(X^{\text{an}}, \mathbb{C})$ gives an isomorphism

$$H_{\text{dR}}^n(X) \xrightarrow{\sim} \bigoplus_{i+j=n} H^{ij}, \quad (8)$$

with $H^{ij} := H^j(X, \Omega_X^i)$ and $H^{ji} = \overline{H^{ij}}$. In particular, if $h^n := \dim H_{\text{dR}}^n(X)$, $h^{ij} := \dim H^{ij}(X, \Omega_X^i)$, we have

$$h^n = \sum_{i+j=n} h^{ij} \quad (9)$$

for all n , and the Hodge to de Rham spectral sequence degenerates at the first page. Moreover, we have the *Hodge symmetry* $h^{ij} = h^{ji}$. It was proved by Deligne [14, Proposition 5.3] that (8) (hence (9) and Hodge symmetry) holds more generally for X/\mathbb{C} only proper and smooth.

2. Algebraic de Rham cohomology in characteristic $p > 0$

Fix a prime number p .

If X is a scheme of characteristic p (i.e., a scheme over \mathbb{F}_p), X admits a natural endomorphism F_X , called the *Frobenius endomorphism*, which is the identity on the underlying space and is $a \mapsto a^p$ on \mathcal{O}_X . If X is a scheme over S , and S is of characteristic p , then, for a in \mathcal{O}_X , $d(a^p) = pa^{p-1}da = 0$ (where $d := d_{X/S}$), and for $\omega \in \Omega_{X/S}^i$, $d(a^p\omega) = a^p d\omega$, so that the de Rham complex $\Omega_{X/S}^\bullet$ is not just \mathcal{O}_S -linear, but also p -linear with respect to \mathcal{O}_X . The conjunction of \mathcal{O}_S -linearity and p - \mathcal{O}_X linearity is best expressed by introducing the pull-back $X^{(1)}/S$ of X by $F_S: S \rightarrow S$, and the S -morphism $F: X \rightarrow X^{(1)}$, called the *relative Frobenius*, through which F_X

canonically factors, and observing that $F_*\Omega_{X/S}^\bullet$, i.e., $\Omega_{X/S}^\bullet$, considered as a complex of $\mathcal{O}_{X^{(1)}}$ -modules, has an $\mathcal{O}_{X^{(1)}}$ -linear differential. In particular, the cohomology sheaves $\mathcal{H}^i(F_*\Omega_{X/S}^\bullet)$ are $\mathcal{O}_{X^{(1)}}$ -modules.

These are trivial remarks. But in 1957 Cartier discovered a new operation on differential forms, which was to later control all differential calculus in positive characteristic (and, consequently, in mixed characteristic, too, for a large part). In a presentation due to Deligne and Katz [38, 7.2], his result is the following.

Theorem 2 ([13]). *The maps*

$$\begin{aligned}\mathcal{O}_X &\longrightarrow \mathcal{O}_X, a \longmapsto a^p, \\ \Omega_{X/S}^1 &\longrightarrow \Omega_{X/S}^1, a df \longmapsto a^p f^{p-1} df\end{aligned}$$

induce $\mathcal{O}_{X^{(1)}}$ -linear homomorphisms

$$\begin{aligned}\mathcal{O}_{X^{(1)}} &\longrightarrow \mathcal{H}^0(F_*\Omega_{X/S}^\bullet), \\ \Omega_{X^{(1)}/S}^1 &\longrightarrow \mathcal{H}^1(F_*\Omega_{X/S}^\bullet),\end{aligned}$$

which uniquely extend to a homomorphism of graded $\mathcal{O}_{X^{(1)}}$ -algebras denoted⁴

$$C^{-1}: \bigoplus_i \Omega_{X^{(1)}/S}^i \longrightarrow \bigoplus_i \mathcal{H}^i(F_*\Omega_{X/S}^\bullet).$$

When X/S is smooth, C^{-1} is an isomorphism.

The proof of the first part is a small calculation (that has recently reappeared in the theory of δ -structures and prisms [12]), and for the second one, an easy dévissage reduces to $X = \operatorname{Spec}(\mathbb{F}_p[t])$, where writing $\mathbb{F}_p[t]$ as the free module over $\mathbb{F}_p[t^p]$ with basis $1, t, \dots, t^{p-1}$, one sees that \mathcal{H}^0 (resp. \mathcal{H}^1) has basis 1 (resp. $t^{p-1}dt$).

It follows from Theorem 2, that if $X = \operatorname{Spec}(A)$ is smooth, affine over a field k of characteristic p , then $H_{\text{dR}}^*(X/k)$ is isomorphic to $\bigoplus \Omega_{A^{(p)}/k}^i$, hence is of infinite dimension over k if X is of positive dimension, in contrast with the case $k = \mathbb{C}$. On the other hand, if X/k is proper and smooth, then the Hodge spaces $H^j(X, \Omega_{X/k}^i)$ have finite dimension h^{ij} , so that, by the Hodge to de Rham spectral sequence

$$E_1^{ij} = H^j(X, \Omega_{X/k}^i) \implies H_{\text{dR}}^{i+j}(X/k)$$

the de Rham cohomology spaces $H_{\text{dR}}^n(X/k)$ are also finite dimensional. However, in the late 1950s and early 1960s it was observed by Serre that Hodge symmetry could fail [50], and by Mumford [45] that a strict inequality $h^n < \sum_{i+j=n} h^{ij}$ was possible. Concerning this second point, a progress was made in 1987.

Theorem 3 ([17]). *Assume k perfect and let $W(k)$ be the ring of Witt vectors on k .⁵ Let X/k be proper and smooth. Assume that $\dim(X) \leq p$, and X is liftable to $W_2(k)$, i.e., there exists a proper, smooth scheme $\mathcal{X}/W_2(k)$ whose reduction $\mathcal{X} \otimes k$ on k is isomorphic to X . Then*

$$h^n(X) = \sum_{i+j=n} h^{ij}(X), \quad \text{for all } n. \quad (10)$$

This is in fact a corollary of a stronger fact, namely, that assuming only X/k smooth of dimension $\leq p$ and liftable to $W_2(k)$, the de Rham complex $F_*\Omega_{X/k}^\bullet$, as an object of the derived category $D(X^{(1)}, \mathcal{O}_{X^{(1)}})$, is the sum of its shifted cohomology sheaves, i.e., thanks to the Cartier isomorphism of Theorem 2, we have an isomorphism in $D(X^{(1)}, \mathcal{O}_{X^{(1)}})$:

$$\bigoplus_i \Omega_{X^{(1)}/k}^i[-i] \xrightarrow{\sim} F_*\Omega_{X/k}^\bullet.$$

⁴For historical reasons: Cartier first defined $C: Z^i F_*\Omega_{X/S}^\bullet \rightarrow \Omega_{X^{(p)}/S}^i$.

⁵ $W(k) = \mathbb{Z}_p$ if $k = \mathbb{F}_p$ and $W_2(k) = W(k)/p^2 W(k)$.

A corollary of Theorem 3 was an algebraic proof of the Hodge degeneration (9). However, there remained the following question: does there exist X/k , proper, smooth of dimension $p+1$, liftable to $W_2(k)$, for which (10) fails for some n ? A positive answer, in a stronger form, was recently given by A. Petrov.

Theorem 4 ([46]). *There exists a projective, smooth scheme $\mathcal{X}/W(k)$ of relative dimension $p+1$, such that its reduction $X := \mathcal{X} \otimes k$ has*

$$h^p < \sum_{i+j=p} h^{ij}.$$

The proof is a tour de force. The identification and description of the obstruction to the degeneration at E_1 of the Hodge to de Rham spectral sequence, or rather, equivalently, of the degeneration at E_2 of the *conjugate spectral sequence* (coming from the canonical filtration of $F_*\Omega_{X/k}^\bullet$) uses the theory of *diffracted Hodge complexes* of Bhatt–Lurie–Drinfeld [10]. The core of the proof relies on an extremely ingenious description of extensions in de Rham and Hodge cohomology complexes of Abelian schemes over $W(k)$ endowed with group actions. See Section 4 for the definition of \mathcal{X} .

Let $\delta_p = \sum_{i+j=p} h_{ij} - h^p$. Petrov can show that, for his example, $\delta_p \geq 2$. He hopes that one could get $\delta_p = 1$ and then that δ_p could probably take any positive integer value.

3. Mixed characteristic: crystalline cohomology

The discovery by Grothendieck [28] (and, independently, by Katz–Oda [40]) of the algebraicity of the classical, analytic, *Gauss–Manin connection*, together with results of Monsky–Washnitzer on de Rham cohomology in the affine case [44], led him to conjecture (in [28]) the following.

Let k be a perfect field of characteristic $p > 0$. Let X/k be proper, smooth, and suppose it admits a proper, smooth lifting $\mathcal{X}/W(k)$. Then:

- (a) $H_{\text{dR}}^*(\mathcal{X}/W(k))$ does not depend on the lifting \mathcal{X} , in the sense that the H_{dR}^* of the various liftings are related by a transitive system of isomorphisms;
- (b) for $n \geq 1$, one can define an inductive system of ringed sites

$$(X/W_n(k))_{\text{crys}}, \quad (11)$$

called the *crystalline sites*, giving a compatible system of isomorphisms

$$H^*((X/W_n(k))_{\text{crys}}, \mathcal{O}) \xrightarrow{\sim} H_{\text{dR}}^*(X_n/W_n(k)), \quad (12)$$

for $X_n := \mathcal{X} \otimes W_n(k)$, yielding an isomorphism

$$H^*((X/W(k))_{\text{crys}}, \mathcal{O}) := \varprojlim H^*((X/W_n(k))_{\text{crys}}, \mathcal{O}) \xrightarrow{\sim} H_{\text{dR}}^*(\mathcal{X}/W(k)). \quad (13)$$

Berthelot proved these conjectures in his thesis [6], where he developed the theory of crystalline sites, topoi, and crystalline cohomology in a much more general framework.

In the case of interest to us, the crystalline site $(X/W_n(k))_{\text{crys}}$ is defined independently of the lifting $\mathcal{X}/W(k)$, which may not exist (see Serre’s examples in [36, 8.6]), and, in fact, for *any* scheme X/k . It is the category of $W_n(k)$ -thickenings $U \hookrightarrow T$ of open subschemes of X , together with a DP (= divided power structure) $x \mapsto x^{[r]}$ on the ideal of T ,⁶ compatible with $p^{[r]} = p^r/r!$, with families $(U_i \hookrightarrow T_i)$ covering $U \hookrightarrow T$ if (T_i) Zariski covers T . In contrast with the étale site, this site is far from the geometric intuition. In particular, it has no final object. If X/k is smooth and $Y/W_n(k)$ is a smooth lifting, then $X \hookrightarrow Y$ covers the final object of the associated topos, and

⁶See [51, Divided powers (07GK)].

cohomology can be calculated by a certain Čech complex, called the *Čech–Alexander complex*. For $n = 1$ and $Y = X$, this complex is the cosimplicial ring

$$\mathcal{O}_{\widetilde{X}^\bullet} \quad (14)$$

formed by the rings of the divided power envelopes $X, \widetilde{X \times_k X}, \widetilde{X \times_k X \times_k X}, \dots$ of the diagonal in the successive powers of X/k . A crucial point is the existence of an isomorphism:

$$\mathcal{O}_{\widetilde{X}^\bullet} \xrightarrow{\sim} \Omega_{X/k}^\bullet \quad (15)$$

in the derived category $D(X, k)$, compatible with the ring structures. This isomorphism is deep. It is a manifestation of Grothendieck's extraordinary insight into differential calculus. Berthelot's proof in [6, V, Théorème 2.3.2], suggested by Grothendieck in [28, 6.4, 6.5], using linearization of differential operators, is rather indirect. It is repeated in [7, 6.14, 7.2]. The construction of an explicit morphism of complexes from (14) to $\Omega_{X/k}^\bullet$ inducing an isomorphism in $D(X, k)$ after divided power completion of the left-hand side is given in [34, VIII, 1.4.4].

All proofs of (15) rely, in one way or another, on the following (trivial) observation.

Lemma 5 (DP Poincaré lemma). *Let A be any commutative ring, let $A\langle t \rangle$ be the divided power algebra $\Gamma_A(At)$ on the free module with basis t , and let $A\langle t \rangle \xrightarrow{d} A\langle t \rangle dt$ be the differential graded algebra with $d(t^{[n]}) = t^{[n-1]} dt$. Then the sequence*

$$0 \longrightarrow A \longrightarrow A\langle t \rangle \xrightarrow{d} A\langle t \rangle dt \longrightarrow 0$$

of A -modules is exact.

The algebra $k\langle t \rangle$ has a natural comultiplication, which makes it the Hopf algebra of a commutative unipotent group scheme⁷ \mathbf{G}_a^\sharp (over k). The morphism of group schemes $\mathbf{G}_a^\sharp \rightarrow \mathbf{G}_a$ given by $k[t] \rightarrow k\langle t \rangle$ defines a ring stack

$$\mathbf{G}_a^{\mathrm{dR}} := [\mathbf{G}_a^\sharp \rightarrow \mathbf{G}_a],$$

which in turn, by Bhatt–Lurie's transmutation [9, 2.3.8], yields a stack

$$X^{\mathrm{dR}}: R \longmapsto X(\mathbf{G}_a^{\mathrm{dR}}(R)),$$

a gerbe on $X^{(1)}$ banded by the DP-envelope $T_{X^{(1)}}^\sharp$ of the tangent sheaf of $X^{(1)}/k$. We have

$$H^*(X^{\mathrm{dR}}, \mathcal{O}) \xrightarrow{\sim} H_{\mathrm{dR}}^*(X/k).$$

The cosimplicial ring (14) can be re-interpreted as the ring of the Čech simplicial complex associated with the cover $X \rightarrow X^{\mathrm{dR}}$ (transmuted from the cover $\mathbf{G}_a \rightarrow \mathbf{G}_a^{\mathrm{dR}}$). This viewpoint leads to a new proof of (15) [9, 2.5.6]. A finer result (taking into account the conjugate filtration of $F_*\Omega_{X/k}^\bullet$) is given in [9, 2.7.2 (3)].

In his famous letter to Tate dated May, 1966, Grothendieck said that the characteristic properties of a crystal were rigidity and ability to grow in an appropriate neighborhood. Crystals in \mathcal{O} -modules on the crystalline site $(X/W_n(k))_{\mathrm{crys}}$ are defined as sheaves of \mathcal{O} -modules for which the transition maps from one thickening to another are isomorphisms [6, IV, 1.1]. A typical example is *de Rham cohomology of liftings* X_n of X/k to $W_n(k)$, incarnating the crystal $R\Gamma((X/W_n(k))_{\mathrm{crys}}, \mathcal{O})$ on $(\mathrm{Spec}(k)/\mathrm{Spec}(W_n(k)))_{\mathrm{crys}}$: vertical rigidity, horizontal growth:

$$\begin{array}{ccc} R\Gamma_{\mathrm{crys}}(X/k) & \longleftarrow & R\Gamma_{\mathrm{crys}}(X/W_n(k)) \\ \downarrow = & & \downarrow = \\ R\Gamma_{\mathrm{dR}}(X/k) & \longleftarrow & R\Gamma_{\mathrm{dR}}(X_n/W_n(k)). \end{array}$$

⁷In the sense of [19, II, 9] (\mathbf{G}_a^\sharp is not of finite type).

A stronger crystalline property of this cohomology was recently proven by S. Mondal [42, Theorem 5.0.1]: $R\Gamma_{\text{crys}}(X/W_n(k))$ is the unique (functorial in X) multiplicative lift of $R\Gamma_{\text{dR}}(X/k)$ to $W_n(k)$.

Crystalline cohomology has been intensely studied during the past 50 years and has been the subject of numerous reviews. I will just briefly recall a few main facts and sketch a couple of recent developments.

Let $K := \text{Frac}(W(k))$. For X/k proper and smooth, $H_{\text{crys}}^*(X/W(k))$ is a finite anti-commutative $W(k)$ -algebra ($R\Gamma_{\text{crys}}(X/W(k))$ in $D(W(k))$ is a perfect complex, and a derived commutative algebra), and

$$X \longrightarrow H_{\text{crys}}^*(X/W(k)) \otimes K$$

is a Weil cohomology (satisfying Künneth, Poincaré duality, and having a cycle class theory), with *good Betti numbers*: by Katz–Messing [39]

$$(b_i)_{\text{crys}}(X) := \dim_K H_{\text{crys}}^i(X/W(k)) \otimes K = \dim_{\mathbb{Q}_\ell} H^i(X_{\bar{k}}, \mathbb{Q}_\ell)$$

($\ell \neq p$), and if X lifts to characteristic zero, then $(b_i)_{\text{crys}}(X)$ is the i th Betti number of the resulting complex variety over \mathbb{C} .

A distinctive feature of crystalline cohomology is its *Frobenius lift*. The endomorphism F_X of X induces an endomorphism φ of $H_{\text{crys}}^*(X/W(k))$ which is semi-linear with respect to the automorphism σ of $W(k)$ deduced from the Frobenius automorphism of k . This endomorphism φ is an *isogeny*, i.e., $\varphi \otimes \mathbb{Q}_p$ is an isomorphism: more precisely, if X/k is of dimension d , there exists a σ^{-1} -linear endomorphism ν of $H_{\text{crys}}^*(X/W(k))$ such that $\varphi\nu = \nu\varphi = p^d$. By [39] again, for X/k proper and smooth, and $k = \mathbb{F}_q$, with $q = p^a$, then $\det(1 - \varphi^a t, H_{\text{crys}}^i(X/W(k)))$ belongs to $\mathbb{Z}[t]$ and coincides with the ℓ -adic analogue $\det(1 - F_q t, H^i(X_{\bar{k}}, \mathbb{Q}_\ell))$ for $\ell \neq p$.

Example. Let A/k be an Abelian variety of dimension g . Then

$$H_{\text{crys}}^*(A/W(k)) = \bigwedge^* H_{\text{crys}}^1(A/W(k)).$$

The $W(k)$ -module $H_{\text{crys}}^1(A/W(k))$ is free of rank $2g$. Equipped with $F := \varphi$ and $V := pF^{-1}$, it is canonically isomorphic to the Dieudonné module $M(A[p^\infty])$ of the p -divisible group associated with A . This result, due to Grothendieck (see [41, p. VI]), was the starting point of the so-called *crystalline Dieudonné theory*, developed by Grothendieck, Messing, Mazur–Messing, Berthelot–Breen–Messing in the 1970s and early 1980s, and recently revived in the prismatic context by several people [4, 23, 43, 49].

Note that there exists an Abelian scheme $\mathcal{A}/W(k)$ lifting A , and we have a canonical isomorphism

$$H_{\text{crys}}^*(A/W(k)) \xrightarrow{\sim} H_{\text{dR}}^*(\mathcal{A}/W(k)).$$

Though the Frobenius endomorphism of A rarely lifts to \mathcal{A} ,⁸ it does produce an endomorphism φ of the de Rham cohomology of $\mathcal{A}/W(k)$.

Together with its isogeny φ , the $W(k)$ -module $H_{\text{crys}}^i(X/W(k))$ becomes an F -crystal. As such, it has a *Dieudonné–Manin slope decomposition*

$$H_{\text{crys}}^i(X/W(k)) \otimes \mathbb{Q}_p = \bigoplus_{\lambda \in \mathbb{Q}} (H_{\text{crys}}^i(X/W(k)) \otimes \mathbb{Q}_p)_\lambda,$$

where for $\lambda = s/r$ in canonical form, $(H_{\text{crys}}^i(X/W(k)) \otimes \mathbb{Q}_p)_\lambda$ is the part of *slope* λ , which over $W(\bar{k}) \otimes \mathbb{Q}_p$ becomes isomorphic to a sum of copies of $(W(\bar{k})_\sigma[F]/(F^r - p^s)) \otimes \mathbb{Q}_p$ (see [19, IV, 3]). The relation between the *Newton polygon* associated with this decomposition and the Hodge numbers of X/k , and their variation in families (starting with Grothendieck’s observation that the Newton polygon rises under specialization), have been the subject of a great number of

⁸It does so only when A is ordinary and \mathcal{A} is its so-called *canonical lifting*.

studies in the 1970s and 1980s, by Katz, Mazur, Ogus, Oort, and many others. The theory of the *de Rham–Witt complex* [35] gave some geometric insight into these questions, which are still actively pursued today with the new technology provided by the Drinfeld–Bhatt–Lurie prismatic stacks [10,20] and the theory of prismatic F -gauges [9].

But undoubtedly the most important development of crystalline cohomology was the study of its relation with p -adic étale cohomology, which became known as *p -adic Hodge theory*.

The starting point is this. Let $\mathcal{X}/W(k)$ be proper and smooth, and let $X := \mathcal{X} \otimes k$, $\mathcal{X}_K := \mathcal{X} \otimes K$. Two different kinds of cohomology groups are associated with \mathcal{X}_K :

- (a) $H_{\mathrm{dR}}^i(\mathcal{X}_K/K)$, canonically isomorphic to $H_{\mathrm{crys}}^i(X/W(k)) \otimes K$, which is a finite dimensional K -vector space, of dimension b_i , equipped with the automorphism φ , and the *Hodge filtration* Fil^\bullet coming from the (degenerate) Hodge to de Rham spectral sequence;
- (b) $H^i(\mathcal{X}_{\bar{K}}, \mathbb{Q}_p)$, a finite dimensional \mathbb{Q}_p -vector space, of the same dimension b_i , equipped with the continuous action of the Galois group $G_K := \mathrm{Gal}(\bar{K}/K)$.

Grothendieck observed that, for \mathcal{X} an Abelian scheme and $i = 1$, both (a) and (b) characterize \mathcal{X} up to isogeny,⁹ and that led him to (boldly) conjecture that in general (a) and (b) should be related by *mysterious functors*. A solution was proposed about ten years later by Fontaine, in the form of a conjectural *period isomorphism* ([21], [22, III, 6.1.4]),¹⁰

$$B_{\mathrm{crys}} \otimes_K H_{\mathrm{dR}}^*(\mathcal{X}_K/K) \xrightarrow{\sim} B_{\mathrm{crys}} \otimes_{\mathbb{Q}_p} H^*(\mathcal{X}_{\bar{K}}, \mathbb{Q}_p), \quad (16)$$

where B_{crys} is a big K -algebra, equipped with the three types of structure $(\varphi, \mathrm{Fil}^\bullet, G_K\text{-action})$. This isomorphism (16) should be compatible with the three structures, and $H_{\mathrm{dR}}^*(\mathcal{X}_K/K)$ (resp. $H^*(\mathcal{X}_{\bar{K}}, \mathbb{Q}_p)$) should be recovered from the left- (resp. right-) hand side by a simple algebraic operation. After several special cases were treated by Fontaine, Fontaine–Messing, Kato, (16) was first proved in general by Tsuji [52] (together with a variant involving semistable reduction). Different proofs were provided later by Faltings, Niziol, Beilinson.

This was not the end of the story. Indeed, (16) is an isomorphism over \mathbb{Q}_p , and Grothendieck was dreaming of a comparison between $H_{\mathrm{dR}}^*(\mathcal{X}/W(k))$ and $H^*(\mathcal{X}_{\bar{K}}, \mathbb{Z}_p)$ that would not neglect torsion. He even asked whether one could compare the lengths of the corresponding torsion submodules. This problem was recently solved by Bhatt–Morrow–Scholze and Bhatt–Scholze, by the theory of *integral p -adic Hodge theory* [11] and *prismatic cohomology* [12]. These new theories somehow interpolate between the two sides of (16). Concerning torsion, the result is simple: one has

$$\mathrm{lg}_{W(k)} H_{\mathrm{dR}}^*(\mathcal{X}/W(k))_{\mathrm{tors}} \geq \mathrm{lg}_{\mathbb{Z}_p} H^*(\mathcal{X}_{\bar{K}}, \mathbb{Z}_p)_{\mathrm{tors}}. \quad (17)$$

The inequality can be strict, and in case of equality, the structures of elementary divisors can be different [11, 2.10, 2.11].

For non-smooth schemes, algebraic de Rham cohomology does not behave well. It was already noted by Grothendieck in EGA IV [24, 0_{IV}, 20.6.26] that for an A -algebra B , written as a quotient of a polynomial algebra P by an ideal J , the complex (in degrees -1 and 0)

$$J/J^2 \xrightarrow{d_{P/A}} B \otimes_P \Omega_{P/A}^1$$

does not depend, up to homotopy, on the choice of the surjective homomorphism $P \rightarrow B$, and was a finer invariant than its $H^0 = \Omega_{B/A}^1$ and H^{-1} (called *imperfection module* in loc. cit.). This is the first appearance of the *cotangent complex* $L_{B/A}$, in the form of its canonical truncation in degree ≥ -1 . Grothendieck returned to this in [30, 5.8], and suggested the use of this (truncated)

⁹In the sense that the contravariant functor from the category of Abelian schemes $\mathcal{X}/W(k)$ lifting X/k , up to isogeny, to the category of data (a) $(H_{\mathrm{dR}}^1(\mathcal{X}_K/K), \varphi, \mathrm{Fil}^1)$ (resp. (b) $(H^1(\mathcal{X}_{\bar{K}}, \mathbb{Q}_p), G_K\text{-action})$) is fully faithful, as follows from results of Serre–Tate, Grothendieck–Messing, Fontaine, Tate.

¹⁰See Note III 4, p. 402 of the second edition of [22] (2020) for remarks on the history of the proof.

complex in the formulation and proof of the Riemann–Roch theorem for locally complete intersection morphisms in SGA 6 [1, Exposé VIII]. He globalized it in [27]. A non-truncated version, in the affine case, was constructed independently by M. André [2] and D. Quillen [47]. Globalization on ringed toposes was made in [33,34] (which contain applications to deformation theory suggested by Grothendieck). Since then, the theory of the cotangent complex has undergone many developments. Let me just mention that a natural extension of it was the construction of *derived de Rham cohomology*. Introduced in [34, VIII], it has been extensively studied since 2011, starting with seminal papers by B. Bhatt [8] and A. Beilinson [5].

4. Petrov’s example

Choose an elliptic curve E over $W(k)$ with supersingular reduction $E_k := E \otimes k$ (i.e.¹¹ $E_k[p](\bar{k}) = 0$, or equivalently, the Frobenius endomorphism is zero on $H^1(E_k, \mathcal{O})$). Then $E[p]$ is a finite, flat, commutative \mathbb{F}_p -module scheme over $W(k)$. Let $q = p^2$ and consider $E[p] \otimes_{\mathbb{F}_p} \mathbb{F}_q^{\oplus p}$, which is a sum of $2p$ copies of $E[p]$. Let $\mathrm{SL}_p(\mathbb{F}_q)$ act on $E[p] \otimes_{\mathbb{F}_p} \mathbb{F}_q^{\oplus p}$ by the trivial action on $E[p]$ and its natural action on $\mathbb{F}_q^{\oplus p}$, and define

$$G := \mathrm{SL}_p(\mathbb{F}_q) \ltimes (E[p] \otimes_{\mathbb{F}_p} \mathbb{F}_q^{\oplus p}).$$

This is a (non-commutative) finite flat group scheme over $W(k)$. By the Godeaux–Serre–Raynaud approximation technique, one can choose a projective, smooth scheme \mathcal{X} over $W(k)$, equipped with a G -torsor P , such that the classifying map $\mathcal{X} \rightarrow BG$ defined by P induces, on the reductions $X = \mathcal{X}_k$ and BG_k on k , an injection $H^{p+1}(BG_k, \mathcal{O}) \hookrightarrow H^{p+1}(X, \mathcal{O})$.

Then, for X , we have $h^p < \sum_{i+j=p} h^{ij}$.

Acknowledgments

These notes are the write-up of colloquium talks given at Clermont-Ferrand, on November 19, 2024, and at the Institut de Mathématiques de Jussieu, on December 12, 2024. I warmly thank the organizers, Jérôme Dubois (Clermont-Ferrand), and Pierre Berger and Olivier Debarre (Jussieu) for their invitations.

I am very grateful to Kęstutis Česnavičius, Olivier Debarre, Hélène Esnault, Shubhodip Mondal, Alexander Petrov, Claire Voisin and Weizhe Zheng for a careful reading of a first draft of these notes and suggestions of corrections and adjustments. I warmly thank the referee for a meticulous examination of the manuscript and the suggestion of several corrections.

Declaration of interests

The author does not work for, advise, own shares in, or receive funds from any organization that could benefit from this article, and has declared no affiliations other than their research organizations.

¹¹As usual, $-[p]$ denotes the kernel of multiplication by p .

References

- [1] P. Berthelot, A. Grothendieck and L. Illusie (eds.), *Théorie des intersections et théorème de Riemann–Roch*, Lecture Notes in Mathematics, Springer, 1971, pp. xii+700. Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967 (SGA 6).
- [2] M. André, *Homologie des algèbres commutatives*, Grundlehren der Mathematischen Wissenschaften, Springer, 1974, pp. xv+341.
- [3] Y. André, “Groupes de Galois motiviques et périodes”, in *Séminaire Bourbaki. Vol. 2015–2016. Exposés 1104–1119*, Astérisque, vol. 390, Société Mathématique de France, 2017, pp. 1–26. Exposé no. 1104.
- [4] J. Anschütz and A.-C. Le Bras, “Prismatic Dieudonné theory”, *Forum Math. Pi* **11** (2023), article no. e2 (92 pages).
- [5] A. Beilinson, “ p -adic periods and derived de Rham cohomology”, *J. Am. Math. Soc.* **25** (2012), no. 3, pp. 715–738.
- [6] P. Berthelot, *Cohomologie cristalline des schémas de caractéristique $p > 0$* , Lecture Notes in Mathematics, Springer, 1974, p. 604.
- [7] P. Berthelot and A. Ogus, *Notes on crystalline cohomology*, Mathematical Notes (Princeton), Princeton University Press; University of Tokyo Press, 1978, pp. vi+243.
- [8] B. Bhatt, “ p -adic derived de Rham cohomology”, 2012. Online at <https://arxiv.org/abs/1204.6560>.
- [9] B. Bhatt, “Prismatic F -gauges”, 2022. Online at <https://www.math.ias.edu/~bhatt/teaching/mat549f22/lectures.pdf>. Notes for a course given at Princeton University.
- [10] B. Bhatt and J. Lurie, “The prismaticization of p -adic formal schemes”, 2022. Online at <https://arxiv.org/abs/2201.06124>.
- [11] B. Bhatt, M. Morrow and P. Scholze, “Integral p -adic Hodge theory”, *Publ. Math., Inst. Hautes Étud. Sci.* **128** (2018), pp. 219–397.
- [12] B. Bhatt and P. Scholze, “Prisms and prismatic cohomology”, *Ann. Math. (2)* **196** (2022), no. 3, pp. 1135–1275.
- [13] P. Cartier, “Une nouvelle opération sur les formes différentielles”, *C. R. Acad. Sci. Paris* **244** (1957), pp. 426–428.
- [14] P. Deligne, “Théorème de Lefschetz et critères de dégénérescence de suites spectrales”, *Publ. Math., Inst. Hautes Étud. Sci.* (1968), no. 35, pp. 107–126.
- [15] P. Deligne, “Théorie de Hodge. II”, *Publ. Math., Inst. Hautes Étud. Sci.* (1971), no. 40, pp. 5–57.
- [16] P. Deligne, “Théorie de Hodge. III”, *Publ. Math., Inst. Hautes Étud. Sci.* (1974), no. 44, pp. 5–77.
- [17] P. Deligne and L. Illusie, “Relèvements modulo p^2 et décomposition du complexe de de Rham”, *Invent. Math.* **89** (1987), no. 2, pp. 247–270.
- [18] P. Deligne, J. S. Milne, A. Ogus and K.-y. Shih, *Hodge cycles, motives, and Shimura varieties*, Lecture Notes in Mathematics, Springer, 1982, pp. ii+414.
- [19] M. Demazure, *Lectures on p -divisible groups*, Lecture Notes in Mathematics, Springer, 1972, pp. v+98.
- [20] V. Drinfeld, “Prismaticization”, *Sel. Math., New Ser.* **30** (2024), no. 3, article no. 49 (150 pages).
- [21] J.-M. Fontaine, “Sur certains types de représentations p -adiques du groupe de Galois d’un corps local; construction d’un anneau de Barsotti–Tate”, *Ann. Math. (2)* **115** (1982), no. 3, pp. 529–577.
- [22] J.-M. Fontaine, “Représentations p -adiques semi-stables”, in *Périodes p -adiques (Bures-sur-Yvette, 1988)*, Astérisque, vol. 223, Société Mathématique de France, 1994, pp. 113–184.

- [23] Z. Gardner and K. Madapusi, “An algebraicity conjecture of Drinfeld and the moduli of p -divisible groups”, 2025. Online at <https://arxiv.org/abs/2412.10226>.
- [24] A. Grothendieck, “Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. 0, I”, *Publ. Math., Inst. Hautes Étud. Sci.* (1964), no. 20, pp. 5–259.
- [25] A. Grothendieck, “On the de Rham cohomology of algebraic varieties”, *Publ. Math., Inst. Hautes Étud. Sci.* (1966), no. 29, pp. 95–103.
- [26] A. Grothendieck, “Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. IV”, *Publ. Math., Inst. Hautes Étud. Sci.* (1967), no. 32, pp. 5–361.
- [27] A. Grothendieck, *Catégories cofibrées additives et complexe cotangent relatif*, Lecture Notes in Mathematics, Springer, 1968, pp. ii+167.
- [28] A. Grothendieck, “Crystals and the de Rham cohomology of schemes”, in *Dix exposés sur la cohomologie des schémas*, Advanced Studies in Pure Mathematics, North-Holland, 1968, pp. 306–358.
- [29] A. Grothendieck, “Théorèmes de dualité pour les faisceaux algébriques cohérents”, in *Séminaire Bourbaki, Vol. 4*, Société Mathématique de France, 1995, pp. 169–193. Exposé no. 149.
- [30] A. Grothendieck, *Résidus et dualité—Prénos pour un “Séminaire Hartshorne”* (R. Hartshorne, ed.), Documents Mathématiques, Société Mathématique de France, 2024, pp. xxiv+165.
- [31] A. Grothendieck, “Techniques de construction en géométrie analytique. VII. Étude locale des morphismes ; éléments de calcul infinitésimal”, in *Familles d’espaces complexes et fondements de la géométrie analytique*, Séminaire Henri Cartan, vol. 13, 1960–1961, pp. 1–27.
- [32] H. Hironaka, “Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II”, *Ann. Math. (2)* **79** (1964), 109–203 and 205–326.
- [33] L. Illusie, *Complexe cotangent et déformations. I*, Lecture Notes in Mathematics, Springer, 1971, pp. xv+355.
- [34] L. Illusie, *Complexe cotangent et déformations. II*, Lecture Notes in Mathematics, Springer, 1972, pp. vii+304.
- [35] L. Illusie, “Complexe de de Rham–Witt et cohomologie cristalline”, *Ann. Sci. Éc. Norm. Supér. (4)* **12** (1979), no. 4, pp. 501–661.
- [36] L. Illusie, “Grothendieck’s existence theorem in formal geometry”, in *Fundamental algebraic geometry*, Mathematical Surveys and Monographs, American Mathematical Society, 2005, pp. 179–233.
- [37] E. Kähler, “Algebra und Differentialrechnung”, in *Bericht über die Mathematiker-Tagung in Berlin, Januar, 1953*, Deutscher Verlag der Wissenschaften, 1953, pp. 58–163.
- [38] N. M. Katz, “Nilpotent connections and the monodromy theorem: applications of a result of Turrittin”, *Publ. Math., Inst. Hautes Étud. Sci.* (1970), no. 39, pp. 175–232.
- [39] N. M. Katz and W. Messing, “Some consequences of the Riemann hypothesis for varieties over finite fields”, *Invent. Math.* **23** (1974), pp. 73–77.
- [40] N. M. Katz and T. Oda, “On the differentiation of de Rham cohomology classes with respect to parameters”, *J. Math. Kyoto Univ.* **8** (1968), pp. 199–213.
- [41] B. Mazur and W. Messing, *Universal extensions and one dimensional crystalline cohomology*, Lecture Notes in Mathematics, Springer, 1974, pp. vii+134.
- [42] S. Mondal, “ $\mathbb{G}_a^{\text{perf}}$ -modules and de Rham cohomology”, *Adv. Math.* **409** (2022), article no. 108691 (72 pages).
- [43] S. Mondal, “Dieudonné theory via cohomology classifying stacks II”, 2024. Online at <https://arxiv.org/abs/2405.12967v2>.

- [44] P. Monsky and G. Washnitzer, “Formal cohomology. I”, *Ann. Math. (2)* **88** (1968), pp. 181–217.
- [45] D. Mumford, “Pathologies of modular algebraic surfaces”, *Am. J. Math.* **83** (1961), pp. 339–342.
- [46] A. Petrov, “Non-decomposability of the de Rham complex and non-semisimplicity of the Sen operator”, 2023. Online at <https://arxiv.org/abs/2302.11389v1>.
- [47] D. Quillen, “On the (co-)homology of commutative rings”, in *Applications of Categorical Algebra (Proc. Sympos. Pure Math., Vol. XVII, New York, 1968)*, Proceedings of Symposia in Pure Mathematics, American Mathematical Society, 1970, pp. 65–87.
- [48] G. de Rham, *Œuvres mathématiques*, L’Enseignement Mathématique, 1981, p. 748.
- [49] P. Scholze and J. Weinstein, *Berkeley lectures on p -adic geometry*, Annals of Mathematics Studies, Princeton University Press, 2020, pp. x+250.
- [50] J.-P. Serre, “Sur la topologie des variétés algébriques en caractéristique p ”, in *Symposium internacional de topología algebraica*, Universidad Nacional Autónoma de México and UNESCO, México, 1958, pp. 24–53.
- [51] The Stacks Project Authors, *Stacks Project*. Online at <http://stacks.math.columbia.edu>.
- [52] T. Tsuji, “ p -adic étale cohomology and crystalline cohomology in the semi-stable reduction case”, *Invent. Math.* **137** (1999), no. 2, pp. 233–411.
- [53] V. Volterra, *Opere matematiche. Memorie e note. Vol. I. 1881–1892*, Accademia Nazionale dei Lincei, 1954, pp. xxxiii+604.